




# Prompt photon production with two jets in POWHEG

Tomas Jezo, Michael Klasen, [Alexander Neuwirth](#)



# Photons

- Many sources of photons in proton-proton and heavy ion collisions:




**Direct-hard:** directly produced in the hard collisions, computed in perturbative non-thermal QCD.

**Thermal:** are expected to be produced in quark gluon plasma


$$dN_{\gamma}^{\text{thermal}} = dN_{\gamma}^{\text{direct}} - dN_{\gamma}^{\text{prompt}}$$

# Prompt photons




$$d\sigma = \sum_{a,b,c} \int_0^1 dx_a dx_b dz \underbrace{f_{a/A}(x_a, \mu_F) f_{b/B}(x_b, \mu_F)}_{\text{PDFs}} \cdot \underbrace{d\hat{\sigma}_{cX}(x_a P_A, x_b P_B, \frac{P_\gamma}{z}, \mu_F)}_{\text{partonic cross section}} \underbrace{D_{\gamma/c}(z, \mu_F)}_{\text{fragmentation}}$$

- ▶ Probe low- $x \sim \frac{2p_T^\gamma}{\sqrt{s}} \exp(-y)$  and  $Q \sim p_T^\gamma$  region where gluon density is large
- ▶ Scale and nuclear PDF uncertainties large at low  $p_T$
- ▶ Real photons are important probes of the QGP (e.g.  $T_{eff}$ .)
- ▶ Fragmentation contributions can be reduced by isolation cones



# POsitive Weight Hardest Emission Generator (POWHEG)



- ▶ Separation between NLO and fragmentation contribution is non-trivial in NLO+PS
- ▶ In order to regulate the divergences we must include pure QCD di-/trijet production at LO


Born	$pp \rightarrow \gamma j \sim \mathcal{O}(\alpha\alpha_s)$		$pp \rightarrow jj \sim \mathcal{O}(\alpha_s^2)$	
Virtual	$\mathcal{O}(\alpha\alpha_s) \cdot \mathcal{O}(\alpha)$	$\mathcal{O}(\alpha\alpha_s) \cdot \mathcal{O}(\alpha_s)$	$\mathcal{O}(\alpha_s^2) \cdot \mathcal{O}(\alpha)$	$\mathcal{O}(\alpha_s^2) \cdot \mathcal{O}(\alpha_s)$
Real	$pp \rightarrow \gamma\gamma j \sim \mathcal{O}(\alpha^2\alpha_s)$	$pp \rightarrow \gamma jj \sim \mathcal{O}(\alpha\alpha_s^2)$		$pp \rightarrow jjj \sim \mathcal{O}(\alpha_s^3)$

## Enhanced QED radiation

$$\Delta_R(p_T) \sim \exp \left[ - \int d\Phi_R \frac{R(\Phi_B, \Phi_R)}{B(\Phi_B)} \theta(k_T(\Phi_B, \Phi_R) - p_T) \right]$$


- ▶ The Born processes of different coupling strength rarely result in direct photon events  $p + p \rightarrow \gamma + j(+j)$ , compared to pure QCD events  $p + p \rightarrow j + j(+j)$ .
- ▶ If no QED emission is attached to the latter, it could still also produce a photon that is not directly produced in the hard process, but in the subsequent shower.

“enhancedrad”  $c$  feature increases the probability of attaching a QED radiation to the pure QCD Born process ( $\sim$  PYTHIA’s “EnhancedSplittings”).



# Photon and jets observables


- Isolated-photon plus two-jet production in  $pp$  collisions at  $\sqrt{s} = 13$  TeV and POWHEG ( $pp \rightarrow \gamma j$ ) + Pythia8 and POWHEG ( $pp \rightarrow \gamma jj$ ) + Pythia8 predictions ( $\Delta R = 0.4$  and  $E_{T,\text{cut}}^{\text{iso}} \equiv 0.0042 \cdot E_T^\gamma + 4.8$  GeV)<sup>1</sup>.



<sup>1</sup>Aad, G. *et al.* *JHEP* **03**, 179. arXiv: 1912.09866 [hep-ex] (2020), Badger, S. *et al.* arXiv: 2304.06682 [hep-ph] (Apr. 2023).

# Outlook

- ▶ Many  $\pi^0$  decays from jets into even more photons.
- ▶ Parton shower valid instead of fragmentation functions
- ▶ Comparison of the different isolation criteria and QCD/QED PS
- ▶ Ingredient for MiNLO(') photon+jets in POWHEG
- ▶ Soft photons in FOrward CALorimeter in ALICE (after LS3) and ALICE3



Thank you!







# Photon Fragmentation

Measured:


- ▶ Prompt photons in  $e^+e^-$  collisions (limited data)
- ▶ Prompt photons in hadronic collisions<sup>2</sup>
- ▶ In vector meson production, assuming dominating hadronic fluctuations of the photon at low scales (VMD)<sup>3</sup>:  $D_{\gamma,i}^{\text{had}}(z, \mu_s) = \sum_{V=\rho,\omega,\phi} \frac{4\pi\alpha}{f_V^2} D_{V/i}(z, \mu_s)$



<sup>2</sup>Adare, A. et al. *Phys. Rev. C* **87**, 054907. arXiv: 1208.1234 [nucl-ex] (2013).


<sup>3</sup>Klasen, M. & König, F. *Eur. Phys. J. C* **74**, 3009. arXiv: 1403.2290 [hep-ph] (2014).

# Photon isolation



Simple boost invariant definition of a fixed-cone (a):

- ▶ Radius around the photon is defined as  $R^{\text{iso}} = \sqrt{\Delta\phi^2 + \Delta\eta^2}$
- ▶ Activity in the cone is measured by the transverse momenta  $p_T$  of the particles
- ▶ Photon is isolated if the sum of the transverse momenta does not exceed a threshold  $E_T^{\text{max}} = p_T^{\text{iso}} \geq \sum_{i=1}^{\Delta R_i < R_{\text{max}}} p_T^i$
- ▶ Isolation suppresses fragmentation photons



<sup>4</sup>d'Enterria, D. & Rojo, J. *Nucl. Phys. B* **860**, 311–338. arXiv: 1202.1762 [hep-ph] (2012).

## Multi-channel phasespace construction

Same algorithm as in trijet<sup>5</sup>:

- ▶ Start from  $2 \rightarrow 2$  massless phase space
- ▶ With 3 massless final state particles there are 6 FSR and 3 ISR divergent regions ( $i, j \geq 3$ )

$$S_{0j}^{\text{ISR}} = S_{1j} + S_{2j} = \frac{1}{E_j^2(1 - \cos^2 \theta_{1j})}, \quad S_{ij}^{\text{FSR}} = \frac{E_i^2 + E_j^2}{2E_i^2 E_j^2(1 - \cos \theta_{ij})} \quad (1)$$



- ▶ POWHEG provides routines to construct  $N + 1$  phase space corresponding to either FSR/ISR divergent regions with good importance sampling ( $\Phi_{2 \rightarrow 3, kj}$ )
- ▶ Pick region randomly and suppress other regions

$$\tilde{S}_{0j} = \frac{S_{ij}}{\sum_j (S_{0j} + \sum_i S_{ij})}, \quad \tilde{S}_{ij} = \frac{S_{ij}}{\sum_j (S_{0j} + \sum_i S_{ij})} \frac{E_j}{E_i + E_j} \quad (2)$$

$$d\Phi_B = \sum_{kj} \tilde{S}_{kj} d\Phi_{2 \rightarrow 3, kj} \quad \tilde{S}_{0j, ij} \rightarrow \begin{cases} 1 & \text{as } E_j \rightarrow 0 \text{ or } \theta_{ij} \rightarrow 0 \\ 0 & \text{else} \end{cases} \quad (3)$$

<sup>5</sup>Kardos, A., Nason, P. & Oleari, C. *JHEP* **04**, 043. arXiv: 1402.4001 [hep-ph] (2014).

# Forward/Focal isolation




6

<sup>6</sup>ALICE Collaboration, C. Tech. rep. (CERN, Geneva, 2020). <https://cds.cern.ch/record/2719928>.

# Virtual comparison

Comparison of the virtuals from FormCalc, RECOLA2, OpenLoops2 and MG5\_aMC@NLO:

► x-axis random phase space point  $\times$  flavour structure



## Going to low scale

Using  $Q = p_T^\gamma$  becomes problematic at low  $p_T$ , some solutions are:

- ▶ Look at ratios since the scale uncertainties are correlated and therefore cancel out
- ▶ Fix the scale to some value  $Q \geq 1.2$  GeV
- ▶ Compute a ratio between  $f^1 = F(Q = Q_0)$  and  $f^2 = F(Q = 2Q_0)$  in a reliable region ( $Q_0 > 2\text{GeV}$ ) and use  $F(Q) \frac{f^1}{f^2}$
- ▶ Theoretically motivated powerlaw fit<sup>7</sup>:


$$\frac{d\sigma^{\text{pp}}}{dp_T dy} = A_{\text{pp}} \left(1 + \frac{p_T^2}{P_0}\right)^{-n} \quad (4)$$

---

<sup>7</sup>Garcia-Montero, O., Löher, N., Mazeliauskas, A., Berges, J. & Reygers, K. *Phys. Rev. C* **102**, 024915. arXiv: 1909.12246 [hep-ph] (2020), Adare, A. et al. *Phys. Rev. C* **91**, 064904. arXiv: 1405.3940 [nucl-ex] (2015).

# Nuclear effects

- Photon is confined to  $|y| < 0.8$  with an isolation cone of  $R^{\text{iso}} = 0.4$  and an isolation energy of  $p_T^{\text{iso}} = 2$  GeV



## Direct photon codes

Shower Monte Carlo Event Generators:

Code	$pp \rightarrow \gamma j$	$pp \rightarrow \gamma jj$	$pp \rightarrow \gamma\gamma$
POWHEG	NLO+PS	[NLO+PS, this project]	[NLO+PS, not public]
Sherpa	NLO+PS	NLO+PS	NLO+PS
MG5,HW7	(NLO+PS)	(NLO+PS)	(NLO+PS)
Pythia	LO	-	LO
⋮			

Integrators:

Code	$pp \rightarrow \gamma j$	$pp \rightarrow \gamma jj$	$pp \rightarrow \gamma\gamma$
NNLOJET[not public]	NNLO+FF(NNLO)	[NLO]	NNLO+FF(NNLO)
JETPHOX/DIPHOX	NLO+FF(NLO)	[LO]	NLO+FF(NLO)
MCFM	NLO+FF(LO)	LO	NLO+FF(LO), NNLO
⋮			

Either FF or Parton showers (PS) is needed to generate physical results.



# Diphoton generators


Name	Final state	Diagrams / order				Various	Isolation		Reference
		Born	Box (*)	1 frag	2 frag		Cone	Frix.	
DIPHOS	YY	NLO	LO	NLO	NLO	Fixed order	X	X	T. Binoth, J.P. Guillet, E. Pilon and M. Werlen, Eur. Phys. J. C16 (2000) 311
GAMMA2MC	YY	NLO	NLO	-	-	Fixed order	X	X	Z. Bern, L. Dixon, and C. Schmidt, Phys. Rev. D66 (2002) 074018.
RESBOS	YY	NLO	NLO	LO	-	NLL	X	X	lots of papers...
MCFM	YY	NLO	NLO	LO	LO?	Fixed order	X	X	J. Campbell, R. Ellis, C. Williams, arXiv:1105.0020
2gNNLO	YY	NNLO	NLO	-	-	Fixed order		X	S. Catani et al, arXiv:1110.2375
PYTHIA / HERWIG	YY	LO	LO	PS	PS	-	X	X	-
ALPGEN	YY+up to 6 jets	LO	-	PS	PS	MLM matching with PYTHIA/HERWIG	X	X	<a href="http://mlm.home.cern.ch/mlm/alpger/">http://mlm.home.cern.ch/mlm/alpger/</a>
SHERPA	YY+X jets	LO	LO	PS	PS	CKKW + truncated shower	X	X	S. Hoeche, S. Schumann, F. Siegert, Phys.Rev.D81:034026,2010
POWHEG/H++	YY+1 jet	NLO	?	PS	PS	truncated shower	X	X	L. d'Errico, P. Richardson, arXiv:1106.3939 (diphoton code not public yet)
GR@PPA	YY+? jets	NLO	?	PS	PS	LL-subtraction	X	X	S. Tsuno et al., Comput. Phys. Commun. 175, 665 (2006) (no specific publi for diphoton yet)

Color code: yellow = integrator; blue = event generator

(\*): formally, (N)LO of box diagram is (N)NNLO - call it (N)LO b/c contributes as much as NLO born...

[https://indico.cern.ch/event/242419/contribution/19/attachments/412185/572744/LHC\\_France\\_SCHWOERER.pdf](https://indico.cern.ch/event/242419/contribution/19/attachments/412185/572744/LHC_France_SCHWOERER.pdf)

# NNLOJET





89

<sup>8</sup>Chen, X., Gehrmann, T., Glover, N., Höfer, M. & Huss, A. *JHEP* **04**, 166. arXiv: 1904.01044 [hep-ph] (2020).

<sup>9</sup>Chen, X. et al. in *16th DESY Workshop on Elementary Particle Physics: Loops and Legs in Quantum Field Theory 2022* (Aug. 2022). arXiv: 2208.02669 [hep-ph].

# NNLOJET




1011

<sup>10</sup>Chen, X., Gehrmann, T., Glover, N., Höfer, M. & Huss, A. *JHEP* **04**, 166. arXiv: 1904.01044 [hep-ph] (2020).

<sup>11</sup>Chen, X. et al. in *16th DESY Workshop on Elementary Particle Physics: Loops and Legs in Quantum Field Theory 2022* (Aug. 2022). arXiv: 2208.02669 [hep-ph].

# QGP temperature



12

<sup>12</sup>Klasen, M., Klein-Bösing, C., König, F. & Wessels, J. P. *JHEP* **10**, 119. arXiv: 1307.7034 [hep-ph] (2013).