

Fermionic Portal to Vector Dark Matter (FPVDM)

Alexander Belyaev




Southampton University & Rutherford Appleton Laboratory

AB, Deandrea, Moretti, Panizzi, Ross, Thongyoi – arXiv:2204.03510,2203.04681
AB, Panizzi, Thongyoi – to appear

Roadmap of Dark Matter Models for Run 3
CERN
May 13-17, 2024

Theory Space with Dark Matter



Vector DM and Vector-Like Fermionic Portal

- Higgs portal : the parameter space for minimal scenarios is almost excluded
- **Vector Like(VL) fermionic portal for Vector Dark Matter**
 - $SU(2)_D$ gauge triplet (new dark gauge) V_μ^D
 - Complex scalar doublet charged under $SU(2)_D$ Φ_D – to break gauge group
 - Vector-Like fermion doublet of $SU(2)_D$ Ψ – to “talk” to SM

Vector DM and Vector-Like Fermionic Portal

- Higgs portal : the parameter space for minimal scenarios is almost excluded
- **Vector Like(VL) fermionic portal for Vector Dark Matter**
 - $SU(2)_D$ gauge triplet (new dark gauge) V_μ^D
 - Complex scalar doublet charged under $SU(2)_D$ Φ_D – to break gauge group
 - Vector-Like fermion doublet of $SU(2)_D$ Ψ – to “talk” to SM
 - we assign the “dark charge” to the components of the doublets, e.g. $Q_D = T_D^3 + Y_D$ and require its conservation
 - $SU(2)_D \times U(1)_{\text{glob}} \rightarrow U(1)_{\text{glob}}^d$
pattern of dark sector breaking
 - \mathbb{Z}_2 subgroup can be defined as: $(-1)^{Q_D}$

Vector DM and Vector-Like Fermionic Portal

	$SU(2)_L$	$U(1)_Y$	$SU(2)_D$	Q_D	Z_2
$\Phi_D = \begin{pmatrix} \varphi_{D+\frac{1}{2}} \\ \varphi_{D-\frac{1}{2}} \end{pmatrix} \rightarrow \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ H_D^+ \nu_D \end{pmatrix}$	1	0	2	+1 0	- +
$\Psi = \begin{pmatrix} \Psi_D \\ \Psi \end{pmatrix} = \begin{pmatrix} \tilde{F} \\ F \end{pmatrix}$	1	Q_{EM}	2	+1 0	- +
$V_M^D = \begin{pmatrix} V_M^{D+} \\ V_M^{D0} \\ V_M^{D-} \end{pmatrix} = \begin{pmatrix} \tilde{V}_D^+ \\ V' \\ \tilde{V}_D^- \end{pmatrix}$	1	0	3	+1 0 -1	- + -

- If we chose $Y_D = +1/2$ for Φ_D and Ψ then we have $y' \bar{\Psi}_L \Phi_D f_R^{SM} + y'' \bar{\Psi}_L \Phi_D^c f_R^{SM} + h.c$

y'' is eliminated under the requirement of Q_D conservation, DM is established!

Fermionic Portal for Vector Dark Matter (FPVDM)

- It is the framework, representing the class of models
(Deandrea, Moretti, Panizzi, Ross, Thongyoi, AB – arXiv:2204.03510,2203.04681)
- Various realisations are possible, including one or several VL fermions

$$\mathcal{L}_{FPVDM} = -\frac{1}{4}(V_{D\mu\nu}^i)^2 + \bar{\Psi}iD\Psi + |D_\mu\Phi_D|^2 - V(\Phi_H, \Phi_D) - \underline{(y'_{\alpha\beta}\bar{\Psi}_L^{i\alpha}\Phi_D f_R^{SM\beta} + h.c.)} - M_\Psi^{ij}\bar{\Psi}^i\Psi^j$$

$$V(\Phi_H, \Phi_D) = -\mu_H^2\Phi_H^\dagger\Phi_H - \mu_D^2\Phi_D^\dagger\Phi_D + \lambda_H(\Phi_H^\dagger\Phi_H)^2 + \lambda_D(\Phi_D^\dagger\Phi_D)^2 + \lambda_{HD}(\Phi_H^\dagger\Phi_H)(\Phi_D^\dagger\Phi_D)$$

- $y'_{\alpha\beta}$ can have a flavour structure – to explain flavour anomalies
- λ_{HD} can be negligible at tree-level, DM can be well-generated via FP

- the model with $\Psi = \begin{pmatrix} \tilde{T} \\ T \end{pmatrix}$ and $\lambda_{HD} = 0$ was explored

Minimal VL top portal VDM: collider signatures

Process	Representative diagrams
mono-jet (only loop)	<p>E_T^{miss} + jet from ISR or from loop</p>
$t\bar{t} + E_T^{\text{miss}}$	<p>E_T^{miss}</p>
$t\bar{t}t\bar{t}$	
hV' and $V'V'$ (only loop)	

FPVDM model with $\Psi_M = \begin{pmatrix} \tilde{M} \\ M' \end{pmatrix}$, **the partner of muon**

$\mathcal{L}_{\mu PVDM} \supset -y' \bar{\Psi}_{ML} \Phi_D \mu_R + h.c$ with $\tilde{V}_D, V', H_D, M', \tilde{M}$

- has potential to explain DM relic density and $(g-2)_\mu$ anomaly

FPVDM model with $\Psi_M = \begin{pmatrix} \tilde{M} \\ M' \end{pmatrix}$, the partner of muon



$$\mathcal{L}_{\mu PVDM} \supset -y' \bar{\Psi}_{ML} \Phi_D \mu_R + h.c. \quad \text{with } \tilde{V}_D, V', H_D, M', \tilde{M}$$

- has potential to explain DM relic density and $(g-2)_\mu$ anomaly
- one should ensure
 - consistency with DD and ID DM search experiments
 - consistency with collider searches

FPVDM model with $\Psi_M = \begin{pmatrix} \tilde{M} \\ M' \end{pmatrix}$, the partner of muon

$\mathcal{L}_{\mu PVDM} \supset -y' \bar{\Psi}_{ML} \Phi_D \mu_R + h.c$ with $\tilde{V}_D, V', H_D, M', \tilde{M}$

- has potential to explain DM relic density and $(g-2)_\mu$ anomaly
- one should ensure
 - consistency with DD and ID DM search experiments
 - consistency with collider searches
- Parameter space ($\lambda_{HD} = 0$ for simplicity): $g_D, m_{V_D}, m_{H_D}, m_{M'}, m_{\tilde{M}}$
- Interactions+mixing:




$$y' \bar{\Psi}_L \Phi_D f_R^{\text{SM}} + h.c$$

FPVDM model with $\Psi_M = \begin{pmatrix} \tilde{M} \\ M' \end{pmatrix}$, the partner of muon

$$\mathcal{L}_{\mu PVDM} \supset -y' \bar{\Psi}_{ML} \Phi_D \mu_R + h.c \quad \text{with } \tilde{V}_D, V', H_D, M', \tilde{M}$$

- has potential to explain DM relic density and $(g-2)_\mu$ anomaly
- one should ensure
 - consistency with DD and ID DM search experiments
 - consistency with collider searches
- Parameter space ($\lambda_{HD} = 0$ for simplicity): $g_D, m_{V_D}, m_{H_D}, m_{M'}, m_{\tilde{M}}$
- Interactions+mass corrections:



The status of $(g-2)_\mu$ and our approach here

- The combined experimental value from BNL +FNAL(from August 2023):

$$a_\mu^{\text{EXP}} = 116592059(22) \times 10^{-11}$$

- The SM Theory Initiative 2020 prediction [arXiv:2006.04822] provides


$$a_\mu^{\text{SM}} = 116591810(43) \times 10^{-11}$$

- Combining above numbers, one concludes one finds **5.1 σ SM vs EXP discrepancy**

$$\Delta a_\mu = a_\mu^{\text{EXP}} - a_\mu^{\text{SM}} = 249(48) \times 10^{-11}$$


- Theory: for three contributions to $(g-2)_\mu$ – QED, EW and Hadronic – the Hadronic Vacuum Polarisation (HVP) is taken from the experimental data and it has the biggest contribution to the uncertainty
- Recent CMD3 results [arXiv:2302.08834] adds and additional intrigue here

- Of course recent Lattice results from BMW [Nature 593, 51 (2021)] must be add here




- $(g-2)_\mu$ is an important puzzle to be solved including discrepancy between HVP from e+e- data and Lattice
- In our study we take Δa_μ as a real effect to be explained within our μ FPVDM model


The contribution to $(g-2)_\mu$ from μ PVDM




a




b



c








d



e

The contribution to $(g-2)_\mu$ from μ PVDM \times

$$\frac{g_D^2}{96\pi^2} \frac{m_\mu^2}{m_{V_D}^2}$$





$$\Delta M^2 = m_{M'}^2 - m_{\tilde{M}}^2$$

$$M = m_{M'}$$

$M \gg m_{V_D}$ was used

$(g-2)_\mu$ results from scan of $g_D, m_{V_D}, m_{H_D}, m_{M'}, m_{\tilde{M}}$ space

$$\Delta \hat{a}_\mu = (\Delta a_\mu^{\mu PVDM} - \Delta a_\mu) / \sigma_{a_\mu} \equiv \frac{\Delta a_\mu^{\mu PVDM} - 249}{48} \times 10^{-11}$$




- Δa_μ can be explained within μ FPVDM model ($g_D/m_{V_D} \sim 0.1$)
- $g_D - m_{V_D}$ correlation can be clearly observed as predicted by analytical calculations
- For $m_{M'} > 1 \text{ TeV}$ it is hard (but possible) to explain Δa_μ because of $1/m_{M'}^2$ suppression

We also aim to explain DM relic density & to be consistent with DM DD and ID as well as with collider searches

Correct Relic density: efficient (co) annihilation
 annihilation to photons can affect CMB



$$\Omega_{\text{DM}}^{\text{Planck}} h^2 = 0.12 \pm 0.0012$$



We also aim to explain DM relic density & to be consistent with DM DD and ID as well as with collider searches

Correct Relic density: efficient (co) annihilation
annihilation to photons can affect CMB


$$\Omega_{\text{DM}}^{\text{Planck}} h^2 = 0.12 \pm 0.0012$$




Tools used

- **DM DD, ID, Relic density**
LanHEP, CalcHEP, micrOMEGAs
- **Collider searches**
CalcHEP, MC@NLO, PYTHIA, DELPHES, MadAnalysis, CHECKMATE


Cosmological constraints on μ PVDM parameter space







- generic DM annihilation
- H_D resonance region
- \tilde{M} co-annihilation region
- underabundant DM region
- overabundant DM region
- excluded by CMB DM ID
- excluded by DM DD





Cosmological constraints on μ PVDM parameter space



Cosmological constraints on μ PVDM parameter space







■ both DM ID and DD exclusions play an important complementary role




$m_{\tilde{M}} > 100$ GeV
Applied to respect LEP constraints

m_{H_D} vs m_{V_D} and m_{V_D} vs m_{μ_D} planes




Combining $(g-2)_\mu$ and DM constraints



$$g_D \sim m_{V_D}$$

“trajectory” to explain Δa_μ


- ★ $DD + ID + \Omega h^2_{PLANCK} + \Delta(g-2)_\mu @ 2\sigma$
- $DD + ID + 15\% \Omega h^2_{PLANCK} + \Delta(g-2)_\mu @ 2\sigma$
- $\Delta(g-2)_\mu$ allowed @ 2σ
- DD+ID allowed + Ωh^2_{PLANCK}
- DD+ID allowed, $\Omega h^2 < 0.12$
- allowed by perturbativity



$$\Omega h^2 \propto \frac{M_{V_D}^2}{g_D^4} \frac{1}{\sqrt{g_{eff}(T_f)}}$$


$$g_D \sim \sqrt{M_{V_D}} / [g_{eff}(T_f)]^{1/2}$$

Ωh^2_{PLANCK}
“trajectory” to explain DM relic density






- $(g-2)$ and DM relic density allowed bands have different slopes crossing at **0.1 – 1 GeV**
 - “dark photon”(V) kind of region
 - New collider signatures (see below)
 - very intriguing to explore further for GW effects and explaining NANOGrav results

Final very set important constraints: colliders



$$pp \rightarrow \tilde{M}^- \tilde{M}^+ \rightarrow \tilde{V}_D \tilde{V}_D \mu^+ \mu^-$$



- Madgraph + PTHIA+Delphes + Madanalysis
- $\tilde{M} > 600$ GeV comes from the main $\mu^+ \mu^- + MET$




- surviving region after all constraints

The parameter space and signatures after all constraints

- ★ $DD + ID + \Omega h_{\text{PLANCK}}^2 + \Delta(g-2)_\mu @ 2\sigma + \text{LHC}$
- $DD + ID \text{ allowed} + \Omega h_{\text{PLANCK}}^2$
- $DD + ID + 15\% \Omega h_{\text{PLANCK}}^2 + \Delta(g-2)_\mu @ 2\sigma$
- $DD + ID \text{ allowed}, \Omega h^2 < 0.12$
- $\Delta(g-2)_\mu \text{ allowed} @ 2\sigma$
- $\text{allowed by perturbativity}$




10μ
final
state



$6\mu + \cancel{E}_T$
final
state

exciting new signatures: 10μ , 8μ , $6\mu(+\text{MET})$, $4\mu(+\text{MET})$, $2\mu+\text{MET}$:
hi-PT pairs of muons (from V' decay) + single high-pT muons

The parameter space and signatures after all constraints




Inputs/Observables	BP1	BP2	BP3
g_D	0.003	0.003	0.003
m_{V_D} [GeV]	0.28	0.28	0.28
m_{μ_D} [GeV]	800	900	1000
$m_{\mu'}$ [GeV]	1000	1200	1400
m_{H_D} [GeV]	0.677	0.677	0.677
$m_{V'}$ [GeV]	0.2756	0.2706	0.2637
$Br(\mu' \rightarrow V' \mu)$	0.383	0.342	0.318
$Br(\mu' \rightarrow H_D \mu)$	0.371	0.319	0.282
$Br(\mu' \rightarrow V_D \mu_D)$	0.246	0.339	0.4
$Br(H_D \rightarrow V_D V_D^*)$	0.639	0.612	0.575
$Br(H_D \rightarrow V' V')$	0.352	0.375	0.409
$Br(H_D \rightarrow \mu^+ \mu^-)$	9.24×10^{-3}	1.31×10^{-2}	1.54×10^{-2}
$Br(V' \rightarrow \mu^+ \mu^-)$	~ 1	~ 1	~ 1
$Br(\mu' \rightarrow V' \mu \rightarrow 3\mu)$	0.383	0.342	0.318
$Br(\mu' \rightarrow H_D \mu \rightarrow 5\mu)$	0.131	0.12	0.115
$\sigma_{\text{tot}}(pp \rightarrow \mu' \mu')$ [fb]	6.499×10^{-2}	1.867×10^{-2}	6.32×10^{-3}
$N_{\text{event}}(pp \rightarrow 6\mu)$	2.86	0.655	0.192
$N_{\text{event}}(pp \rightarrow 8\mu)$	0.978	0.23	0.069
$N_{\text{event}}(pp \rightarrow 10\mu)$	0.335	0.08	0.025

assuming 300 fb^{-1} for the integrated luminosity


The model predicts sub-GeV V' bosons which look like dark-photons, but not quite...

- V' bosons have **kinetic mixing** with photons and Z-bosons similarly to dark-photons
- At the same time V' bosons have **significant coupling to SM fermion** which is the partner of VL dark fermion
- As a result, V' bosons will can promptly decay (if kinematically allowed) to SM fermions **avoiding existing bound on dark-photons: requires dedicated analysis**



JHEP 09 (2023) 035, [2303.08666](https://arxiv.org/abs/2303.08666) [hep-ex]

Inputs/Observables	BP1	BP2	BP3
$\tau_{V'}$ [ns]	1.10×10^{-6}	7.85×10^{-7}	6.77×10^{-7}
$l_{V'}$ [μm]	0.33γ	0.24γ	0.20γ
$\epsilon_{AV'}$	1.13×10^{-5}	1.39×10^{-5}	1.578×10^{-5}



Summary on FPVDM

- FPVDM is a very promising new framework for VDM, not requiring Higgs portal
- Incorporates many possibilities with new collider and cosmological implications
 - great potential to explain dark matter
 - collider signatures: ff+ET miss, V' , $Z'H$, long-lived V'
- Great potential to explore flavour, was not deliberately designed for this
- Models available at hepmdb.soton.ac.uk
- The model with VL partner of muon – the paper to appear (AB, Panizzi, Thongyoi)
 - can explain relic density and Δa_μ
 - collider constraints + simultaneous explanation of DM requires a very specific parameter space: $M_{DM} \sim 0.2 \text{ GeV}$, $M_{HD} \sim 2 M_{DM}$, $g_D \sim 0.01$ to avoid DM ID (CMB constraints)
 - Provides multi-lepton signatures, including di-lepton boosted pairs (up to 10 muons)
 - Has non-standard “dark-photon” (V'), which avoids present experimental constraints and requires dedicated searches
 - The low DM mass range makes it interesting for the connection to GW data from NANOGrav