

Introduction and motivation

Improve muon selection and identification (ID) for muons with **medium p_T range** → key in many CMS analysis. Whether those are:

- Isolated muons from prompt decay of W , Z , H and decay of τ .
- Real muons from heavy flavour decays.

Multivariate analysis (MVA) methods used to construct **two muon ID** discriminators:

- **MVA ID: General muon selection** against spurious hits in the muon system. We construct an ID more flexible and performant than the cut-based ID [1] (used during Run 2) for muons with p_T between 20 and 120 GeV. ID to be used during Run 3.
- **Prompt MVA: select isolated muons from H/W and τ .** Already used during Run 2, crucial in the observation of $t\bar{t}H$, $t\bar{t}t$ and tZq ; the precision measurement of WZ and to increase sensitivity in the search for electroweak production of SUSY.

Input Variables

MVA ID

12 Input variables related to:


- Quality of the track reconstructed in the muons system (stand alone) and in the tracker
- Quality matching between both
- Number of muon station with hits
- Number of good hits in the tracker

Prompt MVA

12 Input variables related to:

- Isolation of the muon
- Information of the jet reconstructed within the same cone
- Impact parameter (IP)

Example of a input variables in each MVA:



Pre-selection and samples

- Used **2018** dataset (59.7 fb⁻¹)
- **MC:**
 - $t\bar{t}$ semileptonic for training
 - DY for performance evaluation
- **Pre-selected muons** are loose muons (cut-based ID) with $p_T > 10$ GeV + **Relaxed isolation and IP selection*** (for the prompt MVA)
- look for geometrical matching traced back using generation information


*Mini-isolation < 0.4 & $d_{xy} < 0.05$ & $d_z < 0.1$ & $d/\sigma_d < 8$

Machine learning models

MVA ID

Random Forest: 200 trees with a maximum depth of 8

Medium working point defined to have same background efficiency as the medium cut-based ID





Prompt MVA:

BDT: 200 trees with a maximum depth of 8

1000 trees with max depth = 4

WP shown optimized for $t\bar{t}H$ measurement



Do you select muons in your analysis?
We have an MVA for you!



arXiv:2310.03844
Accepted by JINST

Performance evaluation

Efficiency in data evaluated using **Tag-and-Probe** method to select muons from Z decay:

- Tag: pass Tight WP (cut-based ID)
- Probe: preselection

Background estimation:

MVA ID: estimated using $t\bar{t}MC$


$$Bkg. Efficiency = \frac{\text{background loose muon \& pass ID selection}}{\text{background loose muon}}$$

Prompt MVA:

- Nonprompt rate measured in a Multi-jet enriched region
- EWK contributions subtracted using a fit to:

$$m_T^{fix} = \sqrt{2 p_T^{fix} p_T^{miss} (1 - \cos \Delta \phi)}$$


$$p_T^{fix} = 35 \text{ GeV}$$



Background efficiency: 50% for $p_T > 20$ GeV

For same background efficiency, systematically higher efficiency than with cut based

A factor of 2 (3) smaller nonprompt rate of the prompt MVA for muons with $|\eta| < 1.2$ ($|\eta| > 1.2$)



Summary

MVA ID

- **Higher efficiency than cut-based ID** (for the same background efficiency)
- Continuous score gives more flexibility for analysers
- MVA shown to be more **resilient as a function of PU** than the cut-based ID → **Great news for run 3!**

Prompt MVA

- Efficiency $> 80\%$ for $p_T > 20$ GeV
- Factor of 2-3 reduction of background from non prompt leptons
- Has been a **key element** in several full **Run 2** CMS published results: $t\bar{t}H$, $t\bar{t}t$, tZq , WZ , EWK SUSY...

Ayudas del Programa Severo Ochoa by



Grant PID2020-113341RB-I00 funded by

