

Improving our understanding of $B \rightarrow D \pi \ell \nu$ and $B \rightarrow \pi \pi \ell \nu$ decays

Tuesday 24 September 2024 14:00 (30 minutes)

In this talk, we present studies of two semileptonic decays with two hadrons in the final state. First, we'll discuss $B \rightarrow D \pi \ell \nu$ decays that not only constitute a sizeable background for $R(D^{(*)})$ determinations, but also comprise a largely unknown signal component for inclusive $B \rightarrow X_c \ell \nu$ decays. The dominant decay chain proceeds through resonant $B \rightarrow D^*/D_2^*(\rightarrow D \pi) \ell \nu$ decays, but also includes a poorly understood broad component. We study the composition of this broad component by using our newly developed formalism for $B \rightarrow D \pi \ell \nu$ decays, together with experimental spectral measurements, and investigate the plausibility of the two-pole structure in the $D \pi$ S-wave. In addition, we provide recommendations for possible future measurements to systematically improve current understanding of $B \rightarrow D \pi \ell \nu$ decays kinematics.

In the second part of the talk we present new results on $B \rightarrow \pi \pi \ell \nu$ decays, relevant to studies of $B \rightarrow \rho \ell \nu$ and V_{ub} determinations. We discuss the separation of signal P-wave from background S-wave contributions in the ρ -region and the use of $e^+e^- \rightarrow \pi^+\pi^-$ data to describe the ρ -lineshape. In addition, we briefly discuss the inclusion of the leading isospin breaking effects in $B \rightarrow \pi \pi \ell \nu$ decays.

Primary author: VAN TONDER, Raynette (McGill University)

Co-author: HERREN, Florian (UZH)

Presenter: VAN TONDER, Raynette (McGill University)

Session Classification: Heavy to heavy exclusive