


Monte Carlo methods for stationary solutions of
general-relativistic Vlasov systems:
Planar accretion onto a moving Schwarzschild black hole

Adam Cieřlik
with
Patryk Mach and Andrzej Odrzywołek





Schwarzschild black hole moving through a cloud of gas
(P. Mach and A. Odrzywołek: 2021,2022)

Our case

- Stationary solution – geodesic trajectories do not depend on time
- Bondi-type accretion – a compact object travelling through the interstellar medium
- Planar accretion
- Asymptotically, the gas is assumed to be homogeneous and described by the two-dimensional Maxwell-Jüttner distribution, boosted with a constant velocity v along the x axis. In the Cartesian coordinates, the asymptotic distribution function is given by

$$\mathcal{F}(x, p) = \alpha \delta(\sqrt{-p_\mu p^\mu} - m_0) \exp\left[\frac{\beta}{m_0} \gamma(p_t + v p_x)\right],$$

and in spherical coordinates

$$\mathcal{F}(x, p) = \alpha \delta(\sqrt{-p_\mu p^\mu} - m_0) \exp\left[\frac{\beta}{m_0} \gamma(p_t + v \cos \varphi p_r)\right].$$

Particle current density

With the help of Hamilton's formalism and action-angle variables, it can be shown that in the vicinity of the Schwarzschild black hole, the distribution function is given by

$$f(x, p) = \alpha \delta(\sqrt{-p_\mu p^\mu} - m_0) \exp \left\{ -\beta \gamma \left[\varepsilon - \epsilon_r v \sqrt{\varepsilon^2 - 1} \cos [\varphi + \epsilon_\varphi \epsilon_r X(\xi, \varepsilon, \lambda)] \right] \right\},$$

where

$$X(\xi, \varepsilon, \lambda) = \lambda \int_\xi^\infty \frac{d\xi'}{\xi'^2 \sqrt{\varepsilon^2 - \left(1 - \frac{2}{\xi'}\right) \left(1 + \frac{\lambda^2}{\xi'^2}\right)}}.$$

Additionally, components of surface particle current density read

$$J_\mu(\xi, \varphi) = \sum_{\epsilon_\varphi = \pm 1} \frac{1}{\xi} \int f(\xi, \varphi, m, \varepsilon, \epsilon_\varphi, \lambda) p_\mu \frac{m^2 dm d\varepsilon d\lambda}{\sqrt{\varepsilon^2 - U_\lambda}}.$$

Monte Carlo approach

Consider a discrete distribution function

$$\mathcal{F}^{(N)}(x^\mu, p_\nu) = \sum_{i=1}^N \int \delta^{(4)}(x^\mu - x_{(i)}^\mu(\tau)) \delta^{(4)}(p_\nu - p_{\nu}^{(i)}(\tau)) d\tau$$

representing a sample of N particles moving along given trajectories $\tau \mapsto (x_{(i)}^\mu(\tau), p_{\nu}^{(i)}(\tau))$, $i = 1, \dots, N$. The particle current density associated with $\mathcal{F}^{(N)}$ is given as

$$\mathcal{J}_\mu^{(N)}(x) = \int_{P_x^+} \mathcal{F}^{(N)}(x, p) p_\mu \sqrt{-\det g^{\alpha\beta}(x)} dp_0 dp_1 dp_2 dp_3.$$

Let $\Sigma \in \mathcal{M}$ be a hypersurface, not necessarily spacelike. We choose a small region $\sigma \in \Sigma$ (a numerical cell) such that $x \in \sigma$. The components of \mathcal{J}_μ are approximated by the averages

$$\langle \mathcal{J}_\mu(x) \rangle = \frac{\int_\sigma \mathcal{J}_\mu^{(N)} \eta_\Sigma}{\int_\sigma \eta_\Sigma},$$

where η_Σ denotes the volume element on Σ .

Intersections of trajectories with arcs of constant radius

For a planar stationary accretion flow in the Schwarzschild spacetime, we select surfaces of constant $r = \bar{r}$ defined by


$$\tilde{\Sigma} = \{(t, r, \theta, \varphi) : t \in \mathbb{R}, r = \bar{r}, \theta = \pi/2, \varphi \in [0, 2\pi)\}$$

and cells

$$\tilde{\sigma} = \{(t, r, \theta, \varphi) : t_1 \leq t \leq t_2, r = \bar{r}, \theta = \pi/2, \varphi_1 \leq \varphi \leq \varphi_2\}.$$

More precisely let $\Phi_\tau(x_0^i)$ denote the orbit of timelike Killing vector field $\xi^\mu = (1, 0, 0, 0)$, passing through x_0^i at $\tau = 0$, i.e., $\Phi_0(x_0^i) = x_0^i$. Then $\tilde{\sigma}$ can be expressed as the image

$$\tilde{\sigma} = \Phi_{[t_1, t_2]}(S).$$



The particle current surface density can now be approximated as

$$\langle J_\mu \rangle = \frac{\int_{\tilde{\sigma}} J_\mu \eta_{\tilde{\Sigma}}}{\int_{\tilde{\sigma}} \eta_{\tilde{\Sigma}}} = \frac{1}{Mm\bar{\xi}(t_2 - t_1)(\varphi_2 - \varphi_1)} \sum_{j=1}^{N_{\text{int}}} \frac{p_\mu^{(j)}}{\sqrt{\varepsilon_{(j)}^2 - (1 - 2/\bar{\xi}) \left(1 + \lambda_{(j)}^2/\bar{\xi}^2\right)}}.$$

For stationary problems, the result should be independent of the choice of t_1 and t_2 in a sense that the number of trajectories that intersects $\tilde{\Sigma}$ should be proportional to the length $t_2 - t_1$, if the latter is sufficiently large. In practice, we omit the factor $t_2 - t_1$ and normalise the results by the number of trajectories taken into account. Moreover, instead of considering complete orbits in the four-dimensional spacetime, it is sufficient to work with projections of trajectories onto surfaces of constant t .





Selection of geodesic parameters

- We select the parameters $\{\xi_0, \varphi_i^{(\text{init})}, \varepsilon_i, \lambda_i\}$, representing the radial and the azimuthal coordinates of the initial position, the energy, and the total angular momentum of i -th particle, respectively
- The first coordinate is the same for all trajectories—all particles start at a fixed radius $r_0 = M\xi_0$. It is important to ensure that this value is sufficiently large
- The coordinate values $\varphi_i^{(\text{init})}$ and ε_i are sampled from the planar asymptotic ($\xi \rightarrow \infty$) distribution function:


$$f(x, p) = \alpha \delta(\sqrt{-p_\mu p^\mu} - m) \exp\left[-\beta \gamma \left(\varepsilon - \varepsilon_r v \sqrt{\varepsilon^2 - 1} \cos \varphi\right)\right] \quad (1)$$

- To randomise the parameters $\varphi_i^{(\text{init})}, \varepsilon_i$ according to the distribution function (1), we use the Markov Chain Monte Carlo (MCMC) method, implemented in the *Wolfram Mathematica*.
- Values of λ_i are distributed uniformly.
- From the selected parameters, we choose those corresponding to the unbound trajectories and then divide them into absorbed and scattered


Results




$$\varphi = 0.5^\circ, v = 0.95c, \beta = 1$$




$$\varphi = 44.5^\circ, v = 0.95c, \beta = 1$$




$$\varphi = 90.5^\circ, v = 0.95c, \beta = 1$$




$$\varphi = 180.5^\circ, v = 0.95c, \beta = 1$$




$$\varphi = 0.5^\circ, v = 0.95c, \beta = 1$$




$$\varphi = 44.5^\circ, v = 0.95c, \beta = 1$$







$$\varphi = 90.5^\circ, v = 0.95c, \beta = 1$$



$$\varphi = 180.5^\circ, v = 0.95c, \beta = 1$$





Conclusions

- We confirmed the analytical results in the case of planar accretion
- We demonstrated that the developed Monte Carlo simulation method can be used for cases that do not have spherical symmetry
- Outlook:
 - Preparation of a three-dimensional simulation (P. Mach and A. Odrzywołek: 2021, 2022)
 - Preparing a simulation of Vlasov gas accretion in Kerr spacetime (A. Cieřlik, P. Mach, A. Odrzywołek: 2022; P. Rioseco, O. Sarbach: 2018, 2023)
 - Generalisation to general-relativistic Vlasov systems coupled with the electromagnetic field (M. Thaller: 2023)