

Model independent Odderon results based on new TOTEM data at 8 TeV

T. Csörgő^{1,2} T. Novák², R. Pasechnik³, A. Ster¹, I. Szanyi^{1,2,4}

¹ Wigner RCP, Budapest, Hungary

² MATE KRC, Gyöngyös, Hungary

³ University of Lund, Lund, Sweden

⁴ Eötvös University, Budapest, Hungary



MATE

WIGNER

Supported by NKFIH:
K 74458, Hungary

Outline

Statistically Significant Observations of Odderon in 2021

Model independent (Hungarian-Swedish Collaboration):

Significance $\geq 6.26 \sigma$: *EPJC (2021) 81:180*

Model dependent (Hungarian-Polish Collaboration):

Significance $\geq 7.08 \sigma$:

EPJC (2021) 81:611 and EPJC (2022) 82:827

Partially model independent (D0-TOTEM Collaboration):

Significance $\geq 5.2 \sigma$: *PRL (2021) 127, 062003*

New in 2022:


TOTEM data at 8 TeV published

Motivation: Einstein said something like: „Even infinite number of experiments can not prove that you are right, but one experiment is enough to prove that you are wrong”

Odderon: 48 years old scientific puzzle

Odderon: L. Lukaszuk, B. Nicolescu,
Lett. Nuovo Cim. 8, 405 (1973)
Received: 31 July 1973

Odderon is an odd component of
elastic scattering:
Changes sign for crossing



СООБЩЕНИЯ
ОБЪЕДИНЕННОГО
ИНСТИТУТА
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
Дубна



E2-6350

A.V.Efremov, R.Peschanski

EVIDENCE FOR NEW SINGULARITIES
IN REGGE PHENOMENOLOGY



1972

ЛАБОРАТОРИЯ ТЕОРЕТИЧЕСКОЙ ФИЗИКИ

Odderon name coined: D. Joynson, E. Leader, B. Nicolescu, C. Lopez,
Nuovo Cim. 30A, 345 (1975) - Well established in QCD by now !
Honorable mention: A. V. Efremov, R. Peschanski, JINR-E2-6350 (1972)

Odderon: elusive experimentally

Odderon search at ISR: indication but no conclusive result
Breakstone et al, Phys. Rev. Lett. 54, 2180 (**1985**): CL = 99.9 %






Indication of Odderon
CL = 99.9 %,
Significance: 3.35σ

2021 observations of Odderon with $> 5 \sigma$

Evidence of Odderon-exchange from scaling properties of elastic scattering at TeV energies #5

T. Csörgő (Wigner RCP, Budapest and CERN), [T. Novák](#) (Unlisted, HU), R. Pasechnik (Lund U., Dept. Theor. Phys.), [A. Sten](#) (Wigner RCP, Budapest), [I. Szanyi](#) (Wigner RCP, Budapest) (Dec 26, 2019)

Published in: *Eur.Phys.J.C* 81 (2021) 2, 180 • e-Print: 1912.11968

 pdf  DOI  cite




Hungarian-Swedish Odderon:

Eur. Phys. J. C (2021) **81**: 180, [Published: 23 February 2021](#)
<https://doi.org/10.1140/epjc/s10052-021-08867-6>

Observation of Odderon effects at LHC energies: a real extended Bialas–Bzdak model study #2

T. Csorgo (Wigner RCP, Budapest and EKV KRC, Gyongyos), I. Szanyi (Eotvos U. and Wigner RCP, Budapest) (May 28, 2020)

Published in: *Eur.Phys.J.C* 81 (2021) 7, 611 • e-Print: 2005.14319

 pdf  DOI  cite

Hungarian-Polish Odderon:

Eur. Phys. J. C (2021) **81**:611, [Published: 13 July 2021](#)
<https://doi.org/10.1140/epjc/s10052-021-09381-5>

Odderon Exchange from Elastic Scattering Differences between pp and $p\bar{p}$ Data at 1.96 TeV and from pp Forward Scattering Measurements #1

TOTEM and D0 Collaborations • V.M. Abazov (Dubna, JINR) et al. (Dec 7, 2020)

Published in: *Phys.Rev.Lett.* 127 (2021) 6, 062003 • e-Print: 2012.03502

 pdf  links  DOI  cite

D0-TOTEM Odderon:




Phys. Rev. Lett. **127** (2021) 6, 062003, [Published: 4 August 2021](#)
<https://doi.org/10.1103/PhysRevLett.127.062003>

2022 observations of Odderon with $> 5 \sigma$

Characterisation of the dip-bump structure observed in proton–proton elastic scattering at $\sqrt{s} = 8 \text{ TeV}$ #1

TOTEM Collaboration • G. Antchev (Pilsen U.) et al. (Nov 23, 2021)

Published in: *Eur.Phys.J.C* 82 (2022) 3, 263 • e-Print: 2111.1195




 pdf  DOI  cite

8 TeV: EPJ C (2022) 82, 263 (2022). Published: March 26, 2022
<https://doi.org/10.1140/epjc/s10052-022-10065-x>
Publishes final data for D0-TOTEM PRL published in 2021

The ReBB model and its $H(x)$ scaling version at 8 TeV: Odderon exchange is a certainty #1

I. Szanyi (Eotvos U. and Wigner RCP, Budapest and Karoly Robert U. Coll.), T. Csörgő (Wigner RCP, Budapest and Karoly Robert U. Coll.) (Apr 21, 2022)

Published in: *Eur.Phys.J.C* 82 (2022) 9, 827, *Eur.Phys.J.C* 82 (2022) 9, 827

 pdf  DOI  cite

New TOTEM 8 TeV data vs ReBB model predictions:
EPJ C 82 (2022) 9, 827. Published: Sept 19, 2022
In the ReBB model, Odderon exchange is a certainty
Presented at Zimányi'22 by I. Szanyi

Model independent observation, 2019 -

Definition of the model independent $H(x)$ scaling function constructed from published data:

$$H(x) = 1/(B \sigma_{el}) d\sigma/dt, \quad x = B t$$

B: slope at $t=0$ (data, published with $d\sigma/dt$)

σ_{el} : total elastic σ (data, published with $d\sigma/dt$)


Definition of χ^2 according to def. in ref. of PHENIX Collaboration:

$$\chi_{2 \rightarrow 1}^2 = \sum_{j=1}^{n_{21}} \frac{(d_1^j + \epsilon_{b,1} e_{B,1}^j - d_{21}^j - \epsilon_{b,21} e_{B,21}^j)^2}{(\tilde{e}_{A,1}^j)^2 + (\tilde{e}_{A,21}^j)^2} + \epsilon_{b,1}^2 + \epsilon_{b,21}^2,$$

$$\tilde{e}_{A,k}^j = e_{A,k}^j \frac{d_k^j + \epsilon_{b,k} e_{B,k}^j}{d_k^j},$$

$$e_{M,k}^j = \sqrt{(\sigma_{M,k}^j)^2 + (d_k^{j,j})^2 (\delta_{M,k}^j x)^2},$$

Model independent observation, 2019 -





$$B \equiv B_0(s)$$

$$x = - B t = - B_0(s) t$$

Data agree within 1 of σ standard deviation.

H(x) scaling of elastic p+p scattering data at ISR energies of 23 - 63 GeV. (See details in the publication)

Model independent observation, 2019 -



$H(x) = 1 / (B \sigma_{el}) \, d\sigma/dt \, \text{vs} \, x = -Bt$

$B \equiv B_0(s)$ from now on

$x = -Bt = -B_0(s) t$


S: Model independent Odderon significance $\geq 6.26 \sigma$
C1: All D0 and TOTEM published data at 1.96, 2.76 and 7.0 TeV
C2: domain of validity is still determined model dependently.

Model dependent observation, 2020-

Observation of Odderon Effects at LHC energies -- A Real Extended Bialas-Bzdak Model Study #2

T. Csorgo (Wigner RCP, Budapest and Eötvös Loránd University, Gyöngyös), I. Szanyi (Eötvös Loránd University, Budapest)
 e-Print: 2005.14319 [hep-ph]

Eur. Phys. J. C (2021) 81:611, published July 2021
<https://doi.org/10.1140/epjc/s10052-021-09381-5>



S: Model dependent Odderon significance $\geq 7.08 \sigma$

C1: All D0 and TOTEM published data at 1.96, 2.76, and 7.0 TeV

C2: domain of validity extended to both pp and pbarp

But limited to $0.37 \leq -t \leq 1.2 \text{ GeV}^2$ and $0.546 \leq \sqrt{s} \leq 7 \rightarrow 8 \text{ TeV}$

Model dependent, Real Extended Bialas-Bzdak theory results,
Odderon significance $\geq 7.08 \sigma$, from 1.96 and 2.76 TeV data only



Partially model independent, 2020-

Odderon Exchange from Elastic Scattering Differences between pp and $p\bar{p}$ Data at 1.96 TeV and from pp Forward Scattering Measurements

#1

TOTEM and D0 Collaborations • V.M. Abazov (Dubna, JINR) et al.
Published in: *Phys.Rev.Lett.* 127 (2021) 6, 062003 • e-Print: 2008.03001

Phys. Rev. Lett. **127** (2021) 6, 062003, Published: 4 August 2021
<https://doi.org/10.1103/PhysRevLett.127.062003>



S: Odderon significance $\geq 5.2\sigma$, C1: *almost* model independently combined with $\sqrt{s} = 13$ TeV data at $t = 0$: σ_{tot} and ρ_0

C2: one additional **pp dataset** at 8 TeV and one additional **data point** at 2.76 TeV,

C3: **8 out of the 17 D0 points** are used

C4: D0 $p\bar{p}$ data and TOTEM pp extrap.data are **assumed** to be **equal** at $t=0$

C5: ρ_0 (1.96 TeV) = 0.145

Some reflections on D0-TOTEM results

Odderon Exchange from Elastic Scattering Differences between pp and $p\bar{p}$ Data at 1.96 TeV and from pp Forward Scattering Measurements #1

TOTEM and D0 Collaborations • V.M. Abazov (Dubna, JINR) et al. (Dec 7, 2020)

Published in: *Phys.Rev.Lett.* 127 (2021) 6, 062003 • e-Print: 2012.03981 [hep-ex]




 pdf  links  DOI  cite


Phys. Rev. Lett. **127** (2021) 6, 062003, Published: 4 August 2021
<https://doi.org/10.1103/PhysRevLett.127.062003>

Lack of evidence for an odderon at small t #1

A. Donnachie (Manchester U.), P.V. Landshoff (Cambridge U.) (Mar 1, 2022)

Published in: *Phys.Lett.B* 831 (2022) 137199 • e-Print: 2203.00290 [hep-ph]




 pdf  DOI  cite


 3 citations

Coulomb-nuclear interference: Theory and practice for pp -scattering at 13 TeV #3


Vladimir A. Petrov (Serpuukhov, IHEP), Nikolai P. Tkachenko (Serpuukhov, IHEP) (Apr 19, 2022)

Published in: *Phys.Rev.D* 106 (2022) 5, 054003 • e-Print: 2204.08815 [hep-ph]

 pdf  DOI  cite

 0 citations

Back to Scaling: Model independently



$H(x|pp)$
s-independent:
2.76 – 7(8) TeV


$H(x|pp, 7 \text{ TeV})$
 \neq
 $H(x|p\bar{p}, 1.96)$

Odderon,
IF scaling holds
in pp down to
1.96 TeV
Domain of validity:
ReBB model
dependent

6.26 σ
Odderon effect


Energy range: tested **both** model independently and with modelling.
Modelling is useful, but model independent tests more important!

H(x) scaling of 2.76 and 8 TeV data




Energy range: H(x) scaling valid between $\sqrt{s} = 8$ and 2.76 TeV.
Uses final, published TOTEM $d\sigma/dt$ data at 8 TeV

H(x) scaling of 7 and 8 TeV data




Energy range: H(x) scaling model independently up to $\sqrt{s} = 8$ TeV.
Uses final, published TOTEM $\text{d}\sigma/\text{d}t$ data at 8 TeV

H(x) scaling of 7 and 8 TeV data



Closer look: systematic effects beyond the reported errors at dataset1 of 7 TeV (regarding type A and B ones, Type_C cancels)

H(x) scaling of 7 and 8 TeV data



Closer look: systematic effects beyond the reported errors at dataset1 of 7 TeV (regarding all types of errors, type_C cancels)


H(x) scaling of 7 and 8 TeV data

This is a first direct observation of systematics in the 7 TeV low $-t$ dataset beyond the reported errors.

Such problems were also seen in many earlier data analyses, but the problems were always attributed to the insufficiency of the methods applied in the analyses. A few examples:


- In this presentation: Model independent analysis
- Fagundes et al.,: Phys. Rev. D88, 094019
- Ster, Jenkovszky and Csörgő. Phys. Rev. D91, 074018; also presented in Bad Honnef at WE Heraeus Physics Scholl, 2015:
„Extracting the Odderon from pp and pp⁻ scattering data”

H(x): Odderon signal, new 8 TeV data



H(x) scaling is violated between $\sqrt{s} = 8$ TeV pp and 1.96 TeV pbarp.
Hungarian-Swedish Odderon signal confirmed with final, published
TOTEM $d\sigma/dt$ data at 8 TeV. Model independently.

$H(x|pbarp)/H(x|pp)$: Odderon peak



$H(x)$ scaling is violated between $\sqrt{s} = 8$ TeV pp and 1.96 TeV $pbarp$. TOTEM $d\sigma/dt$ data at 8 TeV. Odderon exchange, as a peak.


7 TeV: CLOSING DOORS/GATES

7 TeV data shifted

by $\epsilon_{B7,TeV}$ to minimize χ^2

Type A errors are shown only

Both swing and dip regions important!



7 TeV RESULTS, CLOSING GATES

Two sliding gates of size n and size m:
(n,m): Leaving out the first n and last m D0 point

Sliding door technique with two wings (n,m)

Left door excludes the first n, right door excludes the last m D0 points

	n	m	Odderon signal	Background		
	2	2	6.27 σ	1.68 σ		
	3	2	6.33 σ	1.70 σ		
	4	2	6.21 σ	2.37 σ		

MODEL INDEPENDENT RESULT 1:

In best window, optimized Odderon signal is 6.33 σ

MODEL INDEPENDENT RESULT 2:

Best window: leaving out first 3 and last 2 D0 point

MODEL INDEPENDENT RESULT 3:

Outside the best window: $H(x|pp) = H(x,p_{\text{bar}})$
pp and pbarp backgrounds agree within 1.7 σ

SUMMARY: ODDERON DISCOVERED IN 3 PAPERS, NEW: FOCUS ON ITS PROPERTIES

The H(x) analysis of the 8 TeV data CONFIRMED the existence Odderon. The united significance with the 7 and 8 TeV data, using the Stouffer's method is: 7.02σ

Odderon first discovered in three published papers: three different analysis, each with a statistical significance $> 5 \sigma$

0th property: Odderon exists!

Odderon properties: from Bialas-Bzdak model, so far valid in a limited s and $-t > 0.37 \text{ GeV}^2$ range only. 1.96 TeV – 8 TeV: Threshold effect, just appearing.

**There is an ongoing debate in reflective papers about the magnitude of the significance in the D0-TOTEM PRL
For exemple, is there enough evidence for Odderon at $t= 0$?**