

Two-loop calculations of the trilinear Higgs coupling in general renormalisable theories

Based on work in progress

in collaboration with Henning Bahl, Martin Gabelmann and Sebastian Paßehr

Johannes Braathen

SUSY 2024

IFT, Madrid, Spain | 13 June 2024

HELMHOLTZ RESEARCH FOR
GRAND CHALLENGES

DESY.





Why two loops ?

1) Capturing all types of effects entering λ_{hhh}

2) Properly interpreting experimental bounds on κ_λ

Mass-splitting effects in λ_{hhh} – how well established are they?



- First investigation of 1L BSM contributions to λ_{hhh} in 2HDM:
[Kanemura, (Kiyoura), Okada, Senaha, Yuan '02, '04]




- Deviations of tens/hundreds of % from SM possible, for large $g_{h\Phi\Phi}$ or $g_{hh\Phi\Phi}$ couplings
- Mass splitting effects, now found in various models (2HDM, inert doublet model, singlet extensions, etc.)

Mass-splitting effects in λ_{hhh} – how well established are they?


- First investigation of 1L BSM contributions to λ_{hhh} in 2HDM: [Kanemura, (Kiyoura), Okada, Senaha, Yuan '02, '04]



- Large effects **confirmed at 2L** in [JB, Kanemura '19] → **leading 2L corrections involving BSM scalars (H,A,H $^\pm$) and top quark**, computed in effective potential approximation



- Deviations of tens/hundreds of % from SM possible, for large $g_{h\Phi\Phi}$ or $g_{hh\Phi\Phi}$ couplings
- Mass splitting effects**, now found in various models (2HDM, inert doublet model, singlet extensions, etc.)



Scalar contributions to λ_{hhh} in aligned 2HDM

$$g_{hh\Phi\Phi} = -\frac{2(M^2 - m_\Phi^2)}{v^2}$$

BSM scalars:
 $\Phi \in \{H, A, H^\pm\}$
 $m_\Phi^2 = M^2 + \tilde{\lambda}_\Phi v^2$

Coupling/Order	0L	1L	2L	3L
g_{hhhh}			<i>subleading</i>	<i>subleading</i>
$g_{(h)h\Phi\Phi}$	-			
$g_{(h)H\Phi\Phi}$ [$g_{(h)G\Phi\Phi}$ case similar]	-	-		
$g_{\Phi\Phi\Phi'\Phi'}$	-	-		

[NB: 1 h can be replaced by a VEV!]

→ no further type of coupling entering after 2L

→ for each class of diagrams, perturbative convergence can be checked! [Bahl, JB, Weiglein PRL '22]

Why two loops ?


1) Capturing all types of effects entering λ_{hhh}

2) Properly interpreting experimental bounds on κ_λ

A benchmark scenario in the aligned 2HDM

[Bahl, JB, Weiglein PRL '22]

- **Two-Higgs-Doublet Model (2HDM):**
Here: CP conservation assumed, softly-broken Z_2 symmetry to avoid FCNCs, Yukawa couplings of type I
- Mass eigenstates:
 - 2 CP-even Higgs bosons
 h (125-GeV Higgs), H
 - CP-odd Higgs boson A
 - Charged Higgs bosons H^\pm
 - M : new mass term in 2HDM,
- Scenario with **alignment**: couplings of h are SM-like at tree level




A benchmark scenario in the aligned 2HDM

[Bahl, JB, Weiglein PRL '22]

Results shown for aligned 2HDM of type-I, similar for other types (*available in backup*)

We take $m_A = m_{H^\pm}$, $M = m_H$, $\tan\beta = 2$



- **Grey area:** area excluded by other constraints, in particular BSM Higgs searches, boundedness-from-below (BFB), perturbative unitarity
- **Light red area:** area excluded both by other constraints (BFB, perturbative unitarity) and by $\kappa_\lambda^{(2)} > 6.3$ [in region where $\kappa_\lambda^{(2)} < -0.4$ the calculation isn't reliable]
- **Dark red area:** new area that is **excluded ONLY by $\kappa_\lambda^{(2)} > 6.3$** . Would otherwise not be excluded!
- **Blue hatches:** area excluded by $\kappa_\lambda^{(1)} > 6.3 \rightarrow$ impact of including 2L corrections is significant!


A benchmark scenario in the aligned 2HDM

[Bahl, JB, Weiglein PRL '22]

Results shown for aligned 2HDM of type-I, similar for other types (*available in backup*)


We take $m_A = m_{H^\pm}$, $M = m_H$, $\tan\beta = 2$

2HDM type I, $M = m_H$, $m_A = m_{H^\pm}$, $\tan\beta = 2$, $\alpha = \beta - \pi/2$




- **Grey area:** area excluded by other constraints, in particular BSM Higgs searches, boundedness-from-below (BFB), perturbative unitarity
- **Light red area:** area excluded both by other constraints (BFB, perturbative unitarity) and by $\kappa_\lambda^{(2)} > 6.3$ [in region where $\kappa_\lambda^{(2)} < -0.4$ the calculation isn't reliable]
- **Dark red area:** new area that is **excluded ONLY** by $\kappa_\lambda^{(2)} > 6.3$. Would otherwise not be excluded!
- **Blue hatches:** area excluded by $\kappa_\lambda^{(1)} > 6.3$ → impact of including 2L corrections is significant!

Precise predictions for λ_{hhh} in arbitrary BSM theories




1) High-precision (2L) model-specific results
SM + singlet [JB, Kanemura '19]; 2HDMs [Senaha '18], [JB, Kanemura '19]; MSSM [Brucherseifer et al. '13]; NMSSM [Dao et al. '15], [Borschensky et al '22] ; models with classical scale invariance [JB, Kanemura, Shimoda '20], etc

Precise predictions for λ_{hhh} in arbitrary BSM theories



Precise predictions for λ_{hhh} in arbitrary BSM theories



1) High-precision (2L) model-specific results


SM + singlet [JB, Kanemura '19]; 2HDMs [Senaha '18], [JB, Kanemura '19]; MSSM [Brucherseifer et al. '13]; NMSSM [Dao et al. '15], [Borschensky et al '22]; models with classical scale invariance [JB, Kanemura, Shimoda '20], etc

3) High-precision (2L) generic predictions

[Bahl, JB, Gabelmann, Paßehr WIP]
→ [this talk!](#)

2) Generic 1L predictions with anyH3

[Bahl, JB, Gabelmann, Weiglein '23]
[Bahl, JB, Gabelmann, Radchenko Serdula, Weiglein WIP]
→ [c.f. talk in previous session](#)



Generic predictions for λ_{hhh} and λ_{hhhh} at two loops: our setup

Computing λ_{hhh} in general renormalisable theories: method

- All 2L contributions to Higgs/scalar self-energies computed in [Goodsell, Paßehr '19]
- In [Bahl, JB, Gabelmann, Paßehr to appear] we **generalise this with**
 - **new results for full 2L corrections to λ_{iii} and λ_{iiii} (with $p^2 = 0$)**
 - **results for 0-, 1-, 2-point functions in same conventions**
- Generic diagrams generated with FeynArts, computed with TwoCalc and OneCalc [Weiglein '93+]
- Generic results then mapped to specific models via private routines (using FeynArts diagrams and amplitudes)
- Topologies and diagrams reduced, taking benefit of symmetries → details in next slides

Computing λ_{hhh} in general renormalisable theories: method

- All 2L contributions to Higgs/scalar self-energies computed in [Goodsell, Paßehr '19]
- In [Bahl, JB, Gabelmann, Paßehr to appear] we **generalise this with**
 - **new results for full 2L corrections to λ_{iii} and λ_{iiii} (with $p^2 = 0$)**
 - **results for 0-, 1-, 2-point functions in same conventions**
- Generic diagrams generated with FeynArts, computed with TwoCalc and OneCalc [Weiglein '93+]
- Generic results then mapped to specific models via private routines (using FeynArts diagrams and amplitudes)
- Topologies and diagrams reduced, taking benefit of symmetries → details in next slides

- Application of generic results to concrete models:
(for genuine 2L diagrams, and 2L subloop renormalisation diagrams)

Computing λ_{hhh} in general renormalisable theories: method


- All 2L contributions to Higgs/scalar self-energies computed in [Goodsell, Paßehr '19]
- In [Bahl, JB, Gabelmann, Paßehr to appear] we **generalise this with**
 - **new results for full 2L corrections to λ_{iii} and λ_{iiii} (with $p^2 = 0$)**
 - **results for 0-, 1-, 2-point functions in same conventions**
- Generic diagrams generated with FeynArts, computed with TwoCalc and OneCalc [Weiglein '93+]
- Generic results then mapped to specific models via private routines (using FeynArts diagrams and amplitudes)
- Topologies and diagrams reduced, taking benefit of symmetries → details in next slides
- Application of generic results to concrete models:
(for genuine 2L diagrams, and 2L subloop renormalisation diagrams)

Diagrams generated
with FeynArts
[Hahn '01, '09 ++]

Computing λ_{hhh} in general renormalisable theories: method

- All 2L contributions to Higgs/scalar self-energies computed in [Goodsell, Paßehr '19]
- In [Bahl, JB, Gabelmann, Paßehr to appear] we **generalise this with**
 - **new results for full 2L corrections to λ_{iii} and λ_{iiii} (with $p^2 = 0$)**
 - **results for 0-, 1-, 2-point functions in same conventions**
- Generic diagrams generated with FeynArts, computed with TwoCalc and OneCalc [Weiglein '93+]
- Generic results then mapped to specific models via private routines (using FeynArts diagrams and amplitudes)
- Topologies and diagrams reduced, taking benefit of symmetries → details in next slides
- Application of generic results to concrete models:
(for genuine 2L diagrams, and 2L subloop renormalisation diagrams)


Diagrams generated
with FeynArts
[Hahn '01, '09 ++]



Reduced set
of diagrams


Computing λ_{hhh} in general renormalisable theories: method

- All 2L contributions to Higgs/scalar self-energies computed in [Goodsell, Paßehr '19]
- In [Bahl, JB, Gabelmann, Paßehr to appear] we **generalise this with**
 - **new results for full 2L corrections to λ_{iii} and λ_{iiii} (with $p^2 = 0$)**
 - **results for 0-, 1-, 2-point functions in same conventions**
- Generic diagrams generated with FeynArts, computed with TwoCalc and OneCalc [Weiglein '93+]
- Generic results then mapped to specific models via private routines (using FeynArts diagrams and amplitudes)
- Topologies and diagrams reduced, taking benefit of symmetries → details in next slides
- Application of generic results to concrete models:
(for genuine 2L diagrams, and 2L subloop renormalisation diagrams)




Computing λ_{hhh} in general renormalisable theories: method

- All 2L contributions to Higgs/scalar self-energies computed in [Goodsell, Paßehr '19]
- In [Bahl, JB, Gabelmann, Paßehr to appear] we **generalise this with**
 - **new results for full 2L corrections to λ_{iii} and λ_{iiii} (with $p^2 = 0$)**
 - **results for 0-, 1-, 2-point functions in same conventions**
- Generic diagrams generated with FeynArts, computed with TwoCalc and OneCalc [Weiglein '93+]
- Generic results then mapped to specific models via private routines (using FeynArts diagrams and amplitudes)
- Topologies and diagrams reduced, taking benefit of symmetries → details in next slides
- Application of generic results to concrete models:
(for genuine 2L diagrams, and 2L subloop renormalisation diagrams)



Computing λ_{hhh} in general renormalisable theories: method

- All 2L contributions to Higgs/scalar self-energies computed in [Goodsell, Paßehr '19]
- In [Bahl, JB, Gabelmann, Paßehr to appear] we generalise this with
 - new results for full 2L corrections to λ_{iii} and λ_{iiii}
 - results for 0-, 1-, 2-point functions in same context
- Generic diagrams generated with FeynArts, computed
- Generic results then mapped to specific models via p amplitudes)
- Topologies and diagrams reduced, taking benefit of s
- Application of generic results to concrete models: (for genuine 2L diagrams, and 2L subloop renormalis



```

    graph LR
      A[Diagrams generated with FeynArts  
[Hahn '01, '09 ++]] --> B[Reduced set of diagrams]
      B --> C[Analytic expressions in terms of T-integrals  
[Weiglein, Scharf, Böhm '93]]
      C --> D[Reduced analytic expressions in terms of 2L sunrise (T_134) and 1L integrals]
    
```


Reducing topologies and diagrams using symmetries

In our setting:


- External states are identical ($\rightarrow \lambda_{\text{iii}}, \lambda_{\text{iii}}$)
- All external momenta set to zero

} \rightarrow **many diagrams are identical**


For example, double box diagram,
with fermions and scalars:
 \rightarrow 6 topologies, 4 diagrams/topology




T1




T2




T3




T4




T5




T6




T1 G1 N1



T1 G2 N2





T1 G3 N3



T1 G4 N4

\rightarrow **Only need to compute 3 diagrams, rather than 24!**

Automating the reduction



Unique representation of diagram

→ "**canonical edges**"

- list of "edges" (= lines in diagram)
- identical diagrams ↔ permutations of edges
- canonical form = one particular choice of ordering

Reduction algorithm with "canonical edges" in pseudo code:

- identify internal and external indices
- generate permutations of external indices
- generate permutations of internal indices
- combine permutations of internal and external indices
- permute edge list following the combined list of permutations
- sort list of permuted edge lists
- return first element of sorted list of "edge lists"

```
edgelist1 = {edge[v[1], v[4], S[1]], edge[v[2], v[5], S[1]],
            edge[v[3], v[6], S[1]], edge[v[4], v[7], -F[3]],
            edge[v[4], v[8], F[3]], edge[v[5], v[6], F[3]],
            edge[v[5], v[8], -F[3]], edge[v[6], v[7], F[3]],
            edge[v[7], v[8], S[1]]}
edgelist2 = {edge[v[1], v[4], S[1]], edge[v[2], v[5], S[1]],
            edge[v[3], v[6], S[1]], edge[v[4], v[5], F[3]],
            edge[v[4], v[7], -F[3]], edge[v[5], v[8], F[3]],
            edge[v[6], v[7], F[3]], edge[v[6], v[8], -F[3]],
            edge[v[7], v[8], S[1]]}
```

Reducing topologies and diagrams in practice

n	topology-level	field-level
0	2 \rightarrow 2	11 \rightarrow 11
1	3 \rightarrow 3	25 \rightarrow 25
2	9 \rightarrow 8	121 \rightarrow 92 (102)
3	40 \rightarrow 13	936 \rightarrow 229 (291)
4	265 \rightarrow 29	10496 \rightarrow 698 (928)

n-point function

- Count **number of two-loop diagrams before and after reduction** of diagrams using canonical edges algorithm
 - ➔ At the level of topologies
 - ➔ At the level of field insertions

[Numbers in brackets \rightarrow for models with CP violation]

- **Reduction of up to one order of magnitude!**

NB: numbers shown here at generic level, not considering model-specific particle insertions nor summation over generation indices

Example applications


[Bahl, JB, Gabelmann, Paßehr to appear]

1) Cross-checks with existing results → today in SM



(in paper also for Z_2 SSM, 2HDM, NMSSM)

2) New results → today in singlet-extended SM

Leading two-loop $O(\alpha_s \alpha_t)$ and $O(\alpha_t^2)$ corrections in the SM



Reduction of 3-point diagrams, shown here at $O(\alpha_s \alpha_t)$



Leading two-loop $O(\alpha_s \alpha_t)$ and $O(\alpha_t^2)$ corrections in the SM

➤ Genuine 2L contributions at $O(\alpha_s \alpha_t)$

$$\Gamma_h^{(2)}|_{\mathcal{O}(\alpha_s \alpha_t)}^{\text{gen.}} = \frac{\alpha_s m_t^4}{\pi^3 v} \left[-\frac{3}{2\epsilon^2} + \frac{3\overline{\ln}m_t^2 - 2}{\epsilon} - 4 - \frac{\pi^2}{4} + 4\overline{\ln}m_t^2 - 3\overline{\ln}^2 m_t^2 \right],$$

$$\Gamma_{hh}^{(2)}(p^2 = 0)|_{\mathcal{O}(\alpha_s \alpha_t)}^{\text{gen.}} = \frac{\alpha_s m_t^4}{\pi^3 v^2} \left[-\frac{9}{2\epsilon^2} + \frac{9\overline{\ln}m_t^2}{\epsilon} - 4 - \frac{3\pi^2}{4} - 9\overline{\ln}^2 m_t^2 \right],$$

$$\Gamma_{hhh}^{(2)}(0, 0, 0)|_{\mathcal{O}(\alpha_s \alpha_t)}^{\text{gen.}} = \frac{\alpha_s m_t^4}{\pi^3 v^3} \left[-\frac{9}{\epsilon^2} + \frac{18}{\epsilon}(1 + \overline{\ln}m_t^2) - 8 - \frac{3\pi^2}{2} - 18\overline{\ln}m_t^2(2 + \overline{\ln}m_t^2) \right],$$

$$\Gamma_{hhhh}^{(2)}(0, 0, 0, 0)|_{\mathcal{O}(\alpha_s \alpha_t)}^{\text{gen.}} = \frac{\alpha_s m_t^4}{\pi^3 v^4} \left[-\frac{9}{\epsilon^2} + \frac{18}{\epsilon}(3 + \overline{\ln}m_t^2) - 80 - \frac{3\pi^2}{2} - 108\overline{\ln}m_t^2 - 18\overline{\ln}^2 m_t^2 \right]$$

and $O(\alpha_t^2)$

$$\Gamma_h^{(2)}|_{\mathcal{O}(\alpha_t^2)}^{\text{gen.}} = \frac{3m_t^6}{16\pi^4 v^3} \left[\frac{3}{4\epsilon^2} + \frac{1}{\epsilon} \left(2 - \frac{3}{2}\overline{\ln}m_t^2 \right) + \frac{9}{2} + \frac{5\pi^2}{24} - 4\overline{\ln}m_t^2 + \frac{3}{2}\overline{\ln}^2 m_t^2 \right],$$

$$\Gamma_{hh}^{(2)}(p^2 = 0)|_{\mathcal{O}(\alpha_t^2)}^{\text{gen.}} = \frac{3m_t^6}{16\pi^4 v^4} \left[\frac{9}{4\epsilon^2} + \frac{3}{\epsilon} \left(1 - \frac{3}{2}\overline{\ln}m_t^2 \right) + \frac{11}{2} + \frac{5\pi^2}{8} - 6\overline{\ln}m_t^2 + \frac{9}{2}\overline{\ln}^2 m_t^2 \right],$$

$$\Gamma_{hhh}^{(2)}(0, 0, 0)|_{\mathcal{O}(\alpha_t^2)}^{\text{gen.}} = \frac{3m_t^6}{16\pi^4 v^5} \left[\frac{9}{2\epsilon^2} - \frac{3}{\epsilon} \left(1 + 3\overline{\ln}m_t^2 \right) - 1 + \frac{5\pi^2}{4} + 6\overline{\ln}m_t^2 + 9\overline{\ln}^2 m_t^2 \right],$$

$$\Gamma_{hhhh}^{(2)}(0, 0, 0, 0)|_{\mathcal{O}(\alpha_t^2)}^{\text{gen.}} = \frac{3m_t^6}{16\pi^4 v^6} \left[\frac{9}{2\epsilon^2} - \frac{3}{\epsilon} \left(7 + 3\overline{\ln}m_t^2 \right) + 11 + \frac{5\pi^2}{4} + 42\overline{\ln}m_t^2 + 9\overline{\ln}^2 m_t^2 \right].$$

➤ Subloop renormalisation

$$\Gamma_h^{(2)}|_{\text{subloop}} = \frac{3m_t^2}{8\pi^2 v} \left[\frac{\delta_{\text{CT}}^{(1)} v^2}{v^2} - \delta_{\text{CT}}^{(1)} Z_h \right] \mathbf{A}_0(m_t^2)$$

$$- \frac{3m_t \delta_{\text{CT}}^{(1)} m_t}{2\pi^2 v} \left[\mathbf{A}_0(m_t^2) + m_t^2 \mathbf{B}_0(0, m_t^2, m_t^2) \right],$$

$$\Gamma_{hh}^{(2)}(p^2 = 0)|_{\text{subloop}} = \frac{3m_t^2}{4\pi^2 v^2} \left[\frac{\delta_{\text{CT}}^{(1)} v^2}{v^2} - \delta_{\text{CT}}^{(1)} Z_h \right] \left[\mathbf{A}_0(m_t^2) + 2m_t^2 \mathbf{B}_0(0, m_t^2, m_t^2) \right]$$

$$- \frac{3m_t \delta_{\text{CT}}^{(1)} m_t}{2\pi^2 v^2} \left[\mathbf{A}_0(m_t^2) + m_t^2 \left(5\mathbf{B}_0(0, m_t^2, m_t^2) + 4m_t^2 \mathbf{C}_0(m_t^2, m_t^2, m_t^2) \right) \right],$$

$$\Gamma_{hhh}^{(2)}(0, 0, 0)|_{\text{subloop}} = \frac{9m_t^4}{4\pi^2 v^3} \left[\frac{\delta_{\text{CT}}^{(1)} v^2}{v^2} - \delta_{\text{CT}}^{(1)} Z_h \right] \left[3\mathbf{B}_0(0, m_t^2, m_t^2) + 4m_t^2 \mathbf{C}_0(m_t^2, m_t^2, m_t^2) \right]$$

$$- \frac{18m_t^3 \delta_{\text{CT}}^{(1)} m_t}{\pi^2 v^3} \left[\mathbf{B}_0(0, m_t^2, m_t^2) + m_t^2 \left(3\mathbf{C}_0(m_t^2, m_t^2, m_t^2) \right. \right.$$

$$\left. \left. + 2m_t^2 \mathbf{D}_0(m_t^2, m_t^2, m_t^2, m_t^2) \right) \right],$$

$$\Gamma_{hhhh}^{(2)}(0, 0, 0, 0)|_{\text{subloop}} = \frac{9m_t^4}{\pi^2 v^4} \left[\frac{\delta_{\text{CT}}^{(1)} v^2}{v^2} - \delta_{\text{CT}}^{(1)} Z_h \right]$$

$$\times \left[\mathbf{B}_0(0, m_t^2, m_t^2) + 8m_t^2 \left(\mathbf{C}_0(m_t^2, m_t^2, m_t^2) + m_t^2 \mathbf{D}_0(m_t^2, m_t^2, m_t^2, m_t^2) \right) \right]$$


$$- \frac{18m_t^3 \delta_{\text{CT}}^{(1)} m_t}{\pi^2 v^4} \times \left\{ \mathbf{B}_0(0, m_t^2, m_t^2) + m_t^2 \left[13\mathbf{C}_0(m_t^2, m_t^2, m_t^2) \right. \right.$$

$$+ 4m_t^2 \left(7\mathbf{D}_0(m_t^2, m_t^2, m_t^2, m_t^2) \right.$$

$$\left. \left. + 4m_t^2 \mathbf{E}_0(m_t^2, m_t^2, m_t^2, m_t^2, m_t^2) \right) \right] \left. \right\},$$

→ combined with $\overline{\text{OS/MS}}$ counterterms, as desired

Leading two-loop $O(\alpha_s \alpha_t)$ and $O(\alpha_t^2)$ corrections in the SM



Results obtained for 2 electroweak schemes:

- G_F scheme
- $M_W/M_Z/\alpha_{em}$ scheme

[Difference is a 3L effect]

→ reproduces result of [JB, Kanemura '19]

→ new result


→ Remarkable agreement (to ~20 MeV)

New results: Leading two-loop corrections in the SSM

- SSM = Real-singlet extension of SM, without Z_2 symmetry

$$V_{\text{SSM}} = \mu^2 |\Phi|^2 + \frac{\lambda_H}{2} |\Phi|^4 + \frac{m_S^2}{2} S^2 + \frac{\kappa_S}{3} S^3 + \frac{\lambda_S}{2} S^4 + \kappa_{SH} S |\Phi|^2 + \frac{\lambda_{SH}}{2} S^2 |\Phi|^2$$

- Diagrams (reduced) for 2L contributions to λ_{hhh} from BSM scalar S:



New results: Leading two-loop corrections in the SSM


$$V_{\text{SSM}} = \mu^2 |\Phi|^2 + \frac{\lambda_H}{2} |\Phi|^4 + \frac{m_S^2}{2} S^2 + \frac{\kappa_S}{3} S^3 + \frac{\lambda_S}{2} S^4 + \kappa_{SH} S |\Phi|^2 + \frac{\lambda_{SH}}{2} S^2 |\Phi|^2$$

- › Neglect light Higgs mass before BSM scalar mass, $m_s \gg m_h$
- › Provide result in alignment limit (i.e. $\alpha = 0$), but subloop renormalisation still requires a counterterm for mixing angle α
- › Renormalisation scheme chosen as $\underbrace{m_h, m_s, \alpha, t_s, t_h, v}_{\text{on-shell}}, \underbrace{v_S, \kappa_S, \kappa_{SH}}_{\overline{\text{MS}}}$.

$$\begin{aligned} (4\pi)^4 \delta^{(2)} \lambda_{hhh}^{\text{OS}} &= -\frac{9\kappa_{SH}^3 v^3}{2v_S^5} - \frac{3\kappa_{SH}^3 v^3}{2m_s^2 v_S^4} \left[(\kappa_S + 2\kappa_{SH}) \overline{\ln} m_s^2 - 2(\kappa_S - \kappa_{SH}) - 3\kappa_{SH} \frac{v^2}{v_S^2} \right] \\ &\quad - \frac{\kappa_{SH}^3 v^3}{8m_s^4 v_S^3} \left[4\kappa_S^2 + \kappa_{SH} (5\kappa_{SH} - 12\kappa_S) \frac{v^2}{v_S^2} + 9\kappa_{SH}^2 \frac{v^4}{v_S^4} \right], \\ &\approx -\frac{9\kappa_{SH}^3 v^3}{2v_S^5} + \mathcal{O}\left(\frac{m_h^2}{m_s^2}, \frac{\kappa_{SH}^2}{m_s^2}, \frac{\kappa_S^2}{m_s^2}\right) \end{aligned}$$

New results: Leading two-loop corrections in the SSM

Numerical results, in decoupling limit $m_s \rightarrow \infty$



→ inclusion of 2L corrections significantly reduces the renormalisation-scale dependence

Summary

- λ_{hhh} plays a crucial role to understand the **shape of the Higgs potential**, and probe indirectly **signs of New Physics**
- Two-loop corrections to λ_{hhh} can be significant, even for points allowed by theoretical (esp. perturbative unitarity) and experimental constraints → **inclusion of two-loop corrections is important for reliable interpretation of bounds on κ_λ**
- **Full two-loop corrections to λ_{hhh} and λ_{hhhh} computed for general renormalisable theories**
 - Generic results mapped onto diagrams generated by FeynArts for model(s) to consider
 - Extensive cross-checks performed, and new results obtained for λ_{hhh}
 - EFT-like matching of λ_{hhhh} also possible
- Many applications ongoing, countless more possible!

Thank you very much for your attention!

Contact

DESY. Deutsches
Elektronen-Synchrotron

Johannes Braathen
DESY Theory group


www.desy.de

johannes.braathen@desy.de


Backup

Experimental probes of λ_{hhh}

- **Double-Higgs production** $\rightarrow \lambda_{hhh}$ enters at leading order (LO) \rightarrow **most direct probe!**




- **Single-Higgs production** $\rightarrow \lambda_{hhh}$ enters at NLO



[Degrassi, Giardino, Maltoni, Pagani '16] [ATLAS-CONF-2019-049]


- **Electroweak Precision Observables (EWPOs)** $\rightarrow \lambda_{hhh}$ enters at NNLO



[Degrassi, Fedele, Giardino '17]

Accessing λ_{hhh} experimentally

- Double-Higgs production $\rightarrow \lambda_{hhh}$ enters at LO \rightarrow **most direct probe of λ_{hhh}**



- Box and triangle diagrams **interfere destructively**


\rightarrow small prediction in SM

\rightarrow BSM deviation in λ_{hhh} can **significantly alter double-Higgs production!**

- Upper limit on double-Higgs production cross-section

\rightarrow **limits on $\kappa_\lambda \equiv \lambda_{hhh}/(\lambda_{hhh}^{(0)})^{SM}$**

- κ_λ as an effective coupling $\rightarrow \mathcal{L} \supset -\kappa_\lambda \times \frac{3m_h^2}{v^2} \cdot h^3 + \dots$



[Frederix et al., '14]

Accessing λ_{hhh} via double-Higgs production

- Double-Higgs production $\rightarrow \lambda_{hhh}$ enters at LO \rightarrow most direct probe of λ_{hhh}

Recent results from ATLAS hh-searches [ATLAS-CONF-2022-050] yield the limits:

$$-0.4 < \kappa_\lambda < 6.3 \text{ at 95\% C.L.}$$

\rightarrow factor ~ 2 improvement compared to


pre-2021 best ATLAS limits (from single-h prod.)

$$-3.2 < \kappa_\lambda < 11.9 \text{ at 95\% C.L. [ATLAS-PHYS-PUB-2019-009]}$$

(CMS recently gave $-1.2 < \kappa_\lambda < 6.5$ at 95% C.L. [CMS '22])

\rightarrow Can κ_λ now be used to constrain the parameter space of BSM models?

- κ_λ as an effective coupling $\rightarrow \mathcal{L} \supset -\kappa_\lambda \times \frac{\partial \ln v_h}{v^2} \cdot h^3 + \dots$



[Frederix et al., '14]

Future determination of λ_{hhh}

Expected sensitivities in literature, assuming $\lambda_{hhh} = (\lambda_{hhh})^{SM}$

di-Higgs exclusive result


Higgs@FC WG September 2019

di-Higgs	single-Higgs
HL-LHC 50%	HL-LHC 50% (47%)
HE-LHC [10-20]%	HE-LHC 50% (40%)
FCC-ee/eh/hh 5%	FCC-ee/eh/hh 25% (18%)
LE-FCC 15%	LE-FCC n.a.
FCC-eh ₃₅₀₀ -17+24%	FCC-eh ₃₅₀₀ n.a.
	FCC-ee ^{4IP} ₃₆₅ 24% (14%)
	FCC-ee ₃₆₅ 33% (19%)
	FCC-ee ₂₄₀ 49% (19%)
ILC ₁₀₀₀ 10%	ILC ₁₀₀₀ 36% (25%)
ILC ₅₀₀ 27%	ILC ₅₀₀ 38% (27%)
	ILC ₂₅₀ 49% (29%)
	CEPC 49% (17%)
CLIC ₃₀₀₀ -7+11%	CLIC ₃₀₀₀ 49% (35%)
CLIC ₁₅₀₀ 36%	CLIC ₁₅₀₀ 49% (41%)
	CLIC ₃₈₀ 50% (46%)

All future colliders combined with HL-LHC

single-Higgs exclusive

single-Higgs global



Plot taken from
[de Blas et al., 1905.03764]

see also [Cepeda et al., 1902.00134], [Di Vita et al.1711.03978], [Fujii et al. 1506.05992, 1710.07621, 1908.11299], [Roloff et al., 1901.05897], [Chang et al. 1804.07130,1908.00753], etc.

Future determination of λ_{hhh}

Higgs production cross-sections (here double Higgs production) depend on λ_{hhh}





Figure 10. Double Higgs production at hadron (left) [65] and lepton (right) [66] colliders as a function of the modified Higgs cubic self-coupling. See Table 18 for the SM rates. At lepton colliders, the production cross sections do depend on the polarisation but this dependence drops out in the ratios to the SM rates (beam spectrum and QED ISR effects have been included).

Plots taken from
[de Blas et al., 1905.03764]

[Frederix et al.,
1401.7340]

Experimental situation for λ_{hhh}

- Double-Higgs production $\rightarrow \lambda_{hhh}$ enters at LO \rightarrow **most direct probe of λ_{hhh}**




[Note: Single-Higgs production (EW precision observables) $\rightarrow \lambda_{hhh}$ enters at NLO (NNLO)]

- Box and triangle diagrams **interfere destructively**
 \rightarrow small prediction in SM
 \rightarrow BSM deviation in λ_{hhh} can **significantly enhance double-Higgs production!**

- Search limits on double-Higgs production
 \rightarrow **limits on effective coupling $\kappa_\lambda \equiv \lambda_{hhh} / (\lambda_{hhh}^{(0)})^{SM}$**


$$-0.4 < \kappa_\lambda < 6.3$$

[ATLAS PLB '23]




2HDM benchmark plane – individual theoretical constraints


Constraints shown below are independent of 2HDM type




Boundedness from below



EW precision observables computed at 2L




$\kappa_\lambda^{(2)} > 6.6$



Perturbative unitarity at (N)LO


2HDM benchmark plane – experimental constraints

i.e. Higgs physics (via HiggsBounds and HiggsSignals) and b physics (from [Gfitter group 1803.01853])



2HDM benchmark plane – experimental constraints

i.e. Higgs physics (via HiggsBounds and HiggsSignals) and b physics (from [Gfitter group 1803.01853])



2HDM benchmark plane – results for all types

