SUSY24: The 31st International Conference on Supersymmetry and Unification of Fundamental Interactions



Contribution ID: 84

Type: Parallel Talk

## Exploring the effects of leptoquarks in the LFV muon decays

Neutrino oscillation in the matter could be affected by the sub-dominant, yet unknown, non-standard interactions (NSI). The observed shift in  $\delta_{CP}$  value observed for NOvA in case of standard model (SM) and NSIs arising simultaneously from two different off-diagonal sectors,  $e_{-\mu}$  and  $e_{-\tau}$  could be attributed to the beyond standard model physics. We derive constraints on the NSI sectors using the combined datasets of NOvA and T2K. Our analysis reveals a significant impact that dual NSIs may have on the sensitivity of standard CP phase  $\delta_{CP}$  and atmospheric mixing angle  $\theta_{23}$ . Here we assume that the presence of leptoquarks ( $U_3$ ) to account for the apparent difference in the experimental observations of  $\delta_{CP}$  measurement by NOvA and T2K using  $\epsilon_{e\mu}$ . Furthermore, using this input we obtain the Lepton Flavour violation (LFV) muon decay decay branching ratios:  $B(\mu \rightarrow e\gamma) \le 10^{-18}$ ,  $B(\mu \rightarrow eee) \le 10^{-21}$  and  $B(\mu \rightarrow e)$ Ti $\le 10^{-19}$  which could be probed in the future experiments.

**Primary authors:** GIRI, Anjan (IIT Hyderabad); BRAHMA, Barnali (IIT Hyderabad); NAYAK, Lopamudra (IIT Hyderabad)

Presenter: GIRI, Anjan (IIT Hyderabad)

Session Classification: Flavour physics and neutrinos

Track Classification: Flavour physics and neutrinos