

Detecting Rare Species of Dark Matter with Terrestrial Detectors


- i) Phys. Rev. Lett. 131, 011005 (2023) [[arXiv: 2303.03416](#)]
- ii) JCAP 01 029 (2024) [[arXiv: 2309.10032](#)]
- iii) [arXiv: 2402.03431](#)

Anupam Ray


N3AS Fellow, UC Berkeley & University of Minnesota

DPF-PHENOM 2024


05.13.2024



Dark Matter (DM)



From: Bertone and Hooper,
Rev. Mod. Physics (2016)




https://wmap.gsfc.nasa.gov/universe/uni_matter.html

- DM mass?
- DM interactions with baryons?


Strongly-interacting DM Component

- A sub-component of DM can be strongly interacting.



χ makes up a sub-component
of the total DM energy budget.

Strongly-interacting DM Component



Take Away

- “Earth-bound” DM provides a novel powerful probe.

The density of “Earth-bound DM” can be huge, 15 orders of magnitude larger than the Galactic DM density!

Annihilating DM

- Local annihilation inside any large-volume neutrino detectors (such as Super-Kamiokande)

Ray, (with McKeen, Morrissey, Pospelov, Ramani) [PRL, 2023]

- Neutrinos from annihilation of Earth-bound DM.

Pospelov & Ray [JCAP, 2024]

Non-Annihilating DM

- Earth-bound DM can be up-scattered by fast neutrons inside the nuclear reactors, and subsequently detected.



Ray, (with Ema, Pospelov) [2402.03431]

Earth-Bound DM


Press & Spergel (1985,ApJ), Gould (1987, ApJ),...

Small $\sigma_{\chi n} \rightarrow$ single collision,

large $\sigma_{\chi n} \rightarrow$ multiple collisions.



Earth-Bound DM



$$f_c(\sigma_{\chi n}, m_\chi)$$

Earth-Bound DM

- Lets do some estimate:

For DM mass of 1 GeV and $\sigma_{\chi n} = 10^{-28} \text{ cm}^2$

$$C_{\text{geo}} = 1.3 \times 10^{25} \text{ s}^{-1} \quad \text{and} \quad f_c \sim 0.1$$

$f_\chi = 1$


DM density (assuming they uniformly distribute over the Earth-volume)

$$\rho_\chi = m_\chi \frac{f_c \times C_{\text{geo}} \times t_\oplus}{V_\oplus} \sim 3 \times 10^{14} \text{ GeV/cm}^3$$

$f_\chi = 1$

- 15 orders of magnitude larger than the Galactic DM density!

DM Distribution in Stellar Objects



- Dimensionless profile function:

$$G_\chi(R_\oplus) = \frac{n_\chi(R_\oplus)V_\oplus}{N_\chi}$$

- For uniform DM density:

$$G_\chi(R_\oplus) = 1$$

Signal at Super-K

- Earth-bound DM, of mass GeV scale have an enormously large surface density.
- Their detection via scattering is **almost impossible** as they acquire very little amount kinetic energy (0.03 eV).
- How to detect them?


Ray, (with McKeen, Morrissey, Pospelov, Ramani) [PRL, 2023]

Our proposal: simply look at their annihilation signature inside large-volume detectors (annihilation is not limited to the tiny kinetic energy)!

Results

- Using existing di-nucleon annihilation searches at Super-K

Ray, (with McKeen, Morrissey, Pospelov, Ramani) [PRL, 2023]




←
Evaporation

..... →
DM shrinks towards
the Earth-core

Up to $f_\chi = 10^{-10}$


Results

Ray, (with McKeen, Morrissey, Pospelov, Ramani) [PRL, 2023]




Evaporation

DM shrinks towards
the Earth-core



Results

Ray, (with McKeen, Morrissey, Pospelov, Ramani) [PRL, 2023]




χ : Dirac fermion which can couple to a dark photon A'

$$\chi \bar{\chi} \rightarrow A' A'$$


$$A' \rightarrow \text{SM} + \text{SM} (\text{say } e^+ + e^-)$$

Unprecedented sensitivity on parts of the parameter space.

What about heavy DM?



(Can not be improved)



DM shrinks towards
the Earth-core
(Can be improved)

Neutrino Signal

- Earth-bound DM if sufficiently heavy, shrinks towards the core, leading to a negligible surface density.
gravity dominates over the diffusion processes
- Annihilation to neutrinos can occur at the Earth-core, if Earth-bound DM if sufficiently heavy. Since the number density is huge, annihilation rate is also fairly large.
- Neutrinos, because of their feeble interactions, can reach detectors like Super-K, IceCube-DeepCore, and searching these annihilated neutrinos can provide sensitivity to DM interactions.


Pospelov & Ray [JCAP, 2024]

Neutrino Signal

- We consider two phenomenological scenarios:


Lower energy neutrinos from the stopped pion decay

Higher energy neutrino lines from direct annihilation




Low Energy Neutrinos

Super-Kamiokande (APJL, 2023)



Low Energy Neutrinos

Pospelov & Ray [JCAP, 2024]



We use the Super-K DSNB search result with 0.01 wt% gadolinium loaded water ($22.5 \text{ kton} \times 552.2 \text{ days}$) to derive the exclusion limits

Super-Kamiokande (APJL, 2023)

*Gd-loaded water gives competitive limit (as compared to the pure-water limits) although the data is 5 times less.




High Energy Neutrinos

- DM annihilation directly to neutrinos yields a line at $E_\nu = m_\chi$
high-energy neutrinos can also come from $\chi\chi \rightarrow W^+W^-$, $b\bar{b}$, $\tau\bar{\tau}$, giving a continuum spectra up to $E_\nu = m_\chi$ (or $\chi\chi \rightarrow A'A' \rightarrow 4\nu$).
- We search the “neutrino-line” signature in the IceCube DeepCore data with a total live-time of 6.75 years.
- We use the null-detection of the neutrino-line signature in the IceCube DeepCore data to derive the exclusions

Mass (GeV)	$b\bar{b}$ $\Gamma_{\text{ann}} [\text{s}^{-1}] \times 10^{23}$	$\tau\bar{\tau}$ $\Gamma_{\text{ann}} [\text{s}^{-1}] \times 10^{23}$	$\nu\bar{\nu}$ $\Gamma_{\text{ann}} [\text{s}^{-1}] \times 10^{23}$
5	139	139.3	
10	396	7.0	1.37
20	29.7	0.97	0.27
35	7.41	0.22	0.09
50	3.51	0.096	0.05
100	1.39	0.038	0.027

High Energy Neutrinos

Pospelov & Ray [JCAP, 2024]



We probe up to $f_\chi \geq 10^{-8}$ for sufficiently heavy Earth-bound DM.

Summary


- Earth accumulates significant number of DM particles from the Galactic halo, leading to a DM density **15 orders of magnitude larger** than the Galactic DM density!
- Despite their prodigious abundance, their detection is extremely challenging as they acquire **tiny amount** of kinetic energy.
- **Annihilation** of such Earth-bound DM at large-volume neutrino detectors, provides a novel way for their detection and can be used to probe strongly-interacting DM component.

Conclusion

★ How to detect rare species of DM?



★ Look at the
Earth-bound DM!



Thanks!

Questions & Comments: anupam.ray@berkeley.edu