



George W.S. Hou (侯維恕)
National Taiwan University

May 14, 2023, DPF-PHENO @ U. Pittsburgh



National Taiwan University



0. Our current *impasse*: *No New Physics* (*NNP*)

(w/o Z_2)

5 Merits

I. General Two-Higgs Doublet Model (G2HDM)

More *Dim-4*'s (two extra sets of couplings) → **Don't EFT yet!**

II. Decadal Mission of the *New Higgs/Flavor* Era

Midterm Report "my view for *BSM*": $pp \rightarrow t\bar{t}(\text{bar})$ ATLAS & CMS

III. Post-Midterm: $pp \rightarrow bH^+ \rightarrow b\bar{t}(\text{bar}); t\bar{t}(\text{bar})$ @ CMS

[$t \rightarrow ch$ & $t\bar{t}(\text{bar})$ *redux*]

IV. G2HDM as *Next NP?!?*

V. Discussion & Conclusion



Physicists' Nightmare Scenario: The Higgs and Nothing Else

Adrian Cho


Many fear the LHC will cough up only the one particle they've sought for decades.

Some would rather see nothing new at all

fear nothing more than the possibility that you were wrong and the particle doesn't exist, right? Not exactly.

Many particle physicists say their greatest fear is that their grand new machine—the Large Hadron Collider (LHC) under construction at the European particle physics laboratory, CERN, near Geneva, Switzerland—will spot the Higgs boson and nothing else. If so, particle physics could grind to halt, they say. In fact, if the LHC doesn't reveal a plethora of new particles in addition to the Higgs, many say they would rather it see nothing new at all.

That may seem perverse, but put yourself again in the shoes of a particle physicist. In the 1960s and 1970s, researchers hammered out a theory called the Standard Model that, in



SCIENCE VOL 315 23 MARCH 2007



Physicists' Nightmare Scenario: The Higgs and Nothing Else


Adrian Cho

Many fear the LHC will cough up only the one particle they've sought for decades. Some would rather see nothing new at all


fear nothing more than the possibility that you were wrong and the particle doesn't exist, right? Not exactly.

Many particle physicists say their greatest fear is that their grand new machine—the Large Hadron Collider (LHC) under construction at the European particle physics laboratory, CERN, near Geneva, Switzerland—will spot the Higgs boson and nothing else. If so, particle physics could grind to halt, they say. In fact, if the LHC doesn't reveal a plethora of new particles in addition to the Higgs, many say they would rather it see nothing new at all.

That may seem perverse, but put yourself again in the shoes of a particle physicist. In the 1960s and 1970s, researchers hammered out a theory called the Standard Model that, in



SCIENCE VOL 315 23 MARCH 2007



If it has the right mass, the Higgs and nothing else **“would be the real five-star disaster, because that would mean there wouldn't need to be any new physics.”**

—Jonathan Ellis, CERN

Ten years after the Higgs, physicists face the nightmare of finding nothing else



Adrian Cho

Unless Europe's Large Hadron Collider coughs up a surprise, the field of particle physics may wheeze to its end

13 JUN 2022 • 1:30 PM • BY ADRIAN CHO

CPV

Merit

M①: extra top Yukawas ρ_{tt} and $\rho_{tc} \sim 1$ and **complex**, can drive **EWBG**,


Fuyuto, WSH, Senaha PLB'18

Higgs quartic self-couplings η_i at $\mathcal{O}(1)$, $i = 1-7$, provide **1stOPhTr** (\rightarrow primordial GW!)

Kanemura, Okada, Senaha, PLB'05

M②: **CPV @ $\mathcal{O}(1)$** needed for **EWBG** \rightarrow vulnerable to **e EDM** (ACME'18 & JILA'23)

\rightarrow Spectacular 2-loop diagrammatic cancellation



Fuyuto, Hou, Senaha PRD'20 (R)

Higgs- γ - γ^* insertions

$\rightarrow |\rho_{ee}/\rho_{tt}| \sim \lambda_e/\lambda_t$
 the flavor code?

CPV

Merit

M①: extra top Yukawas ρ_{tt} and $\rho_{tc} \sim 1$ and **complex**, can drive **EWBG**,


Fuyuto, WSH, Senaha PLB'18

Higgs quartic self-couplings η_i at $\mathcal{O}(1)$, $i = 1-7$, provide **1stOPhTr** (\rightarrow primordial GW!)

Kanemura, Okada, Senaha, PLB'05

M②: **CPV @ $\mathcal{O}(1)$** needed for **EWBG** \rightarrow vulnerable to **e EDM** (ACME'18 & JILA'23)

\rightarrow Spectacular 2-loop diagrammatic cancellation



Fuyuto, Hou, Senaha PRD'20 (R)

Glashow-Weinberg PRD'77

M③: Glashow worried about **FCNCs**, such as $t \rightarrow ch$; but with $h < t$, it is a "PDG" duty to search!

Higgs- γ - γ^* insertions

$\rightarrow |\rho_{ee}/\rho_{tt}| \sim \lambda_e/\lambda_t$

the flavor code?

Curiously, $t \rightarrow ch$ remains elusive

— Nature threw in **alignment** (*small* h - H mixing)

to hide it so far! Who would have thought?!

WSH, PLB'92
 flavor-protected

CPV

Merit

M①: extra top Yukawas ρ_{tt} and $\rho_{tc} \sim 1$ and **complex**, can drive **EWBG**,


Fuyuto, WSH, Senaha PLB'18

Higgs quartic self-couplings η_i at $\mathcal{O}(1)$, $i = 1-7$, provide **1stOPhTr** (\rightarrow primordial GW!)

Kanemura, Okada, Senaha, PLB'05

M②: **CPV @ $\mathcal{O}(1)$** needed for **EWBG** \rightarrow vulnerable to **eEDM** (ACME'18 & JILA'23)

\rightarrow Spectacular 2-loop diagrammatic cancellation



Fuyuto, Hou, Senaha PRD'20 (R)

Higgs- γ - γ^* insertions

$$\rightarrow |\rho_{ee}/\rho_{tt}| \sim \lambda_e/\lambda_t$$

the flavor code?

Glashow-Weinberg PRD'77

M③: Glashow worried about **FCNCs**, such as $t \rightarrow ch$; but with $h < t$, it is a "PDG" duty to search!

Curiously, $t \rightarrow ch$ remains elusive

— Nature threw in **alignment** (small c_γ h-H mixing) to hide it so far! Who would have thought?!

WSH, PLB'92
 flavor-protected

sub-TeV
 H, A, H⁺

M④: Small c_γ does *Not* contradict **$\mathcal{O}(1)$ quartics**:

\rightarrow Can argue that **H, A, H⁺ populate 300–600 GeV**.

$$c_\gamma \sim \frac{\eta_6 v^2}{m_H^2 - m_h^2}$$

WSH, Kikuchi EPL'18

M⑤: With $t \rightarrow ch$ c_γ -suppressed \rightarrow Natural to pursue $cg \rightarrow tH/tA \rightarrow ttc(\text{bar})$

Kohda, Modak, WSH PLB'18



\rightarrow Better: $cg \rightarrow bH^+ \rightarrow btb(\text{bar})$ [recoil b, not t]

Ghosh, WSH, Modak PRL'20



Same-sign top pair + jet

→ Natural to pursue $cg \rightarrow tH/tA \rightarrow ttc(\text{bar})$



$A/H \rightarrow tc(\text{bar})$



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: July 28, 2023
ACCEPTED: October 5, 2023
PUBLISHED: December 12, 2023

Search for heavy Higgs bosons with flavour-violating couplings in multi-lepton plus *b*-jets final states in *pp* collisions at 13 TeV with the ATLAS detector



The ATLAS collaboration

2307.14759

Phys. Lett. B 850 (2024) 138478



ELSEVIER

Contents lists available at ScienceDirect

Physics Letters B

journal homepage: www.elsevier.com/locate/physletb



Letter

Search for new Higgs bosons via same-sign top quark pair production in association with a jet in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$

The CMS Collaboration* A. Hayrapetyan et al.

2311.03261





* NTUCMS started already 2/2020, but thanks to a good 5-yr grant, manpower could be built-up in time since 8/2021.



Same-sign top pair + jet

→ Natural to pursue $cg \rightarrow tH/tA \rightarrow ttc(\text{bar})$



A/H → tc(bar)

PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: July 28, 2023
ACCEPTED: October 5, 2023
PUBLISHED: December 12, 2023

Search for heavy Higgs bosons with flavour-violating couplings in multi-lepton plus *b*-jets final states in *pp* collisions at 13 TeV with the ATLAS detector

The ATLAS collaboration

Neither ATLAS
2307.14759

Phys. Lett. B 850 (2024) 138478

Contents lists available at ScienceDirect

Physics Letters B

journal homepage: www.elsevier.com/locate/physletb

Letter

Search for new Higgs bosons via same-sign top quark pair production in association with a jet in proton-proton collisions at $\sqrt{s} = 13$ TeV

The CMS Collaboration * A. Hayrapetyan et al.

2311.03261

Neither ATLAS ... **Nor** CMS saw a signal.


* NTUCMS started already 2/2020, but thanks to a good 5-yr grant, manpower could be built-up in time since 8/2021.

III. Post-Midterm: $pp \rightarrow bH^+ \rightarrow b\bar{t}(\bar{b}); t\bar{t}(\bar{b}) @ CMS$ [$t\bar{t}(\bar{b})$ & $t \rightarrow ch$ redux]




→ Better: $cg \rightarrow bH^+ \rightarrow b\bar{t}(\bar{b})$

Ghosh, WSH, Modak PRL'20



V_{tb}/V_{cb} enhanced
 w.r.t. 2HDM-II



$H^+ \rightarrow t\bar{b}$ $\rho_{tt} V_{tb}$



also: $cg \rightarrow tH/tA \rightarrow t\bar{t}(\bar{b}), t\bar{t}(\bar{b}) [H/A \rightarrow t\bar{t}(\bar{b})]$
 redux adding Run 3 data
 t → ch

The elevated current H, A, H⁺ search program @ CMS.



III. Post-Midterm: $pp \rightarrow bH^+ \rightarrow btb(\bar{)}; ttt(\bar{)} @ CMS$ [ttc(bar) & $t \rightarrow ch$ redux]

→ Better: $cg \rightarrow bH^+ \rightarrow btb(\bar{)}$



also: $cg \rightarrow tH/tA \rightarrow ttc(\bar{)}, ttt(\bar{)} [H/A \rightarrow tt(\bar{)}]$
 redux adding Run 3 data
 $t \rightarrow ch$

Ghosh, WSH, Modak PRL'20

V_{tb}/V_{cb} enhanced
 w.r.t. 2HDM-II

The elevated current H, A, H+ search program @ CMS.

IV. G2HDM as *Next NP?!*

WSH, Kohda, Modak, Wong PLB'20

The above H+ CKM enhancement first uncovered in $B \rightarrow \mu\nu, \tau\nu$; ratio $\neq 0.0045$

→ Rule out SM/2HDM-II.


Belle II will study this.

CMS: $B_{s,d} \rightarrow \mu\mu$

$\tau \rightarrow \mu\gamma$ induced by 2-loop diagrams.

$B \rightarrow \mu\nu$: stat. dom. ← takes time

$B \rightarrow \tau\nu$: syst. dom. ← imprv method




V_{tb}/V_{ub} enhanced!

WSH & Kumar PRD'20

Glimpse of the *New Flavor Era*

WSH "Decadal Mission" 2109.02557, CJP'22



Rule of Thumb

$$\rho_{3j}^f \lesssim \lambda_3^f \quad (j \neq 1),$$

$$\rho_{i1}^f \lesssim \lambda_1^f$$

Flavor Control
 “Decadal Mission”
 CJP’22



Thus, our Decadal Mission:

“*Find* the *extra H, A, H⁺ bosons*;

Crack the *Flavor code*;

*Solve** the *Mysterious B.A.U.!*”


Is this it?!

$|P_{ee}/P_{\mu\mu}| \sim \lambda_e/\lambda_t!$
the flavor code?

* We are also conducting a Lattice study of $\mathcal{O}(1)$ quartics for 1stOPhTr → New Scale?


Up to *Nature* whether our “Wish for *Discovery*” is Granted ... or Not ...

Thank You!



a Higgs, and a 2nd Higgs ...

Glimpse of coming **New Flavor Era** in G2HDM



WSH&Kumar, PRD'20
EPIC'21



EWBG ought to be pursued while LHC is still running!

No SUSY, No Nothin'!

Beyond CKM CPV




EW BaryoGenesis (EWBG)

- more testable -



LHC
- No New Physics -

Fuyuto, WSH, Senaha, PRD 101 (2020) 011901(R)



JILA'22
 e EDM: ACME14 → ~~ACME18~~

- L.E. Precision Frontier -

0.41
 $|d_e| < 1.1 \times 10^{-29} \text{ e cm}$

A Natural Cancellation Mechanism!

Soaring to the Starry Heavens

Baryon **A**symmetry of **U**niverse

Enough **CPV** for **BAU**?

$$\lambda_t \text{Im} \rho_{tt}$$

天 **CPV** → **BAU**

Where have all the **Antimatter** Gone?
BAU: an issue as **Big** as the **Universe**.

Billions & billions of **stars** ... all **burning** protons

le Raison d'être

Search H, A, H^+ @ LHC!

地



JILA'22

ACME experiment: Current frontline, Probe **CPV** via **eEDM**, put check on **Baryogenesis**.

Great Reef National Park (c) Wally Pacholka




h(125): observed 7/4/2012

λ_f : Yukawa Couplings

Expt'lly Affirmed!

35.9-137 fb⁻¹ (13 TeV)



m_e ?

$$g \approx 2m_V/v$$

ca. 2015

$$\lambda_f \approx \sqrt{2}m_f/v$$

t/b/ τ : 2018


μ : 2020

Higgs "potential": Simplest!!

$$V(\Phi) \sim -|\mu|^2|\Phi|^2 + \lambda|\Phi|^4$$

$$\Rightarrow |\phi^0|^2 = v^2 \sim \mu^2/\lambda$$

$v \approx 246$ GeV




nobelprize.org (2013)



- Extra Higgs Doublet w/
- Extra Yukawa Couplings
- Extra Quartic Couplings

General 2HDM



Before Top Quark was Discovered: WSH, PLB'92 (PSI-PR-91-34)

Correction on formulation with “SM”-like Higgs h: Chen, WSH, Kao, Kohda, PLB'13

$$\begin{pmatrix} H^+ \\ H + iA \end{pmatrix}$$

PHYSICAL REVIEW LETTERS **129**, 032001 (2022)

Search for Flavor-Changing Neutral Current Interactions of the Top Quark and Higgs Boson in Final States with Two Photons in Proton-Proton Collisions at $\sqrt{s} = 13$ TeV

A. Tumasyan *et al.**
(CMS Collaboration)



(Received 3 November 2021; accepted 13 June 2022; published 13 July 2022)

Proton-proton interactions resulting in final states with two photons are studied in a search for the signature of flavor-changing neutral current interactions of top quarks (t) and Higgs bosons (H). The analysis is based on data collected at a center-of-mass energy of 13 TeV with the CMS detector at the LHC, corresponding to an integrated luminosity of 137 fb^{-1} . No significant excess above the background prediction is observed. Upper limits on the branching fractions (\mathcal{B}) of the top quark decaying to a Higgs boson and an up (u) or charm (c) quark are derived through a binned fit to the diphoton invariant mass spectrum. The observed (expected) 95% confidence level upper limits are found to be 0.019% (0.031%) for $\mathcal{B}(t \rightarrow Hu)$ and 0.073% (0.051%) for $\mathcal{B}(t \rightarrow Hc)$. These are the strictest upper limits yet determined.

CMS (ATLAS similar)

World Best Bound:

$$t \rightarrow ch < 0.00073$$

Explaining
BAU

1705.05034

Physics Letters B 776 (2018) 402–406

Electroweak baryogenesis driven by extra top Yukawa couplings

Kaori Fuyuto^{a,*}, Wei-Shu Hou^b, Eibun Senaha^c

EWBG Driven by $\lambda_t \text{Im} \rho_{tt}$

Grand Motivation!

^a Amherst Center for Fundamental Interactions, Department of Physics, University of Massachusetts Amherst, MA 01003, USA

^b Department of Physics, National Taiwan University, Taipei 10617, Taiwan

^c Center for Theoretical Physics of the Universe, Institute for Basic Science (IBS), Daejeon 34051, Republic of Korea

ARTICLE INFO

Article history:

Received 19 August 2017

Received in revised form 17 November 2017


Accepted 26 November 2017

Available online 2 December 2017

Editor: M. Trodden

ABSTRACT

We study electroweak baryogenesis driven by the top quark in a general two Higgs doublet model flavor-changing Yukawa couplings, keeping the Higgs potential CP invariant. With Higgs sector couplings and the additional top Yukawa coupling ρ_{tt} all of $\mathcal{O}(1)$, one naturally has sizable CP violation that the cosmic baryon asymmetry. Even if ρ_{tt} vanishes, the favor-changing coupling ρ_{tc} can still lead successful baryogenesis. Phenomenological consequences such as $t \rightarrow ch$, $\tau \rightarrow \mu\gamma$ electron electric dipole moment, $h \rightarrow \gamma\gamma$, and hhh coupling are discussed.



$$\begin{pmatrix} H^+ \\ H + iA \end{pmatrix}$$

Fit for LHC


1706.07694

Sub-TeV H, A, H⁺ @ LHC; G2HDM well-hidden so far.


EPL 123 (2018) 11001

Production
Processes

1710.07260



PLB 776 (2018) 379-384



PRL 125 (2020) 221801

unsuppressed
by alignment

ATLAS-CONF-2022-039 (ICHEP)


Search Started 2/2020.

Fruition 2023!



Finite Chance for Discovery!

Extra Higgs Doublet w/
Extra Yukawa Couplings
& Extra Quartic Couplings



Decadal Mission of New Higgs/Flavor Era

1. CMS: **H, A, H[±]** search @ LHC
2. Belle II: $\tau \rightarrow \mu\gamma$; $B \rightarrow \mu\nu, \tau\nu$; $\tau\tau, \tau\mu$
CMS: $B_{s,d} \rightarrow \mu\mu$
3. Lattice: Higgs Potential { 1stEWPT
Landau Pole
4. Steering: Pheno 粒子現象學

54 extra flavor param.
& 7 add'l Higgs param.

Kai-Feng (Jack) Chen

Paoti Chang

David C.J. Lin
(NYCU)

Wish us Luck!