


QED corrections to the Neutral Current Drell-Yan process using jetiness subtraction method

DPF – PHENO 2024, University of Pittsburg /
Carnegie Mellon University

Jiayang Xiao (Co-Author: Dr. Doreen Wackerath)
May 15th, 2024



Drell-Yan process is one of the best theoretically understood, and most precise experimentally studied process




- N3LO QCD corrections
- NNLO EW×QCD corrections

X. Chen, et al. arxiv:2203.01565
 T. Neumann, et al. arxiv:2207.07056
 X. Chen, et al. arxiv:2107.09085

Charged Current: T. Armadillo, et al. arxiv:2405.00612
 Neutral Current: S. Dittmaier, et al. arxiv:2401.15682
 (and references therein)

- Goal: NNLO mixed QED×QCD corrections with N-jettiness subtraction method
- First step: NLO QED corrections

- Subtraction method for NNLO QCD, N-jettiness subtraction using n-jet resolution variable to handle IR divergence
- N-jettiness: A global inclusive event shape to distinguish between N-jet events and M-jet events ($M > N$).



$$\mathcal{T}_N = \sum_{m=1}^M \min_i \left\{ \frac{2p_i \cdot q_m}{P_i} \right\}$$

Goal: apply this method to NNLO mixed QED \times QCD corrections

- IR safe observable $\mathcal{T}_N(\Phi_{\geq N+1} \rightarrow \Phi_N) \rightarrow 0$


- Recycle lower order calculation

$$\begin{aligned}
 \sigma_{NNLO} &= \int d\Phi_N |\mathcal{M}_N|^2 \\
 &+ \int_{\mathcal{T}_N^{cut}} d\Phi_{N+1} |\mathcal{M}_{N+1}|^2 + \int^{\mathcal{T}_N^{cut}} d\Phi_{N+1} |\mathcal{M}_{N+1}|^2 \\
 &+ \int_{\mathcal{T}_N^{cut}} d\Phi_{N+2} |\mathcal{M}_{N+2}|^2 + \int^{\mathcal{T}_N^{cut}} d\Phi_{N+2} |\mathcal{M}_{N+2}|^2 \\
 &= \sigma_{NNLO}(\mathcal{T}_N > \mathcal{T}_N^{cut}) + \sigma_{NNLO}(\mathcal{T}_N < \mathcal{T}_N^{cut})
 \end{aligned}$$


Factorization formula from Soft and Collinear effective field theory (SCET)

$$\sigma(\mathcal{T}_N < \mathcal{T}_N^{cut}) = \int B_a \otimes B_b \otimes S \otimes H \otimes \prod_{i=1}^N J_i + \mathcal{O}(\mathcal{T}_N^{cut})$$

- $B_{a,b}$: Beam function (contains initial state collinear emissions)




- J_i : Jet function (contains final state collinear emissions)



- H : Hard function (contains process specific hard interaction)

- S : Soft function (contains all soft emissions)





- From QCD to QED


Factorization formula from SCET

$$\sigma(\mathcal{T}_N < \mathcal{T}_N^{cut}) = \int B_a \otimes B_b \otimes S \otimes H \otimes \prod_{i=1}^N J_i + \mathcal{O}(\mathcal{T}_N^{cut})$$

- $B_{a,b}$: Beam function (adjust from QCD corrections)
- J_i : Jet function (adjust from QCD corrections)
- H : Hard function (adjust from QCD corrections and new contribution such as box diagram at NLO QED)
- S : Soft function (new calculation)




- QCD case: 0-jettiness all coefficients are constant
- QED case: 2-jettiness all coefficients must be calculated for each phase space point



$$\begin{aligned}
 \frac{\alpha}{2\pi} S^{(1)} &= |M_{ij}^{(1)}|^2 \{ PS^{(1)}(i, j) [F_i^{i,j} + F_j^{i,j}] + PS^{(1)}(k, i) [F_k^{i,j} + F_l^{i,j}] \} \\
 &= \frac{\alpha}{2\pi} [\mathcal{C}_{-1} \delta(\mathcal{T}_N) + \mathcal{C}_0 \mathcal{L}_0(\mathcal{T}_N) + \mathcal{C}_1 \mathcal{L}_1(\mathcal{T}_N)]
 \end{aligned}$$

$$\int_0^1 dx \mathcal{L}_n(x) f(x) = \int_0^1 dx \frac{\ln^n(x)}{x} [f(x) - f(0)]$$

Challenge: Calculation becomes more complex when final state leptons must be taken account!



Measurement Function

$$F = F_i + F_j + F_k + F_l$$

$$F_i = \delta(\mathcal{T}_N - q^i) \theta(q^j - q^i) \theta(q^k - q^i) \theta(q^l - q^i)$$

$$\begin{aligned}
 \mathcal{T}_2 &= \min_i \left\{ \frac{2p_i \cdot q}{P_i} \right\} \\
 &= \min_i \{q^i\}
 \end{aligned}$$

F_i & F_j direction

$$|M_{i,j}^{(1)}|^2 PS^{(1)}(i, j) F_i^{i,j} =$$

$$-\frac{\alpha}{2\pi} Q_i Q_j \mathcal{T}_N^{-1-2\epsilon} e^{\gamma_E \epsilon} \mu^\epsilon \left(\frac{n_{ij}}{2}\right)^\epsilon \frac{2^{-2\epsilon}(-\epsilon)}{\Gamma[1-\epsilon]} \int_0^1 dx_1 dx_2 dx_3 (1-x_3)^{-\epsilon} \sin(\pi x_2)^{-2\epsilon} \left(\frac{\delta(x_1)}{\epsilon} + \mathcal{L}_0(x_1) + \epsilon \mathcal{L}_1(x_1)\right)$$

$$\times \left(-\frac{\delta(x_3)}{\epsilon} + \mathcal{L}_0(x_3) - \epsilon \mathcal{L}_1(x_3)\right)$$

F_k & F_l direction

$$|M_{i,j}^{(1)}|^2 PS^{(1)}(k, i) F_k^{i,j} =$$

$$-\frac{\alpha}{2\pi} Q_i Q_j \mathcal{T}_N^{-1-2\epsilon} e^{\gamma_E \epsilon} \mu^\epsilon n_{ij} n_{ik}^{-1+\epsilon} \frac{2^{-3\epsilon}(-\epsilon)}{\Gamma[1-\epsilon]} \int_0^1 dx_1 dx_2 dx_3 x_1^\epsilon (1-x_3)^{-\epsilon} \sin(\pi x_2)^{-2\epsilon}$$

$$\times \left(-\frac{\delta(x_3)}{\epsilon} + \mathcal{L}(x_3) - \epsilon \mathcal{L}(x_3)\right)$$

$$\int_0^1 dx \mathcal{L}_n(x) f(x) = \int_0^1 dx \frac{\ln^n(x)}{x} [f(x) - f(0)]$$

$$\begin{aligned}
 & -\frac{\alpha}{2\pi} Q_i Q_j \mathcal{T}_N^{-1-2\epsilon} e^{\gamma_E \epsilon} \mu^\epsilon \left(\frac{n_{ij}}{2}\right)^\epsilon \frac{2^{-2\epsilon} (-\epsilon)}{\Gamma[1-\epsilon]} \int_0^1 dx_1 dx_2 dx_3 (1-x_3)^{-\epsilon} \sin(\pi x_2)^{-2\epsilon} \left(\frac{\delta(x_1)}{\epsilon} + \mathcal{L}_0(x_1) + \epsilon \mathcal{L}_1(x_1) \right) \\
 & \times \left(-\frac{\delta(x_3)}{\epsilon} + \mathcal{L}_0(x_3) - \epsilon \mathcal{L}_1(x_3) \right)
 \end{aligned}$$

- Already a 3-dimensional integration for each phase space point.
- Monte Carlo integration is too time consuming!

Time taken for soft function calculation: 20.07s

$$\begin{aligned}
 &-\frac{\alpha}{2\pi} Q_i Q_j \mathcal{T}_N^{-1-2\epsilon} e^{\gamma_E \epsilon} \mu^\epsilon \left(\frac{n_{ij}}{2}\right)^\epsilon \frac{2^{-2\epsilon} (-\epsilon)}{\Gamma[1-\epsilon]} \int_0^1 dx_1 dx_2 dx_3 (1-x_3)^{-\epsilon} \sin(\pi x_2)^{-2\epsilon} \left(\frac{\delta(x_1)}{\epsilon} + \mathcal{L}_0(x_1) + \epsilon \mathcal{L}_1(x_1) \right) \\
 &\quad \times \left(-\frac{\delta(x_3)}{\epsilon} + \mathcal{L}_0(x_3) - \epsilon \mathcal{L}_1(x_3) \right)
 \end{aligned}$$





- Already a 3-dimensional integration for each phase space point.
- Monte Carlo integration is too time consuming!

Time taken for soft function calculation: 20.07s





- Analytic simplification is applied to reduce to one dimensional integration, and Gauss integration is enough to achieve desired precision.

Time taken for soft function calculation: 1.63495541E-03 s

$$S^{(1)} = C_{-1}\delta(\mathcal{T}_2) + C_0\mathcal{L}_0(\mathcal{T}_2) + C_1\mathcal{L}_1(\mathcal{T}_2)$$




- Final step, combining with B , J and H for implementation using ZGRAD Monte Carlo program.
- The N-jettiness subtraction method results are compared with Phase-space-slicing method implemented in ZGRAD as cross check.



Summary

- N-jettiness subtraction method for NLO QED calculation
- Universal Soft function can be applied to other similar processes
- Next step: NNLO QED \times QCD correction to Drell-Yan process



**Thank You for
Your Time and
Attention!**

