

Top-philic ALPs in the elusive mass window

Ken Mimasu

University of Southampton

[S. Blasi, F. Maltoni, A. Mariotti, KM, D. Pagani, S. Tentori; JHEP 06 (2024) 077]

TOP2024, St. Malo

24th September 2024



Science and
Technology
Facilities Council



University of
Southampton

Axion-like particles

Axions: originally motivated by strong CP problem

$$\mathcal{L} \supset \frac{a}{f_a} G_A^{\mu\nu} \tilde{G}_{\mu\nu}^A \Rightarrow m_a f_a = \text{constant}$$

Axion-like particles

Axions: originally motivated by strong CP problem

$$\mathcal{L} \supset \frac{a}{f_a} G_A^{\mu\nu} \tilde{G}_{\mu\nu}^A \Rightarrow m_a f_a = \text{constant}$$

ALPs: model of light, singlet pseudoscalar, a

- Generic, independent interactions

$$\{m_a, f_a\} \text{ independent}$$

Axion-like particles

Axions: originally motivated by strong CP problem

$$\mathcal{L} \supset \frac{a}{f_a} G_A^{\mu\nu} \tilde{G}_{\mu\nu}^A \Rightarrow m_a f_a = \text{constant}$$

ALPs: model of light, singlet pseudoscalar, a

- Generic, independent interactions

$$\{m_a, f_a\} \text{ independent}$$

- SM singlet

\Rightarrow Interactions described by EFT starting at dimension-5, $\mathcal{O}(1/f_a)$

Axion-like particles

Axions: originally motivated by strong CP problem

$$\mathcal{L} \supset \frac{a}{f_a} G_A^{\mu\nu} \tilde{G}_{\mu\nu}^A \Rightarrow m_a f_a = \text{constant}$$

ALPs: model of light, singlet pseudoscalar, a

- Generic, independent interactions

$$\{m_a, f_a\} \text{ independent}$$

- SM singlet

\Rightarrow Interactions described by EFT starting at dimension-5, $\mathcal{O}(1/f_a)$

- Pseudo Nambu-Goldstone boson

\Rightarrow light particle with shift-symmetric interactions

$$a(x) \rightarrow a(x) + c \Rightarrow \mathcal{L} = \mathcal{L}[\partial^\mu a(x)]$$

- Appear in many well-motivated BSM scenarios

ALP interactions

$$\mathcal{L}_{ALP}^{(5)} = \frac{1}{2} \partial^\mu a \partial_\mu a - \frac{1}{2} m_a^2 a^2 + \frac{\partial^\mu a}{f_a} \sum_f \bar{\psi}_f \mathbf{c}_f \gamma_\mu \psi_f + c_H \frac{\partial^\mu a}{f_a} H^\dagger i \overleftrightarrow{D}_\mu H$$

$$+ c_{GG} \frac{\alpha_S}{4\pi} \frac{a}{f_a} G_A^{\mu\nu} \tilde{G}_{\mu\nu}^A + c_{WW} \frac{\alpha_2}{4\pi} \frac{a}{f_a} W_I^{\mu\nu} \tilde{W}_{\mu\nu}^I + c_{BB} \frac{\alpha_Y}{4\pi} \frac{a}{f_a} B_I^{\mu\nu} \tilde{B}_{\mu\nu}$$

f : 5 fermionic SM representations $\{Q, L, u, d, e\}$

- \mathbf{c}_f are matrices in flavor space

ALP interactions

$$\begin{aligned}
 \mathcal{L}_{ALP}^{(5)} = & \frac{1}{2} \partial^\mu a \partial_\mu a - \frac{1}{2} m_a^2 a^2 + \frac{\partial^\mu a}{f_a} \sum_f \bar{\psi}_f \mathbf{c}_f \gamma_\mu \psi_f + c_H \frac{\partial^\mu a}{f_a} H^\dagger i \overleftrightarrow{D}_\mu H \\
 & + c_{GG} \frac{\alpha_S}{4\pi} \frac{a}{f_a} G_A^{\mu\nu} \tilde{G}_{\mu\nu}^A + c_{WW} \frac{\alpha_2}{4\pi} \frac{a}{f_a} W_I^{\mu\nu} \tilde{W}_{\mu\nu}^I + c_{BB} \frac{\alpha_Y}{4\pi} \frac{a}{f_a} B_I^{\mu\nu} \tilde{B}_{\mu\nu}
 \end{aligned}$$

explicit shift symmetry

f : 5 fermionic SM representations $\{Q, L, u, d, e\}$

- \mathbf{c}_f are matrices in flavor space

ALP interactions

$$\begin{aligned}
 \mathcal{L}_{ALP}^{(5)} = & \frac{1}{2} \partial^\mu a \partial_\mu a - \frac{1}{2} m_a^2 a^2 + \frac{\partial^\mu a}{f_a} \sum_f \bar{\psi}_f \mathbf{c}_f \gamma_\mu \psi_f + c_H \frac{\partial^\mu a}{f_a} H^\dagger i \overleftrightarrow{D}_\mu H \\
 & + c_{GG} \frac{\alpha_S}{4\pi} \frac{a}{f_a} G_A^{\mu\nu} \tilde{G}_{\mu\nu}^A + c_{WW} \frac{\alpha_2}{4\pi} \frac{a}{f_a} W_I^{\mu\nu} \tilde{W}_{\mu\nu}^I + c_{BB} \frac{\alpha_Y}{4\pi} \frac{a}{f_a} B_I^{\mu\nu} \tilde{B}_{\mu\nu}
 \end{aligned}$$

explicit shift symmetry

Anomaly induced, 'hidden' shift symmetry

f : 5 fermionic SM representations $\{Q, L, u, d, e\}$

- \mathbf{c}_f are matrices in flavor space

ALP interactions

$$\mathcal{L}_{ALP}^{(5)} = \frac{1}{2} \partial^\mu a \partial_\mu a - \frac{1}{2} m_a^2 a^2 + \frac{\partial^\mu a}{f_a} \sum_f \bar{\psi}_f \mathbf{c}_f \gamma_\mu \psi_f + c_H \frac{\partial^\mu a}{f_a} H^\dagger i \overleftrightarrow{D}_\mu H$$

$$+ c_{GG} \frac{\alpha_S}{4\pi} \frac{a}{f_a} G_A^{\mu\nu} \tilde{G}_{\mu\nu}^A + c_{WW} \frac{\alpha_2}{4\pi} \frac{a}{f_a} W_I^{\mu\nu} \tilde{W}_{\mu\nu}^I + c_{BB} \frac{\alpha_Y}{4\pi} \frac{a}{f_a} B_I^{\mu\nu} \tilde{B}_{\mu\nu}$$

explicit shift symmetry

shift symmetry breaking

Anomaly induced, 'hidden' shift symmetry

f : 5 fermionic SM representations $\{Q, L, u, d, e\}$

- \mathbf{c}_f are matrices in flavor space

ALP interactions

$$\mathcal{L}_{ALP}^{(5)} = \frac{1}{2} \partial^\mu a \partial_\mu a - \frac{1}{2} m_a^2 a^2 + \frac{\partial^\mu a}{f_a} \sum_f \bar{\psi}_f \mathbf{c}_f \gamma_\mu \psi_f + c_H \frac{\partial^\mu a}{f_a} H^\dagger i \overleftrightarrow{D}_\mu H$$

$$+ c_{GG} \frac{\alpha_S}{4\pi} \frac{a}{f_a} G_A^{\mu\nu} \tilde{G}_{\mu\nu}^A + c_{WW} \frac{\alpha_2}{4\pi} \frac{a}{f_a} W_I^{\mu\nu} \tilde{W}_{\mu\nu}^I + c_{BB} \frac{\alpha_Y}{4\pi} \frac{a}{f_a} B_I^{\mu\nu} \tilde{B}_{\mu\nu}$$

explicit shift symmetry

shift symmetry breaking

Anomaly induced, 'hidden' shift symmetry

f : 5 fermionic SM representations $\{Q, L, u, d, e\}$

- \mathbf{c}_f are matrices in flavor space

Not all operators independent: $\{c_H, [c_f]_{ij}\}$

[Bauer, Neubert & Thamm; JHEP 12 (2017) 044]

ALP interactions

$$\mathcal{L}_{ALP}^{(5)} = \frac{1}{2} \partial^\mu a \partial_\mu a - \frac{1}{2} m_a^2 a^2 + \frac{\partial^\mu a}{f_a} \sum_f \bar{\psi}_f \mathbf{c}_f \gamma_\mu \psi_f + c_H \frac{\partial^\mu a}{f_a} H^\dagger i \overleftrightarrow{D}_\mu H$$

$$+ c_{GG} \frac{\alpha_S}{4\pi} \frac{a}{f_a} G_A^{\mu\nu} \tilde{G}_{\mu\nu}^A + c_{WW} \frac{\alpha_2}{4\pi} \frac{a}{f_a} W_I^{\mu\nu} \tilde{W}_{\mu\nu}^I + c_{BB} \frac{\alpha_Y}{4\pi} \frac{a}{f_a} B_I^{\mu\nu} \tilde{B}_{\mu\nu}$$

explicit shift symmetry

shift symmetry breaking

Anomaly induced, 'hidden' shift symmetry

f : 5 fermionic SM representations $\{Q, L, u, d, e\}$

- \mathbf{c}_f are matrices in flavor space


Not all operators independent: $\{c_H, [c_f]_{ij}\}$


[Bauer, Neubert & Thamm; JHEP 12 (2017) 044]

Only one operator at dimension-6:

[Grojean, Kley & Yao; JHEP 11 (2023) 196]

$$\mathcal{L}_{ALP}^{(6)} = \frac{c_{aH}^{(6)}}{f_a^2} (H^\dagger H) \partial^\mu a \partial_\mu a$$





Top philic ALP

Consider an ALP that preferentially couples to the top quark

- e.g. t_R mixing with new sector

$$\mathcal{L}_{top}^{(5)} = c_t \frac{\partial^\mu a}{f_a} \bar{t}_R \gamma_\mu t_R, \quad c_t = [\mathbf{c}_u]_{33}$$

- Consistent with Minimal Flavor Violation

Recent works:

[\[Esser et al.; JHEP 09 \(2023\) 063\]](#)

[\[Biekötter et al.; JHEP 09 \(2023\) 120\]](#)

[\[Rygaard et al.; JHEP 10 \(2023\) 138\]](#)

[\[Bruggisser et al.; JHEP 01 \(2024\) 092\]](#)

[\[Hosseini et al; PRD 110 \(2024\) 5, 055026\]](#)

[\[Phan & Westhoff; JHEP 05 \(2024\) 075\]](#)

[\[Anuar et al.; 2404.19014\]](#)

Top philic ALP

Consider an ALP that preferentially couples to the top quark

- e.g. t_R mixing with new sector

Recent works:

[\[Esser et al.; JHEP 09 \(2023\) 063\]](#)

[\[Biekötter et al.; JHEP 09 \(2023\) 120\]](#)

[\[Rygaard et al.; JHEP 10 \(2023\) 138\]](#)

[\[Bruggisser et al.; JHEP 01 \(2024\) 092\]](#)

[\[Hosseini et al; PRD 110 \(2024\) 5, 055026\]](#)

[\[Phan & Westhoff; JHEP 05 \(2024\) 075\]](#)

[\[Anuar et al.; 2404.19014\]](#)

$$\mathcal{L}_{top}^{(5)} = c_t \frac{\partial^\mu a}{f_a} \bar{t}_R \gamma_\mu t_R, \quad c_t = [\mathbf{c}_u]_{33}$$

- Consistent with Minimal Flavor Violation

Equivalent basis: $t_R \rightarrow t_R e^{-ic_t \frac{a}{f_a}}$

Derivative

Non-derivative (Yukawa-like)

$$c_t \frac{\partial^\mu a}{f_a} \bar{t}_R \gamma_\mu t_R \quad \Leftrightarrow \quad c_t \frac{a}{f_a} \left(-im_t \bar{t} \gamma_5 t + \frac{\alpha_S}{8\pi} G\tilde{G} + \frac{\alpha_Y}{3\pi} B\tilde{B} \right)$$

Top philic ALP

Consider an ALP that preferentially couples to the top quark

- e.g. t_R mixing with new sector

Recent works:

[\[Esser et al.; JHEP 09 \(2023\) 063\]](#)

[\[Biekötter et al.; JHEP 09 \(2023\) 120\]](#)

[\[Rygaard et al.; JHEP 10 \(2023\) 138\]](#)

[\[Bruggisser et al.; JHEP 01 \(2024\) 092\]](#)

[\[Hosseini et al.; PRD 110 \(2024\) 5, 055026\]](#)

[\[Phan & Westhoff; JHEP 05 \(2024\) 075\]](#)

[\[Anuar et al.; 2404.19014\]](#)

$$\mathcal{L}_{top}^{(5)} = c_t \frac{\partial^\mu a}{f_a} \bar{t}_R \gamma_\mu t_R, \quad c_t = [\mathbf{c}_u]_{33}$$

- Consistent with Minimal Flavor Violation

Equivalent basis: $t_R \rightarrow t_R e^{-ic_t \frac{a}{f_a}}$

Derivative

Non-derivative (Yukawa-like)


$$c_t \frac{\partial^\mu a}{f_a} \bar{t}_R \gamma_\mu t_R \quad \Leftrightarrow \quad c_t \frac{a}{f_a} \left(-im_t \bar{t} \gamma_5 t + \frac{\alpha_S}{8\pi} G\tilde{G} + \frac{\alpha_Y}{3\pi} B\tilde{B} \right)$$

Top-philic ALP \neq top-philic pseudo scalar!


Top philic ALP couplings

Top coupling induces light fermion couplings, $c_{f \neq t}$

$$c_b \simeq 5c_t \frac{y_t^2}{16\pi^2} \log \frac{\Lambda}{m_t}$$




$$c_f \simeq -12c_t \frac{y_t^2}{16\pi^2} T_3^f \log \frac{\Lambda}{m_t}$$




Top philic ALP couplings

Top coupling induces light fermion couplings, $c_{f \neq t}$


$$c_b \simeq 5c_t \frac{y_t^2}{16\pi^2} \log \frac{\Lambda}{m_t}$$



$$c_f \simeq -12c_t \frac{y_t^2}{16\pi^2} T_3^f \log \frac{\Lambda}{m_t}$$



Effective vertices between ALP & gauge bosons




$+ \gamma\gamma, Z\gamma, WW, ZZ$

[Neubert et al; JHEP 12 (2017) 044]

[Bonilla et al.; JHEP 11 (2021) 168]

Elusive gauge couplings

Induced $a \rightarrow gg$ amplitude (different from pseudoscalar)




$$\mathcal{A}^{\mu\nu} [a(k) \rightarrow g_a(p) g_b(q)] = i \frac{\alpha_S}{\pi} \frac{c_t}{f_a} \delta_{ab} p_\alpha q_\beta \epsilon^{\mu\nu\alpha\beta} [1 + 2m_t^2 C_0(p, q; m_t^2)]$$

$$k^2 \sim p^2 \sim q^2 \ll m_t^2 \quad \Rightarrow \quad \mathcal{A}^{\mu\nu} \rightarrow 0$$

Elusive gauge couplings

Induced $a \rightarrow gg$ amplitude (different from pseudoscalar)



$$\mathcal{A}^{\mu\nu} [a(k) \rightarrow g_a(p) g_b(q)] = i \frac{\alpha_S}{\pi} \frac{c_t}{f_a} \delta_{ab} p_\alpha q_\beta \epsilon^{\mu\nu\alpha\beta} [1 + 2m_t^2 C_0(p, q; m_t^2)]$$


$$k^2 \sim p^2 \sim q^2 \ll m_t^2 \Rightarrow \mathcal{A}^{\mu\nu} \rightarrow 0$$

Effective $aG\tilde{G}$ contact interaction

$$\text{On-shell } (m_a \ll m_t) \quad c_{GG}^{\text{eff}} \sim \frac{\alpha_S}{\pi} \frac{c_t}{f_a} \frac{m_a^2}{24m_t^2} \quad \text{Off-shell } (k^2 \gg m_t^2) \quad c_{GG}^{\text{eff}} \sim \frac{\alpha_S}{\pi} \frac{c_t}{f_a}$$

Elusive gauge couplings

Induced $a \rightarrow gg$ amplitude (different from pseudoscalar)



$$\mathcal{A}^{\mu\nu} [a(k) \rightarrow g_a(p) g_b(q)] = i \frac{\alpha_S}{\pi} \frac{c_t}{f_a} \delta_{ab} p_\alpha q_\beta \epsilon^{\mu\nu\alpha\beta} [1 + 2m_t^2 C_0(p, q; m_t^2)]$$

$$k^2 \sim p^2 \sim q^2 \ll m_t^2 \Rightarrow \mathcal{A}^{\mu\nu} \rightarrow 0$$

Effective $aG\tilde{G}$ contact interaction


$$\text{On-shell } (m_a \ll m_t) \quad c_{GG}^{\text{eff}} \sim \frac{\alpha_S}{\pi} \frac{c_t}{f_a} \frac{m_a^2}{24m_t^2} \quad \text{Off-shell } (k^2 \gg m_t^2) \quad c_{GG}^{\text{eff}} \sim \frac{\alpha_S}{\pi} \frac{c_t}{f_a}$$

Induced light fermion couplings can be relevant

$$c_{GG}^{\text{eff}} \sim \frac{\alpha_S}{\pi} \frac{c_t}{f_a} \left[\frac{m_a^2}{24m_t^2} + \frac{5}{2} \frac{y_t^2}{16\pi^2} \log \frac{\Lambda}{m_t} \right]$$


*e.g. for ALP production & decay
(same for $a \rightarrow \gamma\gamma$)*

Flavour probes



Flavour probes


[Ebadi et al; PRD 100 (2019) 015016]
 Top chromo-magnetic dipole bound



Stellar/DM bounds

Strong limits from RG-induced ALP-electron coupling

95%CL excluded region from constraints on ALP-electron coupling



Elusive mass window

$$10 \lesssim m_a \lesssim 200 \text{ GeV}$$

Strong bounds from astro/flavour below $m_a \sim \text{few GeV}$

- ALP possibly long-lived

[Esser et al.; JHEP 09 (2023) 063]

[Bruggisser et al.; JHEP 01 (2024) 092]

Elusive mass window

$$10 \lesssim m_a \lesssim 200 \text{ GeV}$$

Strong bounds from astro/flavour below $m_a \sim \text{few GeV}$

- ALP possibly long-lived

[Esser et al.; JHEP 09 (2023) 063]

[Bruggisser et al.; JHEP 01 (2024) 092]

Larger m_a means shift symmetry breaking effects $\propto m_a^2$

- Decays to gauge bosons unsuppressed

Elusive mass window

$$10 \lesssim m_a \lesssim 200 \text{ GeV}$$

Strong bounds from astro/flavour below $m_a \sim \text{few GeV}$

- ALP possibly long-lived


[Esser et al.; JHEP 09 (2023) 063]

[Bruggisser et al.; JHEP 01 (2024) 092]

Larger m_a means shift symmetry breaking effects $\propto m_a^2$

- Decays to gauge bosons unsuppressed

$m_a > 2m_t$, on-shell top decays $\Rightarrow t\bar{t}$ resonance searches




[Anuar et al.; 2404.19014]

See also Laurids Jeppe's poster

- Peak-dip structure from interference in $gg \rightarrow t\bar{t}$
- ALP differs from top-philic pseudo scalar

Branching ratios

dominant $b\bar{b}$ decays



*strong flavour/
astro bounds*




*shift-symmetry
breaking:
 $a \rightarrow gg, \gamma\gamma, t\bar{t}$*



Bounds in elusive window

Recast light scalar resonance searches

- $t\bar{t}a$ production, boosted dijet & $h \rightarrow aa^*$



Exp. searches


- $t\bar{t}a(a \rightarrow \mu^+ \mu^-)$
[\[CMS; PRD 110 \(2024\) 012013\]](#)
- $t\bar{t}a(a \rightarrow \tau^+ \tau^-)$
- Boosted dijet
[\[ATLAS; PLB 788 \(2019\) 316-335\]](#)
- - - $H \rightarrow \text{BSM}$
[\[ATLAS & CMS; Nature \(2022\)\]](#)
- ⋯ $H \rightarrow b\bar{b}\mu^+\mu^-$
[\[ATLAS; PRD 105 \(2022\) 012006\]](#)

We explore top-rich final states

Resonant ALPs in $t\bar{t}b\bar{b}$

Top pair associated production into primary $b\bar{b}$ decay mode

- Constrain contribution to $t\bar{t}b\bar{b}$ signal strength measurements



| Exp. | Channel | $\mu_{t\bar{t}b\bar{b}} \pm \text{stat.} \pm \text{syst.}$ |
|-------|-------------------------|------------------------------------------------------------|
| CMS | dilepton | $1.36 \pm 0.10 \pm 0.34$ |
| CMS | lepton+jets | $1.26 \pm 0.04 \pm 0.31$ |
| ATLAS | dilepton ($e\mu$, 4b) | $1.75 \pm 0.05 \pm 0.56$ |
| ATLAS | lepton+jets (4b) | $1.57 \pm 0.09 \pm 0.49$ |


[\[ATLAS; JHEP 04 \(2019\) 046\]](#)

[\[CMS; JHEP 07 \(2020\) 125\]](#)

Resonant ALPs in $t\bar{t}b\bar{b}$

Top pair associated production into primary $b\bar{b}$ decay mode


- Constrain contribution to $t\bar{t}b\bar{b}$ signal strength measurements



| Exp. | Channel | $\mu_{t\bar{t}b\bar{b}} \pm \text{stat.} \pm \text{syst.}$ |
|-------|-------------------------|------------------------------------------------------------|
| CMS | dilepton | $1.36 \pm 0.10 \pm 0.34$ |
| CMS | lepton+jets | $1.26 \pm 0.04 \pm 0.31$ |
| ATLAS | dilepton ($e\mu$, 4b) | $1.75 \pm 0.05 \pm 0.56$ |
| ATLAS | lepton+jets (4b) | $1.57 \pm 0.09 \pm 0.49$ |

[\[ATLAS; JHEP 04 \(2019\) 046\]](#)

[\[CMS; JHEP 07 \(2020\) 125\]](#)



- Significant b -tagging and modelling systematics
- Resonant channels dominate
- Dedicated resonance search should be beneficial

Non-resonant ALPs in $t\bar{t}t\bar{t}$

Purely non-resonant ($m_a < 2m_t$)


- Corrections to total four top cross section

[ATLAS; EPJC 83 (2023) 496]

[CMS; PLB 847 (2023) 138290]

[CMS; PLB 844 (2023) 138076]

[ATLAS; JHEP 11 (2021) 118]



| Exp. | Channel | $\mu_{t\bar{t}t\bar{t}} \pm \text{stat.} \pm \text{syst.}$ |
|-------|---------|------------------------------------------------------------|
| ATLAS | SSDL+ML | $1.70 \pm 0.40^{+0.7}_{-0.4}$ |
| ATLAS | OSDL+1L | $2.00 \pm 0.70^{+1.5}_{-1.0}$ |
| CMS | SSDL+ML | $1.32 \pm 0.27^{+0.2}_{-0.23}$ |
| CMS | OSDL+1L | $2.20 \pm 0.50 \pm 0.50$ |

Non-resonant ALPs in $t\bar{t}t\bar{t}$

Purely non-resonant ($m_a < 2m_t$)


- Corrections to total four top cross section

[ATLAS; EPJC 83 (2023) 496]


[CMS; PLB 847 (2023) 138290]

[CMS; PLB 844 (2023) 138076]

[ATLAS; JHEP 11 (2021) 118]



| Exp. | Channel | $\mu_{t\bar{t}t\bar{t}} \pm \text{stat.} \pm \text{syst.}$ |
|-------|---------|------------------------------------------------------------|
| ATLAS | SSDL+ML | $1.70 \pm 0.40^{+0.7}_{-0.4}$ |
| ATLAS | OSDL+1L | $2.00 \pm 0.70^{+1.5}_{-1.0}$ |
| CMS | SSDL+ML | $1.32 \pm 0.27^{+0.2}_{-0.23}$ |
| CMS | OSDL+1L | $2.20 \pm 0.50 \pm 0.50$ |



- Included mixed QCD/QED contributions

- Combined 4 recent $t\bar{t}t\bar{t}$ analyses

- Latest SM prediction

[van Beekveld et al.; PRL 131 (2023) 21, 211901]





- Comparable contributions from:

$$\text{interference w/ SM} \propto c_t^2$$

$$\text{ALP diagram squared} \propto c_t^4$$

Indirect ALPs in $t\bar{t}$



Loop corrections to $t\bar{t}$ production: $\sigma = \sigma_{SM} + \sigma_{virt.} + \sigma_{real}$



(Diagrams in non-derivative basis)

Indirect ALPs in $t\bar{t}$

Loop corrections to $t\bar{t}$ production: $\sigma = \sigma_{SM} + \sigma_{virt.} + \sigma_{real}$



- Real corrections negligible
- $\sigma_{virt.} \sim c_t^2$ is interference w/ SM
- Large deviation near threshold

(Diagrams in non-derivative basis)

Indirect ALPs in $t\bar{t}$


Combined several $t\bar{t}$ measurements from LHC experiments

- From `fitmaker` database [\[Ellis, Madigan, KM, Sanz, You; JHEP 04 \(2021\) 279\]](#)
- Non-overlapping combination of $m_{t\bar{t}}$ & p_T distributions (see backup)
- Including correlations where available

Indirect ALPs in $t\bar{t}$

Combined several $t\bar{t}$ measurements from LHC experiments

- From `fitmaker` database [\[Ellis, Madigan, KM, Sanz, You; JHEP 04 \(2021\) 279\]](#)
- Non-overlapping combination of $m_{t\bar{t}}$ & p_T distributions (see backup)
- Including correlations where available




Bounds

| m_a [GeV] | 10 | 50 | 100 | 150 | 200 |
|-------------------------|-----|-----|-----|-----|-----|
| $\frac{f_a}{c_t}$ [GeV] | 201 | 206 | 212 | 221 | 234 |

- Mild dependence on m_a
- Outperforms previous searches
- Better sensitivity than $t\bar{t}b\bar{b}$ & $t\bar{t}t\bar{t}$



Summary of top-rich bounds



The elusive ALP

Top-philic ALP with $10 \lesssim m_a \lesssim 100 \text{ GeV}$ remains elusive...



- Suppressed couplings to gauge bosons make for tough direct searches
- Exotic Higgs decay mode competitive but comes with caveat (see backup)
- Top-rich bounds constrain $f_a/c_t \sim 200 \text{ GeV}$ *EFT valid for strongly coupled scenario*



What next?

Dedicated resonance searches

- $m_a > 90 \text{ GeV}$, $a \rightarrow Z\gamma$
- $m_a > 160 \text{ GeV}$, $a \rightarrow W^+W^-$
- $t\bar{t}a$ production mode dominates at these masses
- Interesting top-rich signatures with small SM backgrounds
- Dedicated studies in progress... stay tuned!



Conclusions

$$10 \lesssim m_a \lesssim 200 \text{ GeV}$$

Top philic ALP is an interesting yet elusive collider target

- Direct $gg \rightarrow a$ production mode suppressed & predominant $b\bar{b}$ decay
- Precision measurements of top-rich final states can be used as searches
- Best bounds come from $t\bar{t}$ differential distributions & $h \rightarrow \mathbf{BSM}$

Conclusions

$$10 \lesssim m_a \lesssim 200 \text{ GeV}$$

Top philic ALP is an interesting yet elusive collider target

- Direct $gg \rightarrow a$ production mode suppressed & predominant $b\bar{b}$ decay
- Precision measurements of top-rich final states can be used as searches
- Best bounds come from $t\bar{t}$ differential distributions & $h \rightarrow \text{BSM}$

Developed a dedicated UFO model for MC simulations

- Currently not public

[*\[Maltoni et al; JHEP 09 \(2024\) 098\]*](#)

Conclusions

$$10 \lesssim m_a \lesssim 200 \text{ GeV}$$

Top philic ALP is an interesting yet elusive collider target

- Direct $gg \rightarrow a$ production mode suppressed & predominant $b\bar{b}$ decay
- Precision measurements of top-rich final states can be used as searches
- Best bounds come from $t\bar{t}$ differential distributions & $h \rightarrow \text{BSM}$

Developed a dedicated UFO model for MC simulations

- Currently not public

[\[Maltoni et al; JHEP 09 \(2024\) 098\]](#)

Dedicated searches would be welcome

- Non-resonant, kinematic distributions in e.g. $t\bar{t}$ & $t\bar{t}t\bar{t}$
- Resonant: $t\bar{t}a$ production with decays to $b\bar{b}/Z\gamma/WW$

Conclusions

$$10 \lesssim m_a \lesssim 200 \text{ GeV}$$

Top philic ALP is an interesting yet elusive collider target

- Direct $gg \rightarrow a$ production mode suppressed & predominant $b\bar{b}$ decay
- Precision measurements of top-rich final states can be used as searches
- Best bounds come from $t\bar{t}$ differential distributions & $h \rightarrow \text{BSM}$

Developed a dedicated UFO model for MC simulations

- Currently not public

[\[Maltoni et al; JHEP 09 \(2024\) 098\]](#)

Dedicated searches would be welcome

- Non-resonant, kinematic distributions in e.g. $t\bar{t}$ & $t\bar{t}t\bar{t}$
- Resonant: $t\bar{t}a$ production with decays to $b\bar{b}/Z\gamma/WW$

Not discussed today: top-philic ALP as DM mediator [\[backup\]](#)

Backup



Science and
Technology
Facilities Council



University of
Southampton

Global approach



Generic ALP should be studied from global perspective

- More than one coupling at a time \Rightarrow correlations [\[Bruggisser et al; JHEP 01 \(2024\) 092\]](#)
- Enable constraints on generic UV completions

ALP-SMEFT mixing:

[\[Biekötter et al; JHEP 09 \(2023\) 120\]](#)

$$\frac{d}{d \log \mu} C_i^{\text{SMEFT}} - \gamma_{ji}^{\text{SMEFT}} C_j^{\text{SMEFT}} = \frac{S_i}{(4\pi f)^2}$$



$t\bar{t}$ data

$m_{t\bar{t}}$

| \sqrt{s} | Collab. | Channel | bins | Ref. |
|------------|---------|--------------|------|---------|
| 8 TeV | ATLAS | Dilepton | 6 | [82] |
| 8 TeV | ATLAS | ℓ +jets | 7 | [83] |
| 8 TeV | CMS | Dilepton | 6 | [84](a) |
| 8 TeV | CMS | ℓ +jets | 7 | [84](b) |
| 13 TeV | ATLAS | ℓ +jets | 9 | [85] |
| 13 TeV | CMS | Dilepton | 7 | [86] |
| 13 TeV | CMS | ℓ +jets | 10 | [87] |
| 13 TeV | CMS | ℓ +jets | 15 | [88] |

p_T

| \sqrt{s} | Collab. | channel | bins | Ref. |
|------------|---------|--------------|------|---------|
| 8 TeV | ATLAS | ℓ +jets | 8 | [83] |
| 8 TeV | CMS | Dilepton | 5 | [84](a) |
| 8 TeV | CMS | ℓ +jets | 8 | [84](b) |
| 13 TeV | ATLAS | ℓ +jets | 8 | [85] |
| 13 TeV | CMS | Dilepton | 6 | [86] |
| 13 TeV | CMS | ℓ +jets | 17 | [88] |

[82] ATLAS collaboration, *Measurement of top quark pair differential cross-sections in the dilepton channel in pp collisions at $\sqrt{s} = 7$ and 8 TeV with ATLAS*, *Phys. Rev. D* **94** (2016) 092003 [Addendum *ibid.* **101** (2020) 119901] [[arXiv:1607.07281](#)] [[INSPIRE](#)].

[83] ATLAS collaboration, *Measurements of top-quark pair differential cross-sections in the lepton+jets channel in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector*, *Eur. Phys. J. C* **76** (2016) 538 [[arXiv:1511.04716](#)] [[INSPIRE](#)].

[84] CMS collaboration, *Measurement of the differential cross section for top quark pair production in pp collisions at $\sqrt{s} = 8$ TeV*, *Eur. Phys. J. C* **75** (2015) 542 [[arXiv:1505.04480](#)] [[INSPIRE](#)].

[85] ATLAS collaboration, *Measurements of top-quark pair differential and double-differential cross-sections in the ℓ +jets channel with pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector*, *Eur. Phys. J. C* **79** (2019) 1028 [Erratum *ibid.* **80** (2020) 1092] [[arXiv:1908.07305](#)] [[INSPIRE](#)].

[86] CMS collaboration, *Measurements of $t\bar{t}$ differential cross sections in proton-proton collisions at $\sqrt{s} = 13$ TeV using events containing two leptons*, *JHEP* **02** (2019) 149 [[arXiv:1811.06625](#)] [[INSPIRE](#)].

[87] CMS collaboration, *Measurement of differential cross sections for the production of top quark pairs and of additional jets in lepton+jets events from pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **97** (2018) 112003 [[arXiv:1803.08856](#)] [[INSPIRE](#)].

[88] CMS collaboration, *Measurement of differential $t\bar{t}$ production cross sections in the full kinematic range using lepton+jets events from proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **104** (2021) 092013 [[arXiv:2108.02803](#)] [[INSPIRE](#)].

Chose independent combination to maximise sensitivity



DM mediator

$$\mathcal{L}_{\text{ALP}} \supset i\bar{\chi}\partial^\mu\gamma_\mu\chi - m_{\text{DM}} - ic_{\text{DM}}\frac{m_{\text{DM}}}{f_a}a\bar{\chi}\gamma^5\chi$$



Indirect searches constrain DM mediator scenario

Dedicated missing E_T searches are more powerful

- $h \rightarrow$ invisible, $t\bar{t}(a \rightarrow \text{MET})$ & monojet





$$pp \rightarrow ajj$$




It is known that this channel probes CP of scalar

- Dijet + MET as a probe for (pseudo)scalar DM mediator





Scalar vs. [\[Haisch & Polesello; JHEP 02 \(2019\)128\]](#)
pseudoscalar



- ALP differs from pseudo scalar
- Very small production cross-section...



Real vs. virt. & s -channel




Assumptions & caveats

Only c_t is generated at matching scale, Λ

- Some couplings generated by RG running assumed to be zero at Λ
- We set $\Lambda=1$ TeV, $c_i \propto \log \frac{\Lambda}{E}$

Indirect bounds arise at order $1/f_a^2 \Rightarrow$ dimension-6

- ALP-EFT & SMEFT operators at dim-6 also assumed to be zero at Λ
- e.g. $h \rightarrow aa$ can also arise at tree level @ dim-6 from $c_{aH}^{(6)}$



Toy model

SM + 2 vector-like fermions: T , Ψ & complex scalar, Φ

- T , Ψ share top quantum numbers (top partners)

$$\mathcal{L}_{\text{UV}} = y\Phi\bar{T}_L T_R + \delta\bar{T}_L t_R + y'\Phi^*\bar{\Psi}_L \Psi_R + \delta'\bar{\Psi}_L t_R + \text{h.c.},$$

- Chirally charged under a global $U(1)$ symmetry, broken by $\langle\Phi\rangle = f_a$

$$Q(\Phi) = 1, \quad Q(T_R) = -1, \quad Q(\Psi_R) = 1, \quad \Phi = \frac{1}{\sqrt{2}}(f_a + \rho)e^{ia/f_a}$$

$$\begin{aligned} T_R &\rightarrow e^{-ia/f_a} T_R \\ \Psi_R &\rightarrow e^{ia/f_a} \Psi_R \end{aligned} \Rightarrow \mathcal{L}_{\text{UV}} = -\frac{1}{f_a}\partial_\mu a \bar{T}_R \gamma^\mu T_R + \frac{1}{f_a}\partial_\mu a \bar{\Psi}_R \gamma^\mu \Psi_R - m_T \bar{T} T - m_\Psi \bar{\Psi} \Psi + (\delta\bar{T}_L t_R + \text{h.c.})$$

- One massless combination before EWSB: $c_\theta t_R + s_\theta T_R$
- Integrating out T_R leads to our ALP-top interaction, $c_t = -\delta^2/m_T^2$

$$\frac{\delta\mathcal{L}_{\text{UV}}}{\delta\bar{T}_L} = \frac{\delta\mathcal{L}_{\text{UV}}}{\delta T_R} = 0 \quad \rightarrow \quad T_R = \frac{\delta}{m_T} t_R, \quad T_L = -\frac{\delta}{m_T^2 f_a} \partial_\mu a \gamma^\mu t_R. \quad \Rightarrow \quad \mathcal{L}_{a,\text{int}} = -\frac{\delta^2}{m_T^2} \frac{1}{f_a} (\partial_\mu a) \bar{t}_R \gamma^\mu t_R$$

Toy model 2

Dimension-6 operator, $\mathcal{O}_{aH}^{(6)}$ depends on Φ potential

$$\mathcal{L}_\Phi = |\partial_\mu \Phi|^2 + \mu^2 |\Phi|^2 - \lambda |\Phi|^4 + \boxed{\kappa |\Phi|^2 \phi^\dagger \phi} \quad \text{Higgs portal}$$

After $U(1)$ symmetry breaking $\Phi = \frac{1}{\sqrt{2}}(f_a + \rho)e^{ia/f_a}$

$$\mathcal{L} \supset \frac{1}{2}(\partial_\mu \rho)^2 + \frac{1}{2}(\partial_\mu a)^2 + \frac{\rho}{f_a}(\partial_\mu a)^2 - \frac{1}{2}m_\rho^2 \rho^2 + \frac{\kappa}{2}(f_a^2 + 2\rho f_a + \rho^2)\phi^\dagger \phi + \mathcal{O}(\rho^3) \quad (m_\rho^2 = \lambda f_a^2)$$

Integrate out ρ at tree level yields dimension-6 operator

$$\frac{\kappa}{\lambda} \frac{1}{f_a^2} (\phi^\dagger \phi) \partial^\mu a \partial_\mu a$$

Depends on $|\Phi|^2 |\phi|^2$ portal relative to $|\Phi|^4$ interaction

UV interpretation

SMEFT-UV connection is model dependent by construction

- Implications on heavy new physics & validity of EFT is ***a posteriori***
- Depends on **sensitivity** & **energy scale** probed by data
- Bottom-up philosophy: new physics scale unknown

In general:

- Larger coupling \Leftrightarrow higher new physics scale
- Better for EFT validity
- Some coefficients typically come with loop factors
- e.g. c_{VV} gauge interactions

UV interpretation

SMEFT-UV connection is model dependent by construction

- Implications on heavy new physics & validity of EFT is ***a posteriori***
- Depends on **sensitivity** & **energy scale** probed by data
- Bottom-up philosophy: new physics scale unknown

arbitrary dimensionful parameter $\frac{c}{f_a} = \frac{\lambda}{M}$ *coupling/mass scale of new physics*

In general:

- Larger coupling \Leftrightarrow higher new physics scale
- Better for EFT validity
- Some coefficients typically come with loop factors
- e.g. c_{VV} gauge interactions


UV interpretation

SMEFT-UV connection is model dependent by construction

- Implications on heavy new physics & validity of EFT is ***a posteriori***
- Depends on **sensitivity** & **energy scale** probed by data
- Bottom-up philosophy: new physics scale unknown

arbitrary dimensionful parameter $\frac{c}{f_a} = \frac{\lambda}{M}$ coupling/mass scale of new physics

constraint: $c/f_a < X$



In general:

- Larger coupling \Leftrightarrow higher new physics scale
- Better for EFT validity
- Some coefficients typically come with loop factors
- e.g. c_{VV} gauge interactions