

**53rd International Symposium
on Multiparticle Dynamics
(ISMD 2025)**

Report of Contributions

Contribution ID: 5

Type: **not specified**

Space-time geometry of small and large collision systems

Jet suppression has long been considered a signal for QGP formation in A-A nuclear collisions. Conventional approaches include a classical Glauber model to estimate the number of nucleon participants N_{part} and N-N binary collisions N_{bin} . A spectrum ratio RAA incorporating N_{bin} and assuming hard-component (jet-related hadrons) factorization is intended to measure the extent of high-pt jet suppression in A-A relative to p-p. The same approach applied to smaller collision systems (p-Pb) returns nontrivial but ambiguous results. Detailed analysis of p-Pb spectra reveals that classical Glauber Monte Carlo substantially overestimate N_{part} and N_{bin} . An alternative method based on ensemble-mean pt data is compatible with spectrum data. Several spectrum features call into question assumed hard-component (jet) factorization implicit in the RAA definition and reveal the importance of “exclusivity” (a projectile nucleon can only interact with one target nucleon *at a time*) and relativistic time dilation. An updated version of a two-component spectrum model (TCM) isolates complete spectrum hard components (jet fragments) down to low hadron pt without assuming factorization. Revised TCM analysis provides surprising new results and places p-Pb and Pb-Pb systems within a common descriptive context.

Author: TRAINOR, Thomas

Presenter: TRAINOR, Thomas

Session Classification: QCD evolution of dense media and minimum-bias jets

Track Classification: QCD evolution of dense media and minimum-bias jets