

FINESS2024: FInite temperature Non-Equilibrium Superfluid Systems

Sunday, 1 September 2024 - Thursday, 5 September 2024
Novotel Surfers Paradise, Gold Coast, Queensland, Australia

Book of Abstracts

Contents

Bose and Fermi Polarons in Atom —Ion Hybrid Systems	1
Imaginary gauge potentials in a non-Hermitian spin-orbit coupled quantum gas	1
Observation of Nonlinear Response and Onsager Regression in a Photon Bose-Einstein Condensate	2
Quantized Vortices in Superfluid Helium Thin Films	3
Collective excitations of a Bose-condensed gas: Fate of second sound in the crossover regime between hydrodynamic and collisionless regimes	3
Coherent fraction of an equilibrium condensate	4
Excitation of Bose-Einstein condensates through the separate field technique	5
Odd-frequency superfluidity from a particle-number-conserving perspective	6
Fractal spectrum and dimension extension of twisted bilayer optical lattices in ultracold atoms	7
Bose polarons in a box: universal features and the effects of finite temperature	7
Emergent Universal Drag Law in a Model of Superflow	8
How to deduce the entropy from atom-atom correlations	9
Microscopic many-body theory of two-dimensional coherent spectroscopy of exciton-polarons in one-dimensional materials	10
A charged impurity in an ultracold gas: observations of cold chemistry	11
Thermal fluctuations in multicomponent quantum gases	12
Dynamics and Thermodynamics of Rabi-driven Fermi gases	12
Vortex spin in a Bose-Einstein condensate	13
A nonequilibrium quantum Otto cycle in a one-dimensional Bose gas	14
Creating and Manipulating Dirac Strings in Spinor Condensate	15
Super Fermi polaron and Nagaoka ferromagnetism in a two-dimensional square lattice	16
Polaron approach to quantum mixtures	16

Observation of stationary turbulence in spinor Bose-Einstein condensates	17
Heteronuclear mixtures: From Efimov effect to heavy Fermi polarons	17
Towards Fermi polarons with heavy impurities	19
Photon BECs in dye-filled microcavities and VCSELs	20
Universal dynamics in strongly interacting Bose gases far from equilibrium	21
Vortices on rotating shell-shaped Bose-Einstein condensates	21
Self organisation and metastability of cavity bosons at very long times, beyond the adiabatic elimination approximation	22
Ferrodark solitons in a spinor superfluid: exact solutions, novel speed limit and anomalous dynamics	23
Hamiltonian engineering using Bragg matter-wave interferometers in an optical cavity .	24
Universal coarsening in 2D and 3D Bose gases	24
Instabilities in Bose-Einstein condensed mixtures, by perturbing initially immiscible con- figurations	25
Bose-Einstein condensation and lasing of low-dimensional semiconductor materials . . .	26
Designing Atomtronic Circuits via Superfluid Dynamics	27
Narrow-linewidth exciton-polariton laser	28
Work, heat and entropy in isolated quantum systems	28
Realising topological phases in the spin-1/2 quantum kicked rotor	29
Equatorial waves in rotating bubble-trapped superfluids	30
Shear-Induced Decaying Turbulence in Bose-Einstein Condensates	31
Exploring the dynamics of polar core vortices in homogeneous spin-1 Bose-Einstein con- densates	32
Vortex matter simulation of the one component plasma	32
On Demand Formation of Topological Defects in Ferromagnetic Spinor Bose Einstein Con- densates	33
Phononic crystal trapping geometries for improved rotational sensing with ultracold atoms	34
Tensor network methods for the Gross-Pitaevskii equation	35
False vacuum decay in an ultracold spin-1 Bose gas	35
An Efficient Quantum Phase-Space Method for Simulating Feedback Control of Interacting Many-Body Quantum Systems	36

Nonequilibrium Transport in a Superfluid Josephson Junction Chain: Is There Negative Differential Conductivity?	37
Non-Hermitian dynamics of a photonic wave packet in a microcavity filled with a liquid crystal	38
Shapiro steps in driven atomic Josephson junctions	39
Incoherent Transport in the detection of Majorana bound states	39
Signatures of many-body localization of quasiparticles in a flat band superconductor	40
Phase transitions and nonequilibrium dynamics in driven quantum matter	41
Exact Results of Fermi Polarons with Ultracold Atoms	41
Non-Hermitian band geometry and dynamics of exciton-polaritons	42
Galactic-Scale Superfluidity: True Macroscopic Condensation with Long-Range Interactions?	43
Observation of Shapiro Steps in an Atomic Superfluid	44
Reynolds similitude of a pure superfluid at low temperatures	45
Universal description of massive vortices in superfluids	45
(Non equilibrium) thermodynamics of classical Integrable models in their thermodynamic limit	46
Active matter in two dimensions	47
Medium-enhanced repulsion between polaron quasiparticles in a quantum gas	47
Tweezers, trapped ions, and Rydbergs: a quantum simulation zoo	48
Non-perturbative corrections to the weakly interacting two component Fermi gas	49
Bogoliubov theory of 1D anyons in a lattice	50
Unravelling Interaction and Temperature Contributions in Unpolarized Trapped Fermionic Atoms in the BCS Regime	50
Shear flow and vortex array instabilities in annular strongly-correlated atomic superfluids	51
Prescod-Weinstein	52
Driven-dissipative spinor superfluids: a compact Kardar-Parisi-Zhang dynamics of the phase	52
Quartet superfluid in mass-imbalanced ultracold Fermi mixtures	52
Wave breaking and multisoliton fission in a chip-scale superfluid waveflume	53

4

Bose and Fermi Polarons in Atom—Ion Hybrid Systems

Authors: R. Pessoa¹; S. Vitiello²; L. Peña Ardila³

¹ Instituto de Física, Universidade Federal de Goiás - UFG, 74001-970 Goiânia, GO, Brazil

² Instituto de Física Gleb Wataghin, Universidade Estadual de Campinas - UNICAMP, 13083-970 Campinas, SP, Brazil

³ Camerino University

Corresponding Author: luis.penaardila@unicam.it

We will discuss ionic polarons and their induced interaction created as a result of charged particles interacting with a Bose-Einstein condensate and a Fermi gas. Here we show that even in a comparatively simple setup consisting of charged impurities in a weakly interacting bosonic medium and an ideal Fermi gas with tunable atom-ion scattering length, the competition of length scales gives rise to a highly correlated mesoscopic state. Using quantum Monte Carlo simulations, we unravel its vastly different polaronic properties compared to neutral quantum impurities. Moreover, we identify a transition between the regime amenable to conventional perturbative treatment in the limit of weak atom-ion interactions and a many-body bound state with vanishing quasi-particle residue composed of hundreds of atoms. Contrary to the case of neutral impurities, ionic polarons can bound many excitations, forming a non-trivial interplay between few and many-body physics, changing the ground-state properties of the polaron radically. Finally, we investigate the specific case of two ions that mediate interactions via the bosonic bath. This interaction can be sizable with respect to the Coulomb interaction, giving rise to notable effects that may have direct consequences in the platform employed for future quantum technologies.

References:

Relevant publications (optional):

1. Charged impurities in a Bose-Einstein condensate: Many-body bound states and induced interactions. Grigory E. Astrakharchik and LAP Ardila, Krzysztof Jachymski & Antonio Negretti. Nat Comm. 14, 1647 (2023). <https://doi.org/10.1038/s41467-023-37153-0>
2. Quantum Behavior of a Heavy Impurity Strongly Coupled to a Bose Gas. Jesper Levinsen, LAP Ardila, Shuhei M. Yoshida, and Meera M. Parish. Phys. Rev. Lett 127, 033401 (2021) - Editors' suggestion. <https://doi.org/10.1103/PhysRevLett.127.033401>.
3. Quantum Droplets of Dipolar Mixtures. R.N. Bisset, LAP Ardila, and L. Santos. Phys. Rev. Lett 126, 025301 (2021). <https://doi.org/10.1103/PhysRevLett.126.025301>.

Career stage:

Professor

Short bio (50 words) or link to website:

<https://scholar.google.dk/citations?user=eYQv9fIAAAAJ&hl=en>

5

Imaginary gauge potentials in a non-Hermitian spin-orbit coupled quantum gas

Author: Ian Spielman^{None}

Corresponding Author: spielman@umd.edu

In 1996 Hatano and Nelson proposed a non-Hermitian lattice model containing an imaginary Peierls phase [Phys. Rev. Lett. **77** 570–573 (1996)], and subsequent analyses revealed it that is an instance of a new class of topological systems.

We experimentally realize a continuum analog to this model containing an imaginary gauge potential in a homogeneous spin-orbit coupled Bose-Einstein condensate (BEC).

The base spin-orbit coupled Hamiltonian is made non-Hermitian by adding tunable spin-dependent loss by microwave coupling to a subspace with spontaneous emission.

We find that the Heisenberg equations of motion for position and momentum with an imaginary gauge potential depend explicitly on the system's phase-space distribution.

In our experiment we first observed the non-Hermitian skin effect by localizing an initial state at the trap boundary, and verifying that it was stationary.

We then revealed collective nonreciprocal transport in real space, where the acceleration is a decreasing function of BECs spatial extent in agreement with non-Hermitian Gross-Pitaevskii equation simulations.

References:

Short bio (50 words) or link to website:

ultracolde.jqi.umd.edu

Relevant publications (optional):

Career stage:

Professor

6

Observation of Nonlinear Response and Onsager Regression in a Photon Bose-Einstein Condensate

Author: Nikolas Longen¹

Co-authors: Alexander Sazhin²; Andris Erglis³; Frank Vewinger²; Göran Hellmann⁴; Julian Schmitt²; Martin Weitz²; Michiel Wouters⁵; Vladimir N. Gladilin⁵

¹ *Institut für Angewandte Physik, Universität Bonn, Wegelerstr. 8, 53115 Bonn, Germany*

² *University of Bonn*

³ *University of Freiburg*

⁴ *Leibniz Institute of Photonic Technology Jena*

⁵ *University of Antwerps*

Corresponding Author: schmitt@iap.uni-bonn.de

The quantum regression theorem states that the correlations of a system at two different times are governed by the same equations of motion as the temporal response of the average values. Such a relation provides a powerful framework for the investigation of physical systems by establishing a formal connection between intrinsic microscopic behaviour and a macroscopic effect due to an external cause, allowing to determine e.g. structure factors. Here I report experiments demonstrating that the two-time second-order correlations of an photon Bose-Einstein condensate inside a dye-filled microcavity exhibit the same eigenvalues of the dynamics as the response of the condensate to a sudden perturbation of the dye molecule bath. This confirms an unconventional form of the regression theorem for a coupled many-body quantum system, where the perturbation acts on the bath and only the condensate response is monitored. For strong perturbations, we observe nonlinear relaxation dynamics back to the steady state which our microscopic theory relates to the equilibrium fluctuations, thereby extending the regression theorem beyond the regime of linear response. The demonstrated nonlinearity of the condensate-bath system paves the way for studies of novel elementary excitations in lattices of driven-dissipative photon condensates.

References:**Short bio (50 words) or link to website:**

<https://www.qfl.uni-bonn.de/en>

Relevant publications (optional):**Career stage:**

Student

Posters I / 7**Quantized Vortices in Superfluid Helium Thin Films**

Authors: Christopher Baker¹; Glen Harris²; Nicole Luu³; Warwick Bowen¹

¹ *The University of Queensland*

² *University of Queensland*

³ *The University of Queensland, Australia*

Corresponding Authors: uqnluu1@uq.edu.au, w.bowen@uq.edu.au, g.harris2@uq.edu.au, c.baker3@uq.edu.au

Quantized vortices are central to two-dimensional superfluidity and quantum turbulence. Though there is great interest in observing and understanding their behaviour, vortices in superfluid helium-4 are particularly challenging due to their Angstrom-sized cores and low refractive indices. I will present my work in the experimental exploration of vortex dynamics in thin films of superfluid helium by direct optical detection. This is achieved by cooling silicon photonic crystals to millikelvin temperatures, at which superfluid helium self-assembles into a nanometre-thick film along the surface of the crystal [1]. Advanced fabrication techniques enable the creation of high-quality silicon photonic crystals with small optical mode volumes that provide the ability to enhance interactions between light and quantized vortices. The presence of vortices creates a dimple in the superfluid film and shifts the resonance frequency of the optical cavity, providing a direct indication of the location of the vortices [2]. Going forward, we will be able to track the position of vortices with sub-nanometre resolution as well as employ optomechanical techniques to trap and control the vortices [3].

[1] W. W. Wasserman et al., *Opt. Express*, 30, 30822 (2022).

[2] Y. P. Sachkou et al., *Science* 366, 1480 (2019).

[3] X. He et al., *Nature Physics* 16, 4 (2020).

References:**Short bio (50 words) or link to website:**

I am in the first year of my PhD at The University of Queensland under supervision by Warwick Bowen and Chris Baker. I have previously worked with Dirk Bouwmeester at The University of California, Santa Barbara and Leiden University, The Netherlands.

Relevant publications (optional):**Career stage:**

Student

Posters I / 8

Collective excitations of a Bose-condensed gas: Fate of second sound in the crossover regime between hydrodynamic and collisionless regimes

Authors: Hoshu Hiyane¹; Shohei Watabe²; Tetsuro Nikuni³

¹ *Okinawa Institute of Science and Technology*

² *Faculty of Engineering, Computer Science and Engineering, Shibaura Institute of Technology*

³ *Department of Physics, Tokyo University of Science*

Corresponding Authors: watabe@shibaura-it.ac.jp, nikuni@rs.tus.ac.jp, hoshu-hiyane@oist.jp

We develop the moment method for Bose-Einstein condensates at finite temperatures that enable us to study collective sound modes from the hydrodynamic to the collisionless regime [1]. In particular, we investigate collective excitations in a weakly interacting dilute Bose gas by applying the moment method to the Zaremba-Nikuni-Griffin equation, which is the coupled equation of the Boltzmann equation with the generalized Gross-Pitaevskii equation. Utilizing the moment method, collective excitations in the crossover regime between the hydrodynamic and collisionless regimes are investigated in detail. In the crossover regime, the second sound mode loses the weight of the density response function because of the significant coupling with incoherent modes, whereas the first sound shows a distinct but broad peak structure. We compare the result obtained by the moment method with that of the Landau two-fluid equations and show that the collective mode predicted by the Landau two-fluid equations well coincides with the result from the moment method even far from the hydrodynamic regime, whereas clear distinction also emerges in the relatively higher momentum regime.

References:

[1] Hoshu Hiyane, Shohei Watabe, and Tetsuro Nikuni. Collective excitations of a Bose-condensed gas: Fate of second sound in the crossover regime between hydrodynamic and collisionless regimes. *Phys. Rev. A*, 109:033302, Mar 2024

Short bio (50 words) or link to website:

My name is Hoshu Hiyane, a fourth year PhD student at the Quantum Systems Unit at the Okinawa Institute of Science and Technology. I received my bachelor's degree at Tokyo University of Science. I will talk about my recent work on the dynamical properties of finite-temperature Bose-Einstein condensates.

Relevant publications (optional):

Career stage:

Student

9

Coherent fraction of an equilibrium condensate

Authors: David W Snoke¹; Hassan Alnatah¹; Jonathan Beaumariage¹; Kenneth W West²; Kirk Baldwin²; Loren N Pfeiffer²; Man Chun Tam³; Qi Yao¹; Shouvik Mukherjee⁴; Zbigniew Wasilewski³

¹ *University of Pittsburgh*

² *Princeton University*

³ *University of Waterloo*

⁴ *University of Maryland and National Institute*

Corresponding Author: haa108@pitt.edu

We report recent progress on the measurement of the coherent fraction of a two-dimensional Bose gas in thermal equilibrium. We have created a homogeneous exciton-polariton gas in equilibrium, realizing the textbook paradigm of a uniform Bose Gas in two-dimensions. Under these conditions, we have measured the coherent fraction of this Bose gas from very low density up to density well above the condensation threshold. These measurements reveal a consistent power law for the coherent fraction over nearly three orders of its magnitude. The same power law is seen in numerical simulations solving the two-dimensional Gross-Pitaevskii equation for the equilibrium coherence; these simulations also show that the power law corresponds to the coherence length in the system growing with a power law of 1.6 as a function of the total density. This power law has not been predicted by prior analytical theories.

This work has been supported by the National Science Foundation through Grant DMR-2306977.

References:

Short bio (50 words) or link to website:

I am a fifth-year graduate student working with Dr. David Snoke at the University of Pittsburgh.

Relevant publications (optional):

<https://arxiv.org/abs/2308.05100>

<https://www.science.org/doi/full/10.1126/sciadv.adi6762>

Career stage:

Student

Posters II / 10

Excitation of Bose-Einstein condensates through the separate field technique

Authors: Leandro Alvares Machado^{None}; Lucas Madeira¹; Mônica Caracanhas¹; Vanderlei Salvador Bagnato¹

¹ *Universidade de São Paulo*

Corresponding Authors: madeira@ifsc.usp.br, caracanhas@ifsc.usp.br, lemachado@usp.br, vander@ifsc.usp.br

In this work, we propose a new approach to excite a Bose-Einstein condensate by introducing perturbations to the trapping potential through the separate fields technique, originally introduced by Norman F. Ramsey. We employed the variational method based on the Thomas-Fermi wave function Ansatz to obtain dynamical equations governing the radii of a Bose-Einstein condensate when under external perturbations. The obtained dynamical equations allow us to construct the resonance curves for the collective modes of Bose-Einstein condensates, specifically focusing on the dipole, quadrupole and breathing modes. We observed that depending on whether the external signal is modulated by the Rabi or Ramsey procedure, the system behaves differently due to its inherent coherence, that reflects on the resonance curve of the collective modes, giving rise to the Ramsey fringes in a clear interference pattern, enabling a more precise frequency measurement. We introduced an alternative model to describe the dynamics of collective modes based on a three-level system, which proved to be consistent with the variational model. Moreover, the three-level model provided a framework for fostering the idea of coherent control by manipulating the population associated with collective modes through the application of perturbation pulses. Finally, to present

ongoing analyses and their perspectives, we used the Rabi excitation protocol to study how the external modulation leads to specific routes in terms of generation of quantum turbulence.

References:

Short bio (50 words) or link to website:

Currently PhD student in the field of non-equilibrium superfluid at the Instituto de Física de São Paulo as part of the Universidade de São Paulo. Master's degree in theoretical and experimental physics and bachelor in Physics for the same institution.

Relevant publications (optional):

<https://journals.aps.org/prresearch/abstract/10.1103/PhysRevResearch.5.043081>

Career stage:

Student

11

Odd-frequency superfluidity from a particle-number-conserving perspective

Authors: Joachim Brand¹; Kadin Thompson²; Michele Governale²; Uli Zülicke²

¹ *Massey University*

² *Victoria University of Wellington*

Corresponding Authors: uli.zuelicke@vuw.ac.nz, michele.governale@vuw.ac.nz, kadinthompson@gmail.com, j.brand@massey.ac.nz

We investigate odd-in-time—or *odd-frequency*—pairing of fermions in equilibrium systems within the particle-number-conserving framework of Penrose, Onsager and Yang, where superfluid order is defined by macroscopic eigenvalues of reduced density matrices. We show that odd-frequency pair correlations are synonymous with even fermion-exchange symmetry in a time-dependent correlation function that generalises the two-body reduced density matrix [1]. Macroscopic even-under-fermion-exchange pairing is found to emerge from conventional Penrose-Onsager-Yang condensation in two-body or higher-order reduced density matrices through the symmetry-mixing properties of the Hamiltonian. We identify and characterise a *transformer* matrix responsible for producing macroscopic even fermion-exchange correlations that coexist with a conventional Cooper-pair condensate, while a *generator* matrix is shown to be responsible for creating macroscopic even fermion-exchange correlations from hidden orders such as a multi-particle condensate. The transformer scenario is illustrated using the spin-imbalanced Fermi superfluid as an example. The generator scenario is demonstrated by the composite-boson condensate arising for itinerant electrons coupled to magnetic excitations. Structural analysis of the transformer and generator matrices is shown to provide general conditions for odd-frequency pairing order to arise in a given system.

References:

[1] Thompson, K., U. Zülicke, M. Governale, and J. Brand. “Odd-Frequency Superfluidity from a Particle-Number-Conserving Perspective.” arXiv, March 10, 2024. <https://doi.org/10.48550/arXiv.2403.06325>.

Short bio (50 words) or link to website:

Professor Brand completed obtained his PhD from the University of Heidelberg. After a postdoctoral research fellowship at the University of Washington and the Max Planck Institute for the Physics of Complex Systems he joined Massey University in 2006.

Relevant publications (optional):

Thompson, K., U. Zülicke, M. Governale, and J. Brand. "Odd-Frequency Superfluidity from a Particle-Number-Conserving Perspective." arXiv, March 10, 2024. <https://doi.org/10.48550/arXiv.2403.06325>.

Reynolds, L. A., E. Schwartz, U. Ebling, M. Weyland, J. Brand, and M. F. Andersen. "Direct Measurements of Collisional Dynamics in Cold Atom Triads." *Physical Review Letters* 124, no. 7 (February 18, 2020): 073401. <https://doi.org/10.1103/PhysRevLett.124.073401>.

Yang, Mingrui, Matija Čufar, Elke Pahl, and Joachim Brand. "Polaron-Depletion Transition in the Yrast Excitations of a One-Dimensional Bose Gas with a Mobile Impurity." *Condensed Matter* 7, no. 1 (January 26, 2022): 15. <https://doi.org/10.3390/condmat7010015>.

Career stage:

Professor

Posters II / 12**Fractal spectrum and dimension extension of twisted bilayer optical lattices in ultracold atoms****Author:** Zheyu Shi^{None}**Co-authors:** Ce Wang ; Chao Gao ; Hui Zhai ; Jing Zhang ; Xu-Tao Wan**Corresponding Author:** zheyushi@gmail.com

The experimental realization of twisted bilayer optical lattices in ultracold atomic gases (*Nature* 615, 231 (2023)) has paved the way towards the investigation of moiré physics in cold quantum gases. I will present two recent theoretical works in this system. In the first work (arXiv:2404.08211), we point out that the geometric moiré effect can induce fractal band structures. The fractals are controlled by the twist angle between two monolayers and are closely connected to the celebrated butterfly spectrum of two-dimensional Bloch electrons in a magnetic field. We also provide numerical evidence on the infinite recursive structures of the spectrum and give an algorithm for computing these structures. In the second work (2404.19608), we propose that by utilizing the current, it is possible to construct a twisted three-dimensional optical lattice in ultracold atomic gases, which extends the moiré physics to higher dimensions. It is worth emphasizing that such lattices cannot be realized in condensed-matter systems, for it is impractical to overlay two pieces of three-dimensional solid-state material. We develop the general theory describing the commensurate conditions, lattice structures, and crystalline symmetries of these lattices. We highlight the fundamental difference between moiré physics in two and three dimensions. That is, three-dimensional moiré lattices can possess more versatile crystalline structures because of the non-commutative nature of $SO(3)$.

References:<https://arxiv.org/abs/2404.08211><https://arxiv.org/abs/2404.19608>**Short bio (50 words) or link to website:**https://faculty.ecnu.edu.cn/_s29/szy_en/main.psp**Relevant publications (optional):****Career stage:**

Professor

Bose polarons in a box: universal features and the effects of finite temperature

Authors: Jiri Etrych¹; Gevorg Martirosyan¹; Alec Cao²; Simon Fischer¹; Seb Morris¹; Christopher Ho¹; Zoran Hadzibabic^{None}; Christoph Eigen¹

¹ *University of Cambridge*

² *JILA*

Corresponding Authors: smf61@cam.ac.uk, sm2357@cam.ac.uk, zh10001@cam.ac.uk, alec.cao@colorado.edu, cjh203@cam.ac.uk, gm572@cam.ac.uk, je382@cam.ac.uk, ce330@cam.ac.uk

An impurity immersed in a quantum bath is a fundamental setting in many-body physics that, in spite of its apparent simplicity, features complex emergent behaviour. I will present our recent experiments in which we measure the spectral properties and real-time dynamics of mobile impurities injected into a homogeneous Bose–Einstein condensate (BEC), using two Feshbach resonances to tune both the impurity-bath and intrabath interactions. We map out the attractive and repulsive branches of polaron quasiparticles and explore the breakdown of the quasiparticle picture for near-resonant interactions. On the repulsive side of the resonance, we resolve both the repulsive polaron and the molecular state associated with the Feshbach resonance in the strongly interacting regime and show that the latter also has a many-body character. Our measurements reveal remarkably universal behavior, controlled by the bath density and a single dimensionless interaction parameter, with no significant dependence on the intrabath interactions. Finally, I will also present an extension of our study to finite temperatures both below and above the BEC phase transition of the bath. In particular, we find that many-body effects are suppressed as the temperature of the bath is increased, which can lead to counterintuitive narrowing of spectral features near the resonance.

References:

Short bio (50 words) or link to website:

2016-2020 University of Cambridge, BA, MSci Natural Sciences (Physics)
2020-2024 University of Cambridge, PhD with Zoran Hadzibabic (current position)
2024- I will be staying as a postdoc with Zoran from the autumn

Career stage:

Student

Relevant publications (optional):

1. J. Etrych, G. Martirosyan, A. Cao, C. Ho, Z. Hadzibabic, and C. Eigen, Universal quantum dynamics of Bose polarons, arxiv.org/pdf/2402.14816
2. G. Martirosyan, C. J. Ho, J. Etrych, Y. Zhang, A. Cao, Z. Hadzibabic, and C. Eigen, Observation of Subdiffusive Dynamic Scaling in a Driven and Disordered Bose Gas, *Phys. Rev. Lett.* 132, 113401 (2024)
3. L. H. Dogra, G. Martirosyan, T. A. Hilker, J. A. P. Glidden, J. Etrych, A. Cao, C. Eigen, R. P. Smith, and Z. Hadzibabic, Universal equation of state for wave turbulence in a quantum gas, *Nature* 620, 521 (2023)

Posters I / 14

Emergent Universal Drag Law in a Model of Superflow

Authors: Arghavan Safavi-Naini¹; Halina Rubinsztein-Dunlop^{None}; Maarten Christenhusz²; Matt Reeves²; Tyler Neely²

¹ *University of Amsterdam*

² *University of Queensland*

Corresponding Authors: m.reeves@uq.edu.au, m.christenhusz@uq.edu.au, hrubinszteindunlop@gmail.com, a.safavinaini@uva.nl, t.neely@uq.edu.au

Despite the fundamentally different dissipation mechanisms, many laws and phenomena of classical turbulence equivalently manifest in quantum turbulence. The Reynolds law of dynamical similarity states that two objects of same geometry across different length scales are hydrodynamically equivalent under the same Reynolds number, leading to a universal drag coefficient law. We confirm the existence of a universal drag law in a superfluid wake, facilitated by the nucleation of quantized vortices. We study superfluid flow across a range of Reynolds numbers for the paradigmatic classical hard-wall and the Gaussian obstacle, popular in experimental quantum hydrodynamics. In addition, we provide a feasible method for measuring superfluid drag forces in an experimental environment using control volumes.

References:

Short bio (50 words) or link to website:

Maarten is a PhD student in the Bose-Einstein Condensation Laboratory at the University of Queensland. He joined the group during his Masters, where he set up a two-dimensional optical trap for Bose-Einstein condensates to study quantum turbulence. Currently, Maarten is working on drag phenomena in superfluids.

Relevant publications (optional):

Career stage:

Student

15

How to deduce the entropy from atom-atom correlations

Author: Karen Kheruntsyan^{None}

Co-authors: Raymon Watson¹; Caleb Coleman¹

¹ *University of Queensland*

Corresponding Authors: caleb.coleman@uq.net.au, karen.kheruntsyan@uq.edu.au, raymon.watson@uq.edu.au

We derive a thermodynamic Maxwell relation by which the entropy of an ultracold atomic gas can be deduced from the local (same point) atom-atom correlation function. The Maxwell relation in question is applicable to many-body systems with short-range interactions that can be characterised by *s*-wave scattering. For such systems, the local atom-atom correlation function represents a thermodynamic quantity that can be calculated from the Helmholtz free energy using the Hellmann-Feynman theorem [1]. Here, we exploit this property to derive a Maxwell relation that relates the atom-atom correlation to the thermodynamic entropy of the system [2]. As a practical application of this Maxwell relation, we utilise it to calculate the entropy of a weakly interacting one-dimensional (1D) Bose gas from its atom-atom correlations in the context of the *c*-field approach of the stochastic projected Gross-Pitaevskii equation (SPGPE). The SPGPE is a well established and widely used numerical approach for computing thermal equilibrium and dynamical properties of finite temperature Bose gases, such as partially condensed Bose-Einstein condensates in 2D and 3D, or phase-fluctuating quasicondensates in 1D. Despite its wide applicability to ultracold quantum gas systems, computing the entropy of such systems within the SPGPE has not been accomplished prior to this work. Our calculations can also be viewed as a numerical experiment that serves as a proof-of-principle demonstration of an experimental method to deduce the entropy of an ultracold quantum gas from the measurements of atom-atom correlations.

References:

- [1] K. V. Kheruntsyan, D. M. Gangardt, P. D. Drummond, and G. V. Shlyapnikov, Phys. Rev. Lett. 91, 040403 (2003).
 [2] R. S. Watson, C. Coleman, and K. V. Kheruntsyan, arXiv:2405.04159 (2024).

Short bio (50 words) or link to website:

<https://people.smp.uq.edu.au/KarenKheruntsyan/index.html>

Relevant publications (optional):**Career stage:**

Professor

Posters I / 16

Microscopic many-body theory of two-dimensional coherent spectroscopy of exciton-polarons in one-dimensional materials

Author: Jia Wang¹

Co-authors: Hui Hu ; Xia-Ji Liu ¹

¹ *Swinburne University of Technology*

Corresponding Authors: xiajiliu@swin.edu.au, jiawanghome@gmail.com, hhu@swin.edu.au

We have developed a microscopic many-body theory for two-dimensional coherent spectroscopy (2DCS) of polarons in one-dimensional (1D) materials [1]. Our theory incorporates contributions from three processes: excited-state emission (ESE), ground-state bleaching (GSB), and excited-state absorption (ESA). While ESE and GSB contributions can be accurately described using Chevy's ansatz with one particle-hole excitation, the ESA process requires information about many-body eigenstates involving two impurities. To address this, we have extended Chevy's ansatz to include double polaron states and verified its validity by comparing our results with exact calculations using Bethe's ansatz.

Our numerical results indicate that in the weak interaction limit, the ESA contribution cancels out the total ESE and GSB contributions, leading to less prominent spectral features. However, under strong interactions, the features of the ESA contribution and the combined ESE and GSB contributions remain observable in the 2DCS spectra, providing valuable insights into polaron interactions. Additionally, we have examined the mixing time dynamics, which characterize the quantum coherences of polaron resonances. Overall, our theory offers a comprehensive framework for understanding and interpreting 2DCS spectra of polarons in 1D materials, shedding light on their interactions and coherent dynamics.

[1] Jia Wang , Hui Hu , and Xia-Ji Liu, PRB 109, 205414 (2024)

References:

- [1] Jia Wang , Hui Hu , and Xia-Ji Liu, PRB 109, 205414 (2024)

Short bio (50 words) or link to website:

I am an ARC Future Fellow working in the Centre for Quantum Technology Theory (CQTT) at Swinburne University. Prior to my current role, I held postdoctoral positions at both Swinburne and the University of Connecticut Storrs. I earned my Ph.D. from the University of Colorado, Boulder.

Relevant publications (optional):

- [1] PRB 109, 205414 (2024) <https://doi.org/10.1103/PhysRevB.109.205414>
 [2] PRA 107, 013305 (2023) <https://doi.org/10.1103/PhysRevA.107.013305>
 [3] PRL 128, 175301 (2022) <https://doi.org/10.1103/PhysRevLett.128.175301>

Career stage:

Postdoc

Posters I / 17**A charged impurity in an ultracold gas: observations of cold chemistry**

Authors: Arghavan Safavi-Naini¹; Egor Kovlakov¹; Eleanor Trimby¹; Henrik Hirzler¹; Jesus Perez-Rios²; Rene Gerritsma¹; Rianne Lous¹

¹ *University of Amsterdam*

² *Stony Brook University*

Corresponding Authors: a.safavinaini@uva.nl, eleanor.trimby@outlook.com

Hybrid ion-atom systems combine the benefits of a single, well-controlled ion with those of a many-body quantum gas, offering prospects for quantum simulation, ultracold chemistry, and charged impurity physics [1, 2]. For the latter, the longer range of the atom-ion interaction, compared to that between two neutrals, is expected to give rise to interesting behaviour, for example the formation of a much larger polaron than in studies of neutral impurities. Furthermore, the excellent degree of control available over a single charged impurity has prospects for its use as a sensitive probe to give information about the bath in which it is immersed. It has been suggested that studies of few-body chemical processes in atom-ion experiments could provide valuable insight into the many-body behaviour of this system [3]. In this talk, I will present recent observations of chemical reactions between a single Yb⁺ ion and Li₂ dimers in an ultracold cloud, leading to the formation of a LiYb⁺ molecular ion [4]. We find this to be an unexpected example of the ion probing the atom cloud for trace quantities of dimers. Furthermore, I will outline recent experimental upgrades used to manipulate our atom cloud, increasing its density and preparing a new mixture of spin states. These upgrades have prospects for future studies of a single ionic impurity in a bath cooled to degeneracy, in particular one that is in the BEC-BCS crossover regime.

References:

- [1] M. Tomza et al., *Rev. Mod. Phys.* 91, 035001 (2019).
 [2] R. S. Lous and R. Gerritsma, *Adv. At. Mol. Opt. Phys.* 71 (2022).
 [3] J. Pérez-Ríos, *Mol. Phys.* 119, 8 (2021).
 [4] H. Hirzler et al., *Phys. Rev. Lett.* 128, 103401 (2022).

Short bio (50 words) or link to website:

For my PhD, I work on experiments in which a single trapped ion is immersed in a cloud of ultracold neutral atoms. My focus is on optimized cooling of the system to energies in which quantum effects begin to dominate, as well as observations of cold chemistry.

Relevant publications (optional):

Observation of chemical reactions between a trapped ion and ultracold Feshbach dimers, H. Hirzler, R.S. Lous, E. Trimby, J. Pérez-Ríos, A. Safavi-Naini and R. Gerritsma, *Phys. Rev. Lett.* 128, 103401 (2022). <https://doi.org/10.1103/PhysRevLett.128.103401>

Buffer gas cooling of ions in time-dependent traps using ultracold atoms, E. Trimby, H. Hirzler, H. Fürst, A. Safavi-Naini, R. Gerritsma and R. S. Lous, *New Journal of Physics* 24, 035004 (2022). <https://doi.org/10.1088/1367-2630/ac5759>

Trap-assisted complexes in cold atom-ion collisions, H. Hirzler, E. Trimby, R. Gerritsma, A. Safavi-Naini and J. Pérez-Ríos, *Phys. Rev. Lett.* 130, 143003 (2023). <https://doi.org/10.1103/PhysRevLett.130.143003>

Career stage:

Student

18

Thermal fluctuations in multicomponent quantum gases**Author:** Arko Roy¹¹ *IIT Mandi***Corresponding Author:** arko@iitmandi.ac.in

In the study of various quantum phase transitions and excitations in Bose-Einstein condensates (BECs), several key findings emerge across different scenarios. For a spin-orbit-coupled quasi-one-dimensional BEC, temperature-induced transitions from a superfluid plane-wave phase to a super-solid phase are observed, contrasting with the behavior in homogeneous systems. The Hartree-Fock-Bogoliubov theory with the Popov approximation is utilized to analyze collective excitation spectra, particularly focusing on the softening of the spin-dipole mode near the quantum critical point. Furthermore, the collective excitations of a spin-orbit-coupled spin-1 BEC in a trapping potential are studied theoretically at both zero and finite temperatures. Density and spin excitations exhibit distinct behaviors with temperature variations, notably affected by spin-orbit coupling.

Transitioning to a two-dimensional coherently-coupled Bose mixture, a paramagnetic-ferromagnetic quantum phase transition at zero temperature is identified, with subsequent investigations into its behavior at finite temperatures. Stochastic Gross-Pitaevskii formalism is employed, revealing a linear shift of the critical point with temperature and power-law scaling of critical quantities, consistent with thermal critical exponents.

Finally, the role of thermal fluctuations in two-dimensional binary Bose mixtures is explored, particularly focusing on the miscible-immiscible transition. Mean-field Hartree-Fock theory predicts a transition instability at non-zero temperatures due to divergent behavior in spin susceptibility, which is partially confirmed by numerical simulations. However, discrepancies between theory and simulations suggest that mean-field approximations struggle to accurately describe the system near the transition, especially concerning thermal fluctuations.

References:

Phys. Rev. A 107, 043301 (2023); Phys. Rev. Lett. 130, 220403 (2023); Phys. Rev. A 109, 033319 (2024); Phys. Rev. Research 3, 013161 (2021)

Short bio (50 words) or link to website:

<https://faculty.iitmandi.ac.in/~arko/>

Relevant publications (optional):

Phys. Rev. A 107, 043301 (2023)[<https://journals.aps.org/pr/abstract/10.1103/PhysRevA.107.043301>]; Phys. Rev. Lett. 130, 220403 (2023) [<https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.130.220403>]; Phys. Rev. A 109, 033319 (2024)[<https://journals.aps.org/pr/abstract/10.1103/PhysRevA.109.033319>]

Career stage:

Professor

Dynamics and Thermodynamics of Rabi-driven Fermi gases

Authors: Brendan Mulkerin^{None}; Jesper Levinsen¹; Meera Parish^{None}; Olivier Bleu¹

¹ *Monash University*

Corresponding Authors: brendan.mulkerin@monash.edu, jesper.levinsen@monash.edu, meera.parish@monash.edu, olivier.bleu@monash.edu

In this work we present our investigations on Rabi coupled Fermi gases. Specifically, the behavior of a mobile spin-1/2 impurity atom immersed in a Fermi gas, where the interacting spin- \uparrow and non-interacting spin- \downarrow states of the impurity are Rabi coupled via an external field. This scenario resembles the classic problem of a two-state system interacting with a dissipative environment but with an added dimension provided by the impurity momentum degree of freedom. In this context, the impurity can become “dressed” by excitations of the Fermi sea to form a Fermi polaron quasiparticle.

For the steady-state system, where the impurity has thermalized with the medium, we derive exact thermodynamic relations that connect the impurity magnetization with quasiparticle properties such as the number of fermions in the dressing cloud. We show how the thermodynamic properties evolve with increasing Rabi coupling and present exact analytical results in the limits of weak and strong Rabi coupling.

For the dynamics of the Rabi-driven Fermi polaron, we formulate a theoretical approach based on correlation functions that respects conservation laws and allows the efficient calculation of Rabi oscillations over a range of time scales and impurity momenta beyond what has been previously achieved. Our results are in good agreement with recent experiments on the Rabi oscillations of the attractive polaron, revealing how these oscillations are influenced by the interplay between the polaron and its dressing cloud.

References:

Short bio (50 words) or link to website:

Graduated with PhD from Melbourne University with Prof. Andy Martin, work as a postdoctoral researcher at Swinburne University with Profs. Xia-Ji Liu and Hui Hu, is a current postdoctoral researcher at Monash University with Pro. Meera Parish and Assoc. Prof Jesper Levinsen.

Relevant publications (optional):

Rabi oscillations and magnetization of a mobile spin-1/2 impurity in a Fermi sea
Brendan C. Mulkerin, Jesper Levinsen, and Meera M. Parish
Phys. Rev. A 109, 023302 (2024)

Crossover from exciton polarons to trions in doped two-dimensional semiconductors at finite temperature
Antonio Tiene, Brendan C. Mulkerin, Jesper Levinsen, Meera M. Parish, and Francesca Maria Marchetti
Phys. Rev. B 108, 125406 (2023)

Exact Quantum Virial Expansion for the Optical Response of Doped Two-Dimensional Semiconductors
Brendan C. Mulkerin, Antonio Tiene, Francesca Maria Marchetti, Meera M. Parish, and Jesper Levinsen
Phys. Rev. Lett. 131, 106901 (2023)

Career stage:

Postdoc

Posters I / 20

Vortex spin in a Bose-Einstein condensate

Author: Tapio Simula^{None}**Co-authors:** Emil Génétay Johansen¹; Chris Vale²¹ *Silicon Quantum Computing, UNSW*² *CSIRO***Corresponding Author:** tsimula@swin.edu.au

General relativity predicts that the curvature of spacetime induces spin rotations on a parallel transported particle. We deploy Unruh's analogue gravity picture and consider a quantised vortex embedded in a two-dimensional superfluid Bose-Einstein condensate. We show that such a vortex behaves dynamically like a charged particle with a spin in a gravitational field in a 2+1 dimensional spacetime [1-3]. The way the fermionic, split-boson, quasiparticle character of the vortex particle emerges out of bosons trapped by the vortices parallels the emergence of Majorana quasiparticles as split-fermions in the vortex cores of topological Fermi superfluids.

References:

[1] Emil Génétay Johansen, Tapio Simula, Vortex spin in a superfluid, arXiv:2305.16016.

[2] E. Génétay Johansen, C. Vale, T. Simula, Quantum double structure in cold atom superfluids, *AVS Quantum Sci.* 5, 033201 (2023).[3] Emil Génétay Johansen, Tapio Simula, Topological quantum computation using analog gravitational holonomy and time dilation, *SciPost Phys. Core* 6, 005 (2023).**Short bio (50 words) or link to website:**

<https://www.swinburne.edu.au/research/our-research/access-our-research/find-a-researcher-or-supervisor/researcher-profile/?id=tsimula>

Relevant publications (optional):

T. Simula, N. Kjærgaard, T. Pfau, Topological transport of a classical droplet in a lattice of time, arXiv:2403.06500.
<https://arxiv.org/abs/2403.06500>

Emil Génétay Johansen, Tapio Simula, Vortex spin in a superfluid, arXiv:2305.16016. <https://arxiv.org/abs/2305.16016>

Rama Sharma, Tapio P Simula, Andrew J Groszek, Fluctuation theorem anomaly in a point-vortex fluid, arXiv:2308.03397.
<https://arxiv.org/abs/2308.03397>

Career stage:

Professor

Posters I / 21**A nonequilibrium quantum Otto cycle in a one-dimensional Bose gas****Author:** Raymon Watson^{None}**Co-author:** Karen Kheruntsyan**Corresponding Authors:** karen.kheruntsyan@uq.edu.au, raymon.watson@uq.edu.au

Theoretical study of nonequilibrium quantum thermodynamics in many-body interacting systems is typically restricted by the complexity of dynamical simulation. Integrable systems, despite their exact solutions, are often no exception to this. However, the recently developed theory of generalized hydrodynamics (GHD) is capable of capturing the large-scale dynamics of integrable and near-integrable models in parameter regimes not accessible through alternative methods. We utilize this

recently developed theory of GHD, applying it to the study of a nonequilibrium quantum thermodynamic device. In particular, we analyse a quantum Otto cycle driven by control over interparticle interactions in an experimentally realistic one-dimensional Bose gas, which can be described by the integrable Lieb-Liniger model in the uniform limit and is nearly integrable in the harmonically trapped configuration. We explore the performance of this Otto engine cycle across the model's rich parameter space—from weak to strong interactions and at temperatures below and above quantum degeneracy. Further, we express the engine performance through a direct link with Glauber's second-order correlation function, which allows both net work and efficiency to be expressed analytically in various regions of the parameter space. Overall, the theory of GHD allows us to study a realistic finite-time operation of this quantum thermodynamic cycle and hence to understand the crossover between the previously studied idealised limits of instantaneous and quasistatic quenches.

References:

Short bio (50 words) or link to website:

I completed my Masters of Experimental Physics at the University of Western Australia under Associate Professor John McFerran, and am soon to complete my PhD under Professor Karen Kheruntsyan at the university of Queensland.

Relevant publications (optional):

Career stage:

Student

22

Creating and Manipulating Dirac Strings in Spinor Condensate

Author: Han Pu¹

¹ *Rice University*

Corresponding Author: hpu@rice.edu

Artificial monopoles have been engineered in various systems, yet there has been no systematic study of the singular vector potentials associated with the monopole field. We show that the Dirac string, the line singularity of the vector potential associated with the monopole field, can be engineered, manipulated, and made manifest in a spinor atomic condensate. We elucidate the connection among spin, orbital degrees of freedom, and the artificial gauge, and show that there exists a mapping between the vortex filament and the Dirac string. We also devise a proposal where preparing initial spin states with relevant symmetries and then adiabatically turn on the effective monopole field can result in different vortex patterns, revealing an underlying correspondence between the internal spin states and the spherical vortex structures [1]. Such a mapping also leads to a new way of constructing spherical Landau levels, and monopole harmonics. Our observation provides insights into the behavior of quantum matter possessing internal symmetries in curved spaces.

[1] arXiv:2402.14705

References:

Short bio (50 words) or link to website:

Han Pu obtained his Ph.D. in physics in 1999 and is currently a professor at Rice University. His research interest lies in theoretical cold atoms, quantum optics and quantum many-body physics. He is a Fellow of the American Physical Society.

Relevant publications (optional):

arXiv:2402.14705

Career stage:

Professor

23

Super Fermi polaron and Nagaoka ferromagnetism in a two-dimensional square lattice**Author:** Xia-Ji LIU¹¹ *Swinburne***Corresponding Author:** xiajiliu@swin.edu.au

We address the Fermi polaron physics of an impurity hopping around a two-dimensional square lattice and interacting with a sea of fermions at given filling factor. When the interaction is attractive, we find standard Fermi polaron quasiparticles, categorized as attractive polarons and repulsive polarons. When the interaction becomes repulsive, interestingly, we observe an unconventional highly-excited polaron quasiparticle, sharply peaked at the corner of the first Brillouin zone. This super Fermi polaron branch arises from the dressing of the impurity's motion with holes, instead of particles of fermions. We show that super Fermi polarons become increasingly well-defined with increasing impurity-fermion repulsions and might be considered as a precursor of Nagaoka ferromagnetism, which would appear at sufficiently large repulsions and at large filling factors. We also investigate the temperature-dependence of super Fermi polarons and find that they are thermally robust against the significant increase in temperature.

References:

H. Hu, J. Wang, X-J. Liu,
Super Fermi polaron and Nagaoka ferromagnetism in a two-dimensional square lattice, arXiv:2312.00385

Short bio (50 words) or link to website:

<https://www.swinburne.edu.au/research/our-research/access-our-research/find-a-researcher-or-supervisor/researcher-profile/?id=xiajiliu>

Relevant publications (optional):**Career stage:**

Professor

Posters I / 24**Polaron approach to quantum mixtures****Authors:** Jesper Levinsen¹; Meera Parish^{None}; Olivier Bleu¹¹ *Monash University***Corresponding Authors:** olivier.bleu@monash.edu, meera.parish@monash.edu, jesper.levinsen@monash.edu

The polaron, a particle dressed by excitations of a quantum medium, has been extensively studied in ultracold atomic gases. It represents the ultimate limit of imbalanced populations in quantum mixtures, and as such has relevance to the phase diagram of a wide range of systems, such as Fermi-Fermi, Bose-Bose, and Bose-Fermi gases. Here, I will present a variational approach to quantum mixtures which is inspired by highly successful variational approaches to the polaron problem.

References:**Short bio (50 words) or link to website:**

<https://research.monash.edu/en/persons/jesper-levinsen>

Relevant publications (optional):**Career stage:**

Professor

25

Observation of stationary turbulence in spinor Bose-Einstein condensates

Author: Yong-il Shin¹

¹ *Seoul National University*

Corresponding Author: yishin@snu.ac.kr

Spinor Bose-Einstein condensates (BECs) of atomic gases represent a quantum fluid characterized by multiple symmetry breaking, providing an interesting platform for the exploration of quantum turbulence. In this talk, I will report our observation of a stationary turbulent state in a spin-1 atomic BEC driven by a radio-frequency magnetic field. The magnetic driving injects energy into the system through spin rotation, leading to the emergence of an irregular spin texture in the condensate. As the driving persists, the spinor condensate evolves into a nonequilibrium steady state marked by distinctive spin turbulence. Remarkably, under specific driving conditions, the turbulence attains its maximum intensity, accompanied by an isotropic spin composition. Through numerical simulations and experimental validation, we find that the turbulence in the BEC is sustained by a mechanism rooted in the chaotic nature of internal spin dynamics induced by the magnetic driving.

References:**Short bio (50 words) or link to website:**

<https://qgl.snu.ac.kr/>

Relevant publications (optional):**Career stage:**

Professor

Heteronuclear mixtures: From Efimov effect to heavy Fermi polarons

Author: Michael Rautenberg¹

Co-authors: Tobias Krom¹; Tilman Enss²; Lauriane Chomaz¹; Matthias Weidemüller¹

¹ *PI Uni Heidelberg*

² *ITP Uni Heidelberg*

Corresponding Author: rautenberg@physi.uni-heidelberg.de

Mixtures of different atomic gases with largely different masses are particularly suited to observe the Efimov effect, a series of three-body bound states obeying a universal scaling law [1, 2]. Indeed, these few-body states have been observed in a thermal mixture of ${}^6\text{Li}$ and ${}^{133}\text{Cs}$ [3]. Considering this mixture now at much colder temperatures and in the limit where Cs atoms act as impurities in a degenerate Fermi sea of Li, the system can be described as quasi-particles known as Fermi polarons. Extending the work of [4], we predict that signatures of the Efimov effect still persist in this scenario where it manifests itself as resonances in the induced impurity interactions at the positions where the Efimov states would cross the scattering continuum [5].

Currently we are working towards the experimental observation of heavy Fermi polarons in the Li-Cs mixture. Besides the aforementioned connections to Efimov physics, this platform enables studies of Anderson orthogonality catastrophe [6]. Most prominently, a universal power law is expected in the real-time response revealed e.g. by Ramsey spectroscopy on the impurity atoms [7]. Our system, being close to the infinitely heavy impurity limit, allows for an extended time window of observability.

To this end, we are working - among other technical aspects - on optimising the sympathetic cooling scheme to obtain a spin-polarized Fermi gas of Li deep in the degenerate regime with a few thermal Cs impurities. We hope to be able to report first spectroscopy results (in frequency and time domain) in September.

References:

1. Efimov, V. Energy levels arising from resonant two-body forces in a three-body system. *Phys. Lett. B* 33, 563–564 (1970).
2. Tran, B. Rautenberg, M. et al. Fermions Meet Two Bosons—the Heteronuclear Efimov Effect Revisited. *Brazilian J. Phys.* 51, 316–322 (2021).
3. Pires, R. et al. Observation of Efimov Resonances in a Mixture with Extreme Mass Imbalance. *Phys. Rev. Lett.* 112, 250404 (2014).
4. Sun, M. & Cui, X. Efimov physics in the presence of a Fermi sea. *Phys. Rev. A* 99, 060701 (2019).
5. Enss, T., Tran, B. Rautenberg, M. et al. Scattering of two heavy Fermi polarons: Resonances and quasibound states. *Phys. Rev. A* 102, 063321 (2020).
6. Anderson, P. W. Infrared Catastrophe in Fermi Gases with Local Scattering Potentials. *Phys. Rev. Lett.* 18, 1049–1051 (1967).
7. Schmidt, R. et al. Universal many-body response of heavy impurities coupled to a Fermi sea: a review of recent progress. *Reports Prog. Phys.* 81, 024401 (2018).

Short bio (50 words) or link to website:

I studied physics at Heidelberg university with a focus on quantum many-body physics, first theoretically during my Bachelor's then also experimentally during my Master's degree. Currently I am working as PhD student in the group of Lauriane Chomaz and Matthias Weidemüller on mixtures of ultracold gases of Lithium and Caesium.

Relevant publications (optional):

1. Enss, T. et al. Scattering of two heavy Fermi polarons: Resonances and quasibound states. *Phys. Rev. A* 102, 063321 (2020). [<https://link.aps.org/doi/10.1103/PhysRevA.102.063321>]

2. Tran, B. et al. Fermions Meet Two Bosons—the Heteronuclear Efimov Effect Revisited. *Brazilian J. Phys.* 51, 316–322 (2021). [<http://dx.doi.org/10.1007/s13538-020-00811-5>]
3. Welz, K. et al. Anomalous loss behavior in a single-component Fermi gas close to a p-wave Feshbach resonance. *Phys. Rev. A* 107, 053310 (2023). [<https://link.aps.org/doi/10.1103/PhysRevA.107.053310>]

Career stage:

Student

Posters I / 29**Towards Fermi polarons with heavy impurities****Author:** Tobias Krom^{None}**Co-authors:** Michael Rautenberg ; Eleonora Lippi ; Lauriane Chomaz ; Matthias Weidemüller**Corresponding Author:** krom@physi.uni-heidelberg.de

The presented ultracold gases experiment uses bosonic ^{133}Cs and fermionic ^6Li . Because of their large mass ratio also qualitatively new observation become accessible. An example is the universal scaling law of Efimov states which has been investigated [1,2]. Currently, we are aiming for the creation of a Fermi polaron. In the infinitely heavy impurity limit the Fermi polaron loses its quasi-particle nature, which is known as the Anderson orthogonality catastrophe [3]. Since the chosen species are close to this limit, some precursors of this effect are predicted to arise [4,5]. For higher mass imbalance also the molecule-polaron crossover shifts to strong interactions which enriches the observable effects in this mixture [6,7,8].

In our apparatus a stable creation of a single species Fermi gas with $T/T_F \approx 0.2$ has been realized. We also set up a scheme with a sequential loading of the two species, which includes the movement of the Cs sample in a large optical dipole trap out of the center, to separate the preparation of the two components. Spectroscopic probes have been built up for ^{133}Cs (via a zero momentum Raman transition) as well as for ^6Li (using a radio frequency antenna). The combination of both spectroscopy methods allows for a direct distinction between molecules and polarons [9].

We will present the current status of the project at the time of the conference, which is expected to include a precise characterization of the sequential loading scheme as well as first spectroscopy signals of a degenerate gas with impurities.

References:

- [1] Pires, R.; Ulmanis, J.; Häfner, S.; Repp, M.; Arias, A.; Kuhnle, E. D.; Weidemüller, M. Observation of Efimov Resonances in a Mixture with Extreme Mass Imbalance. *Phys. Rev. Lett.* 2014, 112 (25), 250404.
- [2] Ulmanis, J.; Häfner, S.; Pires, R.; Kuhnle, E. D.; Wang, Y.; Greene, C. H.; Weidemüller, M. Heteronuclear Efimov Scenario with Positive Intraspecies Scattering Length. *Phys. Rev. Lett.* 2016, 117 (15), 153201.
- [3] Anderson, P. W. Infrared Catastrophe in Fermi Gases with Local Scattering Potentials. *Phys. Rev. Lett.* 1967, 18 (24), 1049–1051.
- [4] Schmidt, R.; Knap, M.; Ivanov, D. A.; You, J.-S.; Cetina, M.; Demler, E. Universal Many-Body Response of Heavy Impurities Coupled to a Fermi Sea: A Review of Recent Progress. *Rep. Prog. Phys.* 2018, 81 (2), 024401.
- [5] Adlong, H. S.; Liu, W. E.; Turner, L. D.; Parish, M. M.; Levinsen, J. Signatures of the Orthogonality Catastrophe in a Coherently Driven Impurity. *Phys. Rev. A* 2021, 104 (4), 043309.
- [6] Massignan, P. Polarons and Dressed Molecules near Narrow Feshbach Resonances. *EPL* 2012, 98 (1), 10012.
- [7] Trefzger, C.; Castin, Y. Impurity in a Fermi Sea on a Narrow Feshbach Resonance: A Variational Study of the Polaronic and Dimeronic Branches. *Phys. Rev. A* 2012, 85 (5), 053612.
- [8] Parish, M. M.; Adlong, H. S.; Liu, W. E.; Levinsen, J. Thermodynamic Signatures of the Polaron-Molecule Transition in a Fermi Gas. *Phys. Rev. A* 2021, 103 (2), 023312.

[9] Schirotzek, A.; Wu, C.-H.; Sommer, A.; Zwerlein, M. W. Observation of Fermi Polarons in a Tunable Fermi Liquid of Ultracold Atoms. *Phys. Rev. Lett.* 2009, 102 (23), 230402.

Short bio (50 words) or link to website:

- B. Sc. (2018) and M.Sc. (2021) in physics: Karlsruhe (Germany); Bachelor and Master thesis: optical fiber-based micro resonators for quantum networking (Prof. David Hunger)
- since 2021: PhD student in Heidelberg (Germany). Ultracold mixture experiment with Lithium and Cesium to explore polaron physics (Prof. Matthias Weidemüller/Prof. Lauriane Chomaz)

Relevant publications (optional):

Career stage:

Student

30

Photon BECs in dye-filled microcavities and VCSELs

Author: Axel Pelster¹

¹ *RPTU Kaiserslautern-Landau*

Corresponding Author: axel.pelster@physik.uni-kl.de

The talk provides an overview of current theoretical challenges for describing a photon Bose-Einstein condensate (BEC), which represents a modern prime example for an open dissipative quantum many-body system. In the original experimental platform of dye-filled microcavities [1] the technique of direct laser writing [2] allows to microstructure potentials with different geometries on the mirror surfaces. In this way soon lattices of coupled photon condensates containing hundreds of individual sites are realizable, which are expected to have spiral vortices [3]. We show that their shape can be approximately determined analytically with a projection optimization method, which extends the variational optimization method for BECs of closed systems to open-dissipative condensates [4]. Furthermore, quite recently photon BECs have also been observed in vertical cavity surface-emitting lasers (VCSELs) [5,6]. Here frequent photon absorption and emission processes occur due to the creation and annihilation of excitons in the semiconductor device, yielding a thermalization of photons. But it was found experimentally that the extracted spectral temperatures are significantly lower than those of the device, which warrants a theoretical explanation.

References:

- [1] J. Klaers, J. Schmitt, F. Vewinger and M. Weitz, *Nature* 468, 545 (2010).
 [2] J. Schulz, J. Noh, W. A. Benalcazar, G. Bahl, and G. von Freymann, *Nature Comm.* 13, 6597 (2022).
 [3] V. N. Gladilin and M. Wouters, *Phys. Rev. Lett.* 125, 215301 (2020).
 [4] J. Krauß, M.A.G. dos Santos Filho, F.E.A. dos Santos, and A. Pelster, arXiv:2311.10027 (2023).
 [5] R. C. Schofield, M. Fu, E. Clarke, I. Farrer, A. Trapalis, H. S. Dhar, R. Mukherjee, J. Heffernan, F. Mintert, R. A. Nyman, and R. F. Oulton, *Nature Photonics* (in press), arXiv:2306.15314 (2023).
 [6] M. Pieczarka, M. Gębski, A. N. Piasecka, J. A. Lott, A. Pelster, M. Wasiak, and T. Czyszanowski, *Nature Photonics* (in press), arXiv:2307.00081 (2023).

Short bio (50 words) or link to website:

<https://www-user.rhrk.uni-kl.de/~apelster>

Relevant publications (optional):

Career stage:

Postdoc

31

Universal dynamics in strongly interacting Bose gases far from equilibrium

Authors: Qi Liang¹; RuGway WU¹; Sebastian Erne¹; Jörg Schmiedmayer¹

¹ *Atominstitut, Technische Universität Wien*

Corresponding Author: qi.liang@tuwien.ac.at

We prepare gases of Feshbach molecules of lithium 6 far from equilibrium by introducing broad excitations using an optical speckle potential, and study the relaxation of the strongly interacting isolated many-body system going through the formation of a molecular Bose-Einstein condensates (mBEC). We report for the first time, universal spatio-temporal scaling behaviour in a strongly interacting Bose gas.

By varying the interaction strength of the system and observing the duration of scaling evolution, we show how the dimensionality of the system affects the stability of soliton-like excitations, and the non-equilibrium evolution rate. Comparison of the observed dependence on interaction with the earlier rubidium experiment suggests a possible general scaling law governing the self-similar evolution.

References:

Short bio (50 words) or link to website:

PhD at the Australian National University, group of Andrew Truscott, metastable helium experiment
Postdoc researcher at Atominstitut, Technische Universität Wien, Vienna, Austria, Schmiedmayer group, building and managing Lithium 6 quantum gas experiment.

Relevant publications (optional):

Career stage:

Postdoc

32

Vortices on rotating shell-shaped Bose-Einstein condensates

Author: Angela White¹

¹ *The University of Queensland*

Corresponding Author: a.white5@uq.edu.au

The recent realisation of hollow-core bubbles of Bose-Einstein condensates on the Cold Atom Lab aboard the International Space Station has rendered this intriguing geometry accessible, motivating study into the behaviour of vortices on curved surfaces [1]. In shell geometries, superfluid vortex behaviour promises interesting responses to rotation as the continuity of the velocity field required across the closed-curved surface of the shell imposes additional restrictions on the condensate phase. We study the response of a bubble condensate to an externally imposed rotation, demonstrating that for small rotation rates, a familiar triangular Abrikosov lattice of vortices is formed, with two aligned vortex lattices appearing in each hemispherical shell. An elliptical deformation of the spherically symmetric condensate shape occurs at larger rotation rates due to the centrifugal force. As the

driving rotation frequency is increased, a multi-charge vortex and its anti-vortex pair is formed at the poles, surrounded by singly charged vortices in the bulk condensate density. strong text

References:

[1] R.A. Carollo et al. Nature 606 (2022) 281-286.

Short bio (50 words) or link to website:

I am a Deborah Jin Research Fellow at the University of Queensland. Prior to this, I have held post-doctoral research positions at the Australian National University, Okinawa Institute of Science and Technology Graduate University, Japan and Newcastle University, UK.

Relevant publications (optional):

- [1] A. C. White, Triangular vortex lattices and giant vortices in rotating bubble Bose-Einstein condensates, Phys. Rev. A 109 (2024) 013301. <https://doi.org/10.1103/PhysRevA.109.013301>
 [2] T. Zhang et. al, Chaotic few-body vortex dynamics in rotating Bose-Einstein condensates, Phys. Rev. Fluids 4 (2019) 054701. <https://doi.org/10.1103/PhysRevFluids.4.054701>
 [3] A. White, Y. Zhang and T. Busch, Odd-petal-number states and persistent flows in spin-orbit-coupled Bose-Einstein condensates, Phys. Rev. A 95 (2017) 041604(R). <https://doi.org/10.1103/PhysRevA.95.041604>

Career stage:

Postdoc

33

Self organisation and metastability of cavity bosons at very long times, beyond the adiabatic elimination approximation

Author: Piotr Deuar¹

Co-authors: Giuliano Orso²; Jakub Zakrzewski³

¹ *Institute of Physics, Polish Academy of Sciences*

² *Université Paris Cité, Laboratoire Matériaux et Phénomènes Quantiques (MPQ), CNRS*

³ *Jagiellonian University in Kraków*

Corresponding Authors: jakub.zakrzewski@uj.edu.pl, giuliano.orso@u-paris.fr, deuar@ifpan.edu.pl

Phase-space formulations of quantum mechanics like the positive-P and truncated Wigner can give access to the full quantum behaviour of very large systems. In particular, the full distribution of single-shot configurations can be obtained from a stochastic simulation. This is particularly useful for dissipative systems for which direct simulation is harder but phase space methods become stable [1].

In a recent work [2] we have looked at the very long-time behaviour and self-organisation of weakly interacting bosons in a 2d optical lattice coupled to a lossy cavity, in the regime of high filling similar to experiments at ETH. The truncated Wigner representation allows us to go orders of magnitude longer in time compared to earlier numerical work. It takes into account the dynamics and correlation of the cavity mode, quantum fluctuations, and self-organization of individual runs. We observe metastability at very long times and superfluid quasi-long range order, in sharp contrast with the true long range order found in the ground state of the Bose-Hubbard model with extended interactions obtained by adiabatically eliminating the cavity. The metastability appears to be dependent on the relaxation of the adiabatic elimination constraint. As the strength of the cavity coupling increases in a superfluid, the system first becomes (lattice) supersolid at the superradiant transition and then turns into a charge-density wave via the BKT mechanism. Notably, experimental preparation times have often been comparable with the very long times simulated here, so the metastable effects may be relevant in practice.

References:

- [1] G. Orso, J. Zakrzewski, P. Deuar, Self-organized cavity bosons beyond the adiabatic elimination approximation, arXiv:2312.10502
- [2] P. Deuar, A. Ferrier, M. Matuszewski, G. Orso, M. H. Szymanska, Fully quantum scalable description of driven-dissipative lattice models, PRX Quantum 2, 010319 (2021)

Short bio (50 words) or link to website:

<http://info.ifpan.edu.pl/~deuar/people.html>

Relevant publications (optional):

G. Orso, J. Zakrzewski, P. Deuar,
Self-organized cavity bosons beyond the adiabatic elimination approximation, arXiv:2312.10502
<http://arxiv.org/abs/2312.10502>

W. Verstraelen, P. Deuar, M. Matuszewski, and T.C.H. Liew
Analog spin simulators: How to keep the amplitude homogeneous
Physical Review Applied 21, 024057 (2024)
<https://doi.org/10.1103/PhysRevApplied.21.024057>

P. Deuar, A. Ferrier, M. Matuszewski, G. Orso, M. H. Szymanska,
Fully quantum scalable description of driven-dissipative lattice models,
PRX Quantum 2, 010319 (2021)
<http://dx.doi.org/10.1103/PRXQuantum.2.010319>

Career stage:

Professor

34

Ferrodark solitons in a spinor superfluid: exact solutions, novel speed limit and anomalous dynamics

Author: Xiaoquan Yu¹

¹ *Graduate School of China Academy of Engineering Physics*

Corresponding Author: yuxq2008@gmail.com

Exact propagating topological solitons are found in the easy-plane phase of ferromagnetic spin-1 Bose-Einstein condensates, manifesting themselves as kinks in the transverse magnetization. Propagation is only possible when the symmetry-breaking longitudinal magnetic field is applied. Such solitons have two types: a low energy branch with positive inertial mass and a higher energy branch with negative inertial mass. Both types become identical at the maximum speed, a new speed bound that is different from speed limits set by the elementary excitations. The physical mass, which accounts for the number density dip, is negative for both types. In a finite one-dimensional system subject to a linear potential, the soliton undergoes oscillations caused by transitions between the two types occurring at the maximum speed.

References:

- 1) Xiaoquan Yu and P.B. Blakie, Dark soliton-like magnetic domain walls in a two-dimensional ferromagnetic superfluid, Phys. Rev. Research 3, 023043,(2021)
- 2) Xiaoquan Yu and P. B. Blakie, Propagating Ferrodark Solitons in a Superfluid: Exact Solutions and Anomalous Dynamics, Phys. Rev. Lett. 128, 125301,(2022)
- 3) Xiaoquan Yu and P. B. Blakie, Core structure of static ferrodark solitons in a spin-1 Bose-Einstein condensate, Phys. Rev. Research 4, 033056, (2022)

Short bio (50 words) or link to website:

<https://xiaoquanyu.github.io/resaerch-group-on-quantum-liquid/>

Relevant publications (optional):**Career stage:**

Professor

36

Hamiltonian engineering using Bragg matter-wave interferometers in an optical cavity

Author: Ana Rey¹

¹ *JILA*

Corresponding Author: arey@jilau1.colorado.edu

Cavity-QED systems have emerged as a powerful platform for generating highly entangled states, with significant implications for both quantum metrology and quantum simulation. A particular setting that has gained great interest for Bragg matterwave interferometry, and thus inertial navigation and fundamental science, is an array of free-falling atoms inside a cavity where one can encode a pseudo-spin $\frac{1}{2}$ degree of freedom using a pair of selected momentum states, which are coupled by the cavity photons.

I will show this system not only is suitable for the generation of a spin exchange Hamiltonians via two photon processes, but also arbitrary collective tunable Heisenberg XYZ models without the need of Floquet engineering. This can be done via cavity-mediated four-photon or higher photon interactions with two dressing lasers in a high finesse cavity. In particular I will report on how to dynamically generate the so-called two-axis counter-twisting model, a special type of collective XYZ model proposed more than 30 years ago for the fast generation of spin squeezed states that saturate the Heisenberg bound. In general I will discuss why this system opens a unique pathway for the use of momentum states for quantum enhanced interferometry and quantum simulation.

References:

Momentum-exchange interactions in a Bragg atom interferometer suppress Doppler dephasing
Luo C., H. Zhang, V.P.W. Koh, J.D. Wilson, A. Chu, M.J. Holland, A.M. Rey, and J.K. Thompson, *Science* 384, 551-556 (2024).

Hamiltonian Engineering of collective XYZ spin models in an optical cavity: From one-axis twisting to two-axis counter twisting models
Chengyi Luo, Haoqing Zhang, Anjun Chu, Chitose Maruko, Ana Maria Rey, James K. Thompson, arXiv:2402.19429

Short bio (50 words) or link to website:

<https://jila.colorado.edu/arey>

Relevant publications (optional):**Career stage:**

Professor

37

Universal coarsening in 2D and 3D Bose gases

Authors: Martin Gazo^{None}; Andrey Karailiev¹; Gevorg Martirosyan¹; Tanish Satoor¹; Christoph Eigen¹; Maciej Gałka²; Zoran Hadzibabic^{None}

¹ *University of Cambridge*

² *Heidelberg University*

Corresponding Authors: zh10001@cam.ac.uk, ak2026@cam.ac.uk, ce330@cam.ac.uk, galka@physi.uni-heidelberg.de, gm572@cam.ac.uk, mg816@cam.ac.uk, tss35@cam.ac.uk

Coarsening of an isolated far-from-equilibrium quantum system is a paradigmatic many-body phenomenon, relevant from subnuclear to cosmological lengthscales, and predicted to feature universal dynamic scaling. It is hypothesised that the associated scaling exponents would allow for the classification of nonequilibrium phenomena into an out-of-equilibrium analogue of equilibrium universality classes. In this talk, I will present our recent observations of universal scaling in the coarsening of isolated homogeneous two- and three-dimensional Bose gases.

We start by preparing a degenerate gas in a far-from-equilibrium state, and then observe the relaxation towards an equilibrium condensate. We reveal universal scaling in the experimentally accessible finite-time dynamics by elucidating and accounting for initial-state-dependent prescaling effects. The observed scaling exponents match analytical predictions, and are independent of both the initial state and the strength of interparticle interactions. The methods we introduce establish a direct comparison between cold-atom experiments and non-equilibrium field theory, and are applicable to any study of universality far from equilibrium.

Finally, we also investigate the timescales associated with coarsening. While stronger interactions generally speed up the thermalisation dynamics, we find that at sufficiently high interactions the coarsening dynamics becomes interaction-independent, hinting at the existence of a universal ‘speed limit’ for coarsening.

References:

Short bio (50 words) or link to website:

2016 –2020: University of Cambridge, MSci in Natural Sciences

2020 –now: University of Cambridge, PhD at Zoran Hadzibabic group

Relevant publications (optional):

1. Gazo, Martin, Andrey Karailiev, Tanish Satoor, Christoph Eigen, Maciej Gałka, and Zoran Hadzibabic. “Universal Coarsening in a Homogeneous Two-Dimensional Bose Gas.” arXiv preprint arXiv:2312.09248 (2023). <https://arxiv.org/abs/2312.09248>
2. Karailiev, Andrey, Martin Gazo, Maciej Gałka, Christoph Eigen, Tanish Satoor, and Zoran Hadzibabic. “Observation of an inverse turbulent-wave cascade in a driven quantum gas.” arXiv preprint arXiv:2405.01537 (2024). <https://arxiv.org/abs/2405.01537>
3. Gałka, Maciej, Panagiotis Christodoulou, Martin Gazo, Andrey Karailiev, Nishant Dogra, Julian Schmitt, and Zoran Hadzibabic. “Emergence of isotropy and dynamic scaling in 2D wave turbulence in a homogeneous Bose gas.” *Physical Review Letters* 129, no. 19 (2022): 190402. <https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.129.190402>

Career stage:

Student

Instabilities in Bose-Einstein condensed mixtures, by perturbing initially immiscible configurations

Author: Lauro Tomio^{None}

Corresponding Author: lauro.tomio@gmail.com

By considering initially immiscible configuration of homogeneous Bose-Einstein condensates confined in a two-dimensional circular box, I am reporting an investigation considering the emergence of three-kind of instabilities generated by different kind of time-dependent perturbations. By keeping the mixture in immiscible configurations, Rayleigh-Taylor (RT) and Kelvin-Helmholtz (KH) kind of instabilities can be generated. Another kind of instability in the mixture is verified by considering a sudden transition from the initially immiscible to miscible configuration, which can be obtained by reducing the inter-species interactions such that, along the following dynamics it will be kept smaller than the intra-species interactions. In this third case, it was also verified how the dynamics change by modifying the initial immiscible configurations of the mixture, by considering centrally and axially phase separated states before applying the sudden transition to miscible systems. For the binary mixture, it has been considered the rubidium isotopes ^{85}Rb and ^{87}Rb . In all the reported cases, it will be shown the behavior of the kinetic energy spectrum as a function of the wave number k , which follows approximated the $k^{-5/3}$ Kolmogorov behavior.

Note: Work in collaboration with R. Kishor Kumar, A. S. Bradley, S. Sabari, and A. Gammal

References:

Short bio (50 words) or link to website:

<http://lattes.cnpq.br/8621258845956348>

Relevant publications (optional):

Sabari, S.; Kumar, Kumar, R. K.; Tomio, Lauro. Vortex dynamics and turbulence in dipolar Bose-Einstein condensates. *Phys. Rev. A*109, 023313, 2024.

Nunes da Silva, Anacé ; Kumar, R. K.; Bradley, A. S.; Tomio, Lauro. Vortex generation in stirred binary Bose-Einstein condensates. *Phys. Rev. A*107, 03314, 2023.

Brito, L.; Tomio, Lauro; Gammal, A. Faraday waves on a bubble-trapped Bose-Einstein-condensed binary mixture. *Phys. Rev. A*108, 053315, 2023.

Career stage:

Professor

39

Bose-Einstein condensation and lasing of low-dimensional semiconductor materials

Author: Zheng Sun¹

¹ *East China Normal University*

Corresponding Author: zsun@lps.ecnu.edu.cn

Exciton-polaritons are unique quasiparticles formed by the interaction of excitons and optical modes, offering promising applications in coherent light sources and optical control devices. Our presentation focuses on the observation of Bose-Einstein condensation of upper polariton branch in a WS₂ monolayer microcavity. As the condensation threshold is reached, we note a nonlinear increase in upper polariton intensity, reduced linewidth, and enhanced temporal coherence, characteristic of the condensation phenomenon. Through simulations, we determine the specific particle density

range necessary for this condensation based on excitonic properties and pumping conditions. This discovery opens avenues for exploring condensate competition and its practical use in polaritonic lasers. Additionally, we explore the potential of Van der Waals homostructures consisting of stacked WS₂ layers for enhancing optical properties. Our experiments demonstrate ultra-low threshold laser emission from triple WS₂ layers separated by hBN, indicating efficient laser operation possibilities with such structures.

References:

Short bio (50 words) or link to website:

https://faculty.ecnu.edu.cn/_s29/sz2_en/main.psp

Relevant publications (optional):

Career stage:

Professor

40

Designing Atomtronic Circuits via Superfluid Dynamics

Author: Sarah Jährling^{None}

Co-authors: Ludwig Mathey ; Vijay Singh

Corresponding Authors: sjaehrli@physnet.uni-hamburg.de, lmathey@physnet.uni-hamburg.de, vsingh@physnet.uni-hamburg.de

We propose an implementation concept for atomtronic circuit elements based on the criticality of superfluid dynamics in specially designed Bose-Einstein condensates (BECs). Specifically, to obtain a logical 2-input AND-gate, we employ a T-shaped BEC together with two mobile and one stationary Gaussian barrier, functioning as Josephson junctions. The transistor-like behavior of the AND-gate can be identified by studying the resulting non-equilibrium density distributions around the stationary barrier for different scenarios of the deployable mobile barriers. Extending the original setup, we present a logical 4-input AND-gate in an attempt to realize an advanced connected atomtronic circuit. In addition, we discuss the possibility of a universal set of logical gates by establishing a connection to a logical NOT-gate motivated by the recent studies of Singh et al. [1] by exploiting Josephson oscillations. Lastly, we illustrate the potential by merging NOT- and AND-gate elements into an atomtronic NAND-gate.

References:

[1] V. P. Singh, N. Luick, L. Sobirey, and L. Mathey, Josephson junction dynamics in a two-dimensional ultracold bose gas, *Phys. Rev. Res.* 2, 033298 (2020).

Short bio (50 words) or link to website:

I completed my Bachelor of Science (Physics) with the thesis ‘Latent Symmetry Analysis of Small & Medium-Sized Graphs’ (2022), under the supervision of Prof. Dr. Peter Schmelcher (<https://www.physik.uni-hamburg.de/en/iqp/schmelcher.html>). I am currently approaching the end of my Master’s degree, under supervision of Prof. Dr. Ludwig Mathey (<https://www.physik.uni-hamburg.de/en/iqp/mathey.html>).

Relevant publications (optional):

Career stage:

Student

Posters I / 41

Narrow-linewidth exciton-polariton laser

Authors: Andrew Truscott¹; Bianca Rae Fabricante^{None}; David Snoke²; Elena Ostrovskaya¹; Eliezer Estrecho¹; Kenneth West³; Loren Pfeiffer³; Maciej Pieczarka⁴; Mark Steger²; Mateusz Król⁵; Matthias Wurdack⁶

¹ *Department of Quantum Science and Technology, Research School of Physics, The Australian National University, Australia*

² *Department of Physics and Astronomy, University of Pittsburgh, USA*

³ *Department of Electrical Engineering, Princeton University, USA*

⁴ *Department of Experimental Physics, Faculty of Fundamental Problems of Technology, Wrocław University of Science and Technology, Wrocław, Poland*

⁵ *Australian National University*

⁶ *The Australian National University*

Corresponding Authors: elena.ostrovskaya@anu.edu.au, bianca.fabricante@anu.edu.au, eliezer.estrecho@anu.edu.au, mwurdack@stanford.edu, mateusz.krol@anu.edu.au

Exciton-polariton condensates are non-equilibrium quantum fluids formed by short-lived hybrid light-matter particles in a semiconductor microcavity. In the steady-state regime, these particles decay via photon emission that inherits the coherence properties of the condensate. This so-called exciton-polariton laser is a promising source of coherent light for low-energy applications due to its low-threshold operation. However, a detailed experimental study of its spectral purity, which directly affects its coherence properties, is still missing. Here, we present a high-resolution spectroscopic investigation of the energy and linewidth of an exciton-polariton laser in the single-mode regime, which derives its coherent emission from an optically pumped exciton-polariton condensate. We report an ultra-narrow linewidth of 56 MHz or 0.24 μeV , the narrowest on record [1], corresponding to a coherence time of 5.7 ns. The narrow linewidth is achieved by using an exciton-polariton condensate with a high photonic content confined in an optically induced trap that minimizes an overlap between the condensate and the excitonic reservoir [2]. Contrary to previous reports [2,3], we observe that the excitonic reservoir injected by the pump and responsible for creating the trap does not strongly affect the emission linewidth, as long as the condensate is trapped and the pump power is well above the condensation (lasing) threshold. The long coherence time of the exciton-polariton system uncovered here opens opportunities for manipulating its macroscopic quantum state, which is essential for applications in classical and quantum computing.

References:

- [1] B.R. Fabricante, M. Król, M. Wurdack, M. Pieczarka, M. Steger, D. W. Snoke, K. West, L. N. Pfeiffer, A. G. Truscott, E. A. Ostrovskaya, and E. Estrecho, arXiv:2404.18354 (2024).
- [2] A. Askitopoulos, L. Pickup, S. Alyatkin, A. Zasedatelev, K.G. Lagoudakis, W. Langbein, and P. G. Lagoudakis. Giant increase of temporal coherence in optically trapped polariton condensate. arXiv preprint arXiv:1911.08981 (2019).
- [3] A. P. D. Love, D. N. Krizhanovskii, D. M. Whittaker, R. Bouchekioua, D. Sanvitto, S. Al Rizeiqi, R. Bradley, M. S. Skolnick, P. R. Eastham, R. Andre, and Le Si Dang. Intrinsic decoherence mechanisms in the micro-cavity polariton condensate. Phys. Rev. Lett., 101(1):067404 (2008).

Short bio (50 words) or link to website:

I am a PhD Physics student in the Polariton BEC Group of the Department of Quantum Science and Technology, Research School of Physics, College of Science, The Australian National University.

Relevant publications (optional):**Career stage:**

Student

Posters II / 42

Work, heat and entropy in isolated quantum systems

Author: Lewis Williamson^{None}

Corresponding Author: lewis.williamson@uq.edu.au

Work, heat and entropy are three of the most fundamental concepts in thermodynamics. Over the past 30 years, the discovery of fluctuation theorems in both classical and quantum systems have extended these concepts from equilibrium (slow) to non-equilibrium (fast) processes. To date, almost all this exploration has defined thermal equilibrium in terms of the canonical thermal distribution. Coincident with this progress, our understanding of thermal states at the microscopic level has evolved substantially, with profoundly new insights provided by the Eigenstate Thermalization Hypothesis. This new theory describes how a pure quantum state may look thermal, despite the absence of chaos-inducing non-linear dynamics. Aside from a few isolated studies, there is a notable absence of research into fluctuation theorems and notions of heat and work from the perspective of the Eigenstate Thermalization Hypothesis.

Here we explore the concepts of heat, work and entropy in a quantum spin-chain undergoing unitary evolution starting from a pure state. This system can conveniently be tuned from integrable to non-integrable by changing the combination of external fields incident on the system. We define notions of heat, work and entropy in this system and explore their dependence on the rate of work extraction and the integrability of the system. Our results provide new connections between the Eigenstate Thermalization Hypothesis and thermodynamic fluctuation theorems, with broad relevance for finite temperature quantum systems.

References:

Short bio (50 words) or link to website:

From 2020-present I have done research in quantum thermodynamics as a Postdoctoral Research Fellow at the University of Queensland. Prior to this I did my PhD at the University of Otago researching spinor Bose-Einstein condensates and then worked as a Research Associate at Lancaster University researching strongly-correlated light-matter systems.

Relevant publications (optional):

Career stage:

Postdoc

Posters I / 43

Realising topological phases in the spin-1/2 quantum kicked rotor

Authors: Anushka Thenuwara¹; Andrew Groszek²; Matthew Edmonds³; Matthew Davis^{None}; Kristian Helmer-son¹

¹ Monash University

² The University of Queensland

³ University of Queensland

Corresponding Authors: m.edmonds@uq.edu.au, anushka.thenuwara@monash.edu, kristian.helmerson@monash.edu, a.groszek@uq.edu.au, mdavis@uq.edu.au

The quantum kicked rotor (QKR) is an archetypal system in the study of quantum chaos, and can be realised by periodically delta-kicking a cloud of ultracold atoms. This system is mathematically equivalent to a tight-binding model - up to an exchange of position and momentum space - and therefore exhibits behaviour analogous to electrons evolving in a lattice. Early work focused on

“dynamical localisation” in the QKR, a phenomenon equivalent to Anderson localisation, but in momentum space rather than position space. More recently, interest has arisen in realising a QKR with a spin-1/2 degree of freedom. With the appropriate form of kicking, this system supports topological phases, analogous to a topological insulator. We propose an experiment to realise this setup using ultracold atoms, allowing for the detection of these topological phases via their associated topological invariants.

References:

Short bio (50 words) or link to website:

<https://smp.uq.edu.au/profile/9399/andrew-groszek>

Relevant publications (optional):

Career stage:

Postdoc

Posters II / 44

Equatorial waves in rotating bubble-trapped superfluids

Author: Dmitry Efimkin¹

Co-author: Guangyao Li¹

¹ *Monash University*

Corresponding Author: dmitry.efimkin@monash.edu

As the Earth rotates, the Coriolis force causes various oceanic and atmospheric waves to be trapped along the equator, including Kelvin, Yanai, Rossby, and Poincaré modes. It has been demonstrated that the mathematical origin of these waves is related to the nontrivial topology of the underlying hydrodynamic equations. Inspired by recent observations of Bose-Einstein condensation (BEC) in bubble-shaped traps in microgravity ultracold quantum gas experiments, we demonstrate that equatorial modes are supported by a rapidly rotating condensate in a spherical geometry. Using a zero-temperature coarse-grained hydrodynamic framework, we reformulate the coupled oscillations of the superfluid and the Abrikosov vortex lattice resulting from rotation as a Schrödingerlike eigenvalue problem. The resulting non-Hermitian Hamiltonian is topologically nontrivial. We also solve the hydrodynamic equations for a spherical geometry and find that the rotating superfluid hosts Kelvin, Yanai, and Poincaré equatorial modes, but not the Rossby mode. Our predictions can be tested with state-of-the-art bubble-shaped trapped BEC experiments.

References:

G. Li and D. K. Efimkin - Phys. Rev. A 107, 023319 (2023)

Short bio (50 words) or link to website:

<https://sites.google.com/a/monash.edu/dmitry-efimkin-research-group/publications?authuser=0>

Relevant publications (optional):

1. Equatorial Waves in Rotating Bubble-Trapped Superfluids - G. Li and D. K. Efimkin - Phys. Rev. A 107, 023319 (2023)
2. Topological hybrid electron-hole Cooper pairing - A. Chansky and D.K. Efimkin - Phys. Rev. B 108, 075433 (2023)
3. Anomalous drag in electron-hole condensates with granulated order - H. Liu, A.H. MacDonald, and D.K. Efimkin - Phys. Rev. Lett. 127, 166801 (2021)

Career stage:

Professor

Posters I / 45**Shear-Induced Decaying Turbulence in Bose-Einstein Condensates****Authors:** Guillaume Gauthier¹; Halina Rubinsztein-Dunlop^{None}; Simeon Simjanovski¹; Tyler Neely²¹ *The University of Queensland*² *University of Queensland***Corresponding Authors:** hrubinszteindunlop@gmail.com, g.gauthier@uq.edu.au, s.simjanovski@uq.net.au, t.neely@uq.edu.au

In this talk, we experimentally consider the problem of decaying turbulence in a Bose-Einstein condensate (BEC) superfluid. We begin with a shear layer comprised of quantum vortices formed between a stationary BEC and a stirred-in persistent current. This structure breaks down rapidly (<150 ms) through vortex pairing which we characterise through simple crystal structure analysis [1,2]. Subsequently decaying turbulence is established, through the progressive clustering of the vortices [3], which follows a power law decay with time, similar to decaying turbulence in other two-dimensional systems under the classical Kelvin-Helmholtz instability (KHI) [4,5]. We extend this investigation using a point-vortex model that matches experimental conditions [6]. From this, we observe a convergence of the power-law exponent to a fixed value.

[1] H. Aref, On the equilibrium and stability of a row of point vortices, *Journal of Fluid Mechanics* 290, 167–181 (1995).

[2] D. Hernández-Rajkov et al., Connecting shear flow and vortex array instabilities in annular atomic superfluids, *Nature Physics*. (2024)

[3] A. W. Baggaley and N. G. Parker, Kelvin-Helmholtz instability in a single-component atomic superfluid, *Physical Review A* 97, 053608 (2018).

[4] D. A. Schecter, D. H. E. Dubin, K. S. Fine, and C. F. Driscoll, Vortex crystals from 2D Euler flow: Experiment and simulation, *Physics of Fluids* 11, 905 (1999).

[5] Y. Pomeau, Vortex dynamics in perfect fluids, *Journal of Plasma Physics* 56, 407–418 (1996)

[6] M. T. Reeves et al., Turbulent Relaxation to Equilibrium in a Two-Dimensional Quantum Vortex Gas, *Physical Review X* 12, 011031 (2022)

References:**Short bio (50 words) or link to website:**

I am a PhD student at the University of Queensland, Bose-Einstein condensate laboratory. My research is primarily related to 2D quantum turbulence and quantum vorticity.

Relevant publications (optional):

S. Simjanovski, G. Gauthier, M. J. Davis, H. Rubinsztein-Dunlop, and T. W. Neely, Optimizing persistent currents in a ring-shaped Bose-Einstein condensate using machine learning, *Phys. Rev. A* 108, 063306 (2023). <https://journals.aps.org/pr/abstract/10.1103/PhysRevA.108.063306>

Career stage:

Student

Posters II / 46

Exploring the dynamics of polar core vortices in homogeneous spin-1 Bose-Einstein condensates

Authors: Matthew Davis^{None}; Matthew Edmonds¹

¹ *University of Queensland*

Corresponding Authors: mdavis@uq.edu.au, m.edmonds@uq.edu.au

Atomic superfluids with internal spin degrees of freedom exhibit a rich phenomenology, which in turn heralds their utility in both fundamental research and applications with emerging quantum technologies such as metrology and atomtronics. Complementary to this, it is now feasible to make quantum gases in homogeneous potentials, allowing a stronger connection between existing theoretical methodology and state-of-the-art experiments [1].

In this work, we explore the topological nature of the superfluid vortices present in the ferromagnetic phase of spin-1 Bose-Einstein condensates [2, 3]. In particular we examine the static and dynamic properties of polar core vortices that exist in the so-called easy-plane phase of the spinor system. Comprehensive numerical simulations reveal the structure of the individual vortices, while the dynamics of pairs of these excitations are shown to depend strongly on their individual phase windings as well as the atomic interactions and confining geometry of the homogeneous system. Our findings provide both useful insight as well as being accessible to the current generation of experiments with spinor condensate systems [4].

References:

- [1] G. Gauthier, I. Lenton, N. M. Parry, M. Baker, M. J. Davis, H. R.-Dunlop, T. W. Neely, *Optica* 3, 1136 (2016).
- [2] H. Saito, Y. Kawaguchi, and M. Ueda *Phys. Rev. Lett.* 96, 065302 (2006).
- [3] L. A. Williamson and P. B. Blakie, *Phys. Rev. A* 94, 063615 (2016).
- [4] Y. Xiao, M. O. Borgh, L. S. Weiss, A. A. Blinova, J. Rustekoski, and D. S. Hall, *Commun. Phys.* 4, 52 (2021).

Short bio (50 words) or link to website:

<https://smp.uq.edu.au/profile/13035/matthew-edmonds>

Relevant publications (optional):

Career stage:

Postdoc

Posters II / 47

Vortex matter simulation of the one component plasma

Authors: Charles Glasspool¹; Guillaume Gauthier²; Matthew Davis^{None}; Matthew Reeves¹; Tyler Neely¹

¹ *University of Queensland*

² *The University of Queensland*

Corresponding Authors: mdavis@uq.edu.au, g.gauthier@uq.edu.au, m.reeves@uq.edu.au, t.neely@uq.edu.au

In this work, we explore the low-energy states of vortex matter in a quasi-2D uniform BEC superfluid [1,2]. Mapping this system to 2D charges, we realize a vortex matter simulator of the one-component plasma (OCP) a fundamental minimal model in condensed matter. While the OCP is broadly considered a toy model, our system realizes its equilibrium states exactly. To benchmark our simulator, starting from the minimum energy state, a Wigner crystal, and we observe the melting of the lattice under systematic heating. We observe several predicted features of melting transition, including excess density at the edge of the vortex cluster, spatial squeezing of the density distribution, and persistent crystallization at the cluster edge [3]. These states of vortex matter have gained prominence in the theory of the fractional quantum Hall effect, where the 2D electron gas moves analogously to vortices in an incompressible fluid, and the vortex density maps to the density of the quantum Hall droplet.

References:

Short bio (50 words) or link to website:

uq-bec.org

Relevant publications (optional):

Career stage:

Professor

Posters II / 48

On Demand Formation of Topological Defects in Ferromagnetic Spinor Bose Einstein Condensates

Authors: Guillaume Gauthier¹; Halina Rubinsztein-Dunlop^{None}; Matthew Davis^{None}; Tyler Neely²; Zac Kerr¹

¹ *The University of Queensland*

² *University of Queensland*

Corresponding Authors: z.kerr@uqconnect.edu.au, g.gauthier@uq.edu.au, mdavis@uq.edu.au, t.neely@uq.edu.au, hrubinszteindunlop@gmail.com

Spinor Bose-Einstein condensates (sBECs) are quantum superfluids with a spin degree of freedom arising from interactions between atoms in different magnetic sublevels $m_F = +1, 0, -1$. These novel ultracold atomic systems can exhibit ferromagnetic order and offer enhanced opportunities for exploring phenomena beyond those accessible in scalar BEC's, such as new classes of topological defects. The polar core vortex (PCV) is an example of a unique defect occurring in a transversely magnetized, ferromagnetic sBECs. PCVs exhibits opposing spin circulation in the $m_F = \pm 1$ components and an unmagnetized vortex core populated by atoms in the $m_F = 0$ state. This results in a defect with a topologically protected winding of the transverse magnetization and a flat density profile. The first experimental observation of a PCV was achieved by Sadler et al [1] in 2006, where the vortex sporadically formed following a magnetic field quench. Due to the non-deterministic nature of creating PCV's, further experimental study of their properties and dynamics has been limited, thus leaving a wide range of PCV applications left to be explored. In this presentation, I report on the apparatus developed to realize the first on-demand creation of PCVs in a uniform 2D ^{87}Rb sBEC. We also demonstrate our fine experimental control of density and spin profiles via the use of DMDs that lead our investigations into PCV dynamics, PCV driven turbulence and tests of ultracold spintronic devices

References:

[1] L. E. Sadler, J. M. Higbie, S Leslie, M. Vengalattore, D. Stamper-Kurn, Spontaneous symmetry breaking in a quenched ferromagnetic spinor Bose-Einstein condensate, *Nature* 443 (2006) 312.

Short bio (50 words) or link to website:

<https://www.uq-bec.org/zachary-kerr>

Relevant publications (optional):

Career stage:

Student

Posters II / 49

Phononic crystal trapping geometries for improved rotational sensing with ultracold atoms

Author: Lachlan Miller^{None}

Co-authors: Charles Woffinden ; Halina Rubinsztein-Dunlop ; Tyler Neely ¹

¹ *University of Queensland*

Corresponding Authors: h rubinszteindunlop@gmail.com, s4587645@student.uq.edu.au, t.neely@uq.edu.au

Ultracold atom interferometry for inertial sensing in GPS/GNSS-denied environments presents a compact and more sensitive alternative to traditional methods such as light interferometry. While free-space interferometers have approached commercial scale implementation, trapping the atomic sample throughout the measurement may offer greater robustness to accelerations of the apparatus, although it is potentially limited by increased phase-diffusion due to two-body interactions at high densities.

Marti et al. [1] developed a sensor that measured rotation using the interference of counterpropagating standing-wave phonons in a ring geometry. Building on this work, we recently explored the sensing limits of this trapped atom system, experimentally finding an improved sensitivity of 0.3 rad s^{-1} . Numerical modelling found that higher harmonic generation by phonon mode mixing and thermal damping limit the lifetime of the imprinted phonons and the Q value [2].

To address this issue, we explore the implementation of a resonant phononic crystal geometry, extending our findings with atomtronic Helmholtz resonators [3]. We simulate the evolution of the system in this new geometry with the Gross-Pitaevski Equation, investigating the role of geometry on the lifetime of imprinted phonons and hence the theoretical Q value. Experimentally, confining the atoms to novel 2D geometries and limiting the thermal background may improve the sensitivity of the rotational sensor.

References:

1. G. E. Marti, R. Olf and D. M. Stamper-Kurn, Collective excitation interferometry with a toroidal Bose-Einstein condensate, *Phys. Rev. A* 91, 013602 (2015), doi:10.1103/PhysRevA.91.013602.
2. C. W. Woffinden, A. J. Groszek, G. Gauthier, B. J. Mommers, M. W. J. Bromley, S. A. Haine, H. Rubinsztein-Dunlop, M. J. Davis, T. W. Neely and M. Baker, Viability of rotation sensing using phonon interferometry in Bose-Einstein condensates, *SciPost Phys.* 15, 128 (2023), doi: 10.21468/SciPostPhys.15.4.128
3. G. Gauthier, S. S. Szigeti, M. T. Reeves, M. Baker, T. A. Bell, H. Rubinsztein-Dunlop, M. J. Davis and T. W. Neely, Quantitative Acoustic Modes for Superfluid Circuits, *Phys. Rev. Lett.* 123, 260402 (2019), doi: 10.1103/PhysRevLett.123.260402

Short bio (50 words) or link to website:

<https://www.uq-bec.org/lachlan-miller-1>

Relevant publications (optional):**Career stage:**

Student

Posters II / 50**Tensor network methods for the Gross-Pitaevskii equation**

Author: Ryan Connor¹

Co-authors: Callum Duncan¹; Andrew Daley²

¹ *University of Strathclyde*

² *University of Oxford, University of Strathclyde*

Corresponding Author: ryan.connor.2019@uni.strath.ac.uk

Numerically simulating partial differential equations can be a challenging task. Often one requires huge simulation grids to be able to correctly resolve all physical length scales, leading to huge memory and CPU time requirements. Recently, there has been a focus in extending the applications of Tensor Networks (TNs) into simulations of challenging non-linear partial differential equations [1,2,3]. TNs have been widely successful, in the study of quantum many-body physics and strongly correlated systems [4], providing a framework to obtain physically motivated data compression. In this work, we extend the application of TNs to simulate quantum fluids and turbulence through the Gross-Pitaevskii Equation (GPE). We introduce a procedure to implement the split-step Fourier method for time evolution [5], and use this to demonstrate vortex formation in the GPE and dipolar condensates. We show that by encoding our problem in a TN format, one can perform simulations on large spatial grids in 2D and 3D, which would be unfeasible with standard direct numerical simulations.

References:

- [1] - N.Gourianov et.al, Nat Comput Sci 2, 30–37 (2022), <https://doi.org/10.1038/s43588-021-00181-1>
- [2] - E. Ye and N. F. G. Loureiro, Physical Review E 106, 10.1103, <https://link.aps.org/doi/10.1103/PhysRevE.106.035208>
- [3] - M.Kiffner and D.Jaksch, Preprint, arXiv:2303.03010, <https://doi.org/10.48550/arXiv.2303.03010>
- [4] - U.Schollwoeck, Annals of Physics 326, 96 (2011), <https://doi.org/10.1016/j.aop.2010.09.012>
- [5] - M.D Feit et.al, Journal of Computational Physics, Volume 47, Issue 3 (1982), [https://doi.org/10.1016/0021-9991\(82\)90091-2](https://doi.org/10.1016/0021-9991(82)90091-2)

Short bio (50 words) or link to website:

I am a second year PhD student at the University of Strathclyde, within a quantum many body physics group led by Prof Andrew Daley. Within this group, I am focused on quantum inspired methods and quantum algorithms for solving partial differential equations and strongly correlated systems.

Relevant publications (optional):**Career stage:**

Student

51

False vacuum decay in an ultracold spin-1 Bose gas

Author: Thomas Billam^{None}

Corresponding Author: thomas.billam@newcastle.ac.uk

Cold atomic gases offer multiple prospects for simulating the physics of the very early universe in the laboratory, and an ultracold atom analogue of early universe false vacuum decay was recently observed in a ferromagnetic superfluid [1]. In this talk I will discuss theoretical modelling of false vacuum decay analogues using c-field methods [2]. Specifically, I will describe truncated-Wigner and stochastic projected Gross-Pitaevskii simulations of false vacuum decay, modelling zero and finite temperatures respectively, in a spin-1 Bose gas analogue [3, 4]. I discuss the comparison of these simulations to the bubble nucleation rates predicted by the non-perturbative instanton method, and areas of potential future work to refine stochastic simulations of these non-equilibrium superfluid systems.

References:

- [1] Observation of false vacuum decay via bubble formation in ferromagnetic superfluids, A. Zenesini, A. Berti, R. Cominotti, C. Rogora, I. G. Moss, T. P. Billam, I. Carusotto, G. Lamporesi, A. Recati, and G. Ferrari, *Nature Physics* 20, 558 (2024).
- [2] P.B. Blakie, A.S. Bradley, M.J. Davis, R.J. Ballagh, and C.W. Gardiner, Dynamics and statistical mechanics of ultra-cold Bose gases using c-field techniques, *Advances in Physics* 57:5, 363 (2008).
- [3] False-vacuum decay in an ultracold spin-1 Bose gas, T.P. Billam, K. Brown, and I.G. Moss, *Physical Review A* 105, L041301 (2022).
- [4] Bubble nucleation in a cold spin 1 gas, T.P. Billam, K. Brown, and I.G. Moss, *New Journal of Physics* 25, 043028 (2023).

Short bio (50 words) or link to website:

<https://www.ncl.ac.uk/math-physics/people/profile/thomasbillam.html>

Relevant publications (optional):

Career stage:

Professor

52

An Efficient Quantum Phase-Space Method for Simulating Feedback Control of Interacting Many-Body Quantum Systems

Author: Simon Haine^{None}

Co-authors: Joe Hope¹; Kaiwen Zhu¹; Zain Mehdi¹

¹ *Australian National University*

Corresponding Author: simon.haine@anu.edu.au

Accurately modelling measurement and control of ultracold Bose gases has so far proved unfeasible due the prohibitively large size of the numeric simulations, and problems with under-sampling. We present a new field-theoretic technique based on existing phase-space methods, and use it model feedback cooling of a Bose gas subject to measurements via phase-contrast imaging.

is developed for scalable numerical simulations of controlled quantum systems, and used to model feedback cooling of a Bose gas subject to periodic, non-destructive measurements via phase-contrast imaging. We check the validity of our approach in a two-mode system, which permits an exact solution due to its low-dimensional nature, and observe exceptional agreement across various moments of pseudospin operators. In addition, we benchmark our approach with existing techniques such as the Number-Phase Wigner particle filter, which has been the leading choice for existing simulations of controlled quantum systems.

Finally, we present preliminary results demonstrating successful cooling of a thermal state with low condensate fraction to condensate formation in both quasi-1D and 2D geometries, correctly accounting for measurement induced backaction and spontaneous emission effects. It is shown that the final achievable condensate fraction is dependent upon experimental parameters such as the measurement strength, rate, and detector resolution, and a simple model is constructed to derive optimal values for the parameters above.

References:

Short bio (50 words) or link to website:

- Completed PhD at the Australian National University
- Moved to University of Queensland where I undertook ARC APD and DECRA fellowships
- Moved to University of Sussex UK for a Marie Curie fellowship
- Currently at ANU on an ARC Future Fellowship

Relevant publications (optional):

Career stage:

Postdoc

Posters II / 53

Nonequilibrium Transport in a Superfluid Josephson Junction Chain: Is There Negative Differential Conductivity?

Authors: Samuel Begg¹; Matthew Davis^{None}; Matt Reeves²

¹ *Oklahoma State University*

² *University of Queensland*

Corresponding Authors: mdavis@uq.edu.au, samuel.ewan.begg@gmail.com, m.reeves@uq.edu.au

We consider the far-from-equilibrium quantum transport dynamics in a 1D Josephson junction chain of multi-mode Bose-Einstein condensates. We develop a theoretical model to examine the experiment of R. Labouvie et al. [Phys. Rev. Lett. 115, 050601 (2015)], wherein the phenomenon of negative differential conductivity (NDC) was reported in the refilling dynamics of an initially depleted site within the chain. We demonstrate that a unitary c -field description can quantitatively reproduce the experimental results over the full range of tunnel couplings, and requires no fitted parameters. With a view towards atomtronic implementations, we further demonstrate that the filling is strongly dependent on spatial phase variations stemming from quantum fluctuations. Our findings suggest that the interpretation of the device in terms of NDC is invalid outside of the weak coupling regime. Within this restricted regime, the device exhibits a hybrid behaviour of NDC and the AC Josephson effect. I will also discuss open questions and future directions.

References:

Begg, S. E., Davis, M. J., Reeves, M. T., Phys. Rev. Lett. 132, 103402 (2024)

Short bio (50 words) or link to website:

Undergraduate at University of Western Australia, before a PhD at King's College London with Joe Bhaseen. Then a post-doc at the Asia Pacific Center for Theoretical Physics in South Korea with Ryo Hanai and a post-doc at Oklahoma State University with Thomas Bilitewski.

Relevant publications (optional):

PRL 132, 103402 (2024) <https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.132.103402>

Other works on non-equilibrium systems (not superfluids):

PRL 132, 120401 (2024) <https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.132.120401>

PRR 5, 043288

<https://journals.aps.org/prresearch/abstract/10.1103/PhysRevResearch.5.043288>

Career stage:

Postdoc

Posters II / 54

Non-Hermitian dynamics of a photonic wave packet in a micro-cavity filled with a liquid crystal

Author: Mateusz Król¹

Co-authors: Y.-M. Robin Hu¹; Rafał Mazur²; Przemysław Morawiak²; Wiktor Piecek²; Przemysław Kula²; Andrew G. Truscott¹; Barbara Piętka³; Jacek Szczytko³; Eliezer Estrecho¹; Elena A. Ostrovskaya¹

¹ *The Australian National University*

² *Military University of Technology, Poland*

³ *University of Warsaw*

Corresponding Author: mateusz.krol@anu.edu.au

Planar optical microcavities provide an excellent platform for studying the evolution of two-dimensional (2D) photonic wave packets in the presence of synthetic fields. Numerous experimental methods of probing the light that escapes the cavity, including its phase and polarization, enable direct comparison between the measurements and theoretical predictions. Recently, theoretical description of wave packet evolution in optical microcavities was extended to take into account their inherent open-dissipative nature. Such theoretical investigations use a non-Hermitian Hamiltonian to describe the system and predict novel, up to now unobserved, effects arising directly from the non-Hermiticity like self-acceleration [1] and corrections to the anomalous Hall drift [2].

In this work, we focus on experimental verification of these predictions using a planar microcavity filled with a birefringent liquid crystal (LC). Optical anisotropy of the LC results in separate cavity modes with specific polarizations and differing decay rates. In addition, the sensitivity of the LC to the external electric fields brings unique possibility to tune the cavity modes by voltage applied to the transparent electrodes built-in into the cavity. All these properties make this system uniquely suitable for investigation of non-Hermitian effects on the propagation of photonic wave packets inside the cavity. Specifically, we measure the centre-of-mass position as well as the pseudospin of a laser pulse injected and propagating inside a cavity and observe clear deviations from Hermitian dynamics, such as nonharmonic pseudospin oscillations leading to nonharmonic Zitterbewegung effect.

References:

[1] Y.-M. Robin Hu et al., Phys. Rev. B 108, 115404 (2023).

[2] Y.-M. Robin Hu et al., Opt. Mater. Express 14, 664-686 (2024).

Short bio (50 words) or link to website:

Mateusz Król is a post-doc in Polariton BEC group at Australian National University. He finished his PhD in 2022 at University of Warsaw, where he worked on exciton-polaritons and microcavities filled with liquid crystals.

Relevant publications (optional):**Career stage:**

Postdoc

55

Shapiro steps in driven atomic Josephson junctions

Author: Vijay Singh^{None}

Corresponding Author: vijay24phy@gmail.com

We study driven atomic Josephson junctions realized by coupling two two-dimensional atomic clouds with a tunneling barrier. By moving the barrier at a constant velocity, dc and ac Josephson regimes are characterized by a zero and nonzero atomic density difference across the junction, respectively. Here, we monitor the dynamics resulting in the system when, in addition to the above constant velocity protocol, the position of the barrier is periodically driven. We demonstrate that the time-averaged particle imbalance features a plateau behavior that is the analog of Shapiro steps observed in driven superconducting Josephson junctions. The underlying dynamics reveals an intriguing interplay of the vortex and phonon excitations, where Shapiro steps are induced via suppression of vortex growth. We study the system with a classical-field dynamics method, and benchmark our findings with a driven circuit dynamics. We report on the LENS and Kaiserslautern experiments that realize Shapiro steps in driven ultracold atom Josephson junctions in close collaboration.

References:

V. P. Singh, J. Polo, L. Mathey, and L. Amico. arXiv:2307.08743 (2023)
[<https://arxiv.org/abs/2307.08743>]

Short bio (50 words) or link to website:

I am a senior researcher at Quantum Research Center, Technology Innovation Institute, Abu Dhabi, focussing on the development of Atomtronic circuits. My PhD work was on superfluid properties of ultracold atoms.

Relevant publications (optional):

1. S. Sunami, V. P. Singh, et al., Science 382, 443–447 (2023) [<https://www.science.org/doi/abs/10.1126/science.abq6753>].
2. H. P. Zahn, V. P. Singh, et al., Phys. Rev. X 12, 021014 (2022) [<https://journals.aps.org/prx/abstract/10.1103/PhysRevX.12.021014>].
3. V. P. Singh et al., Phys. Rev. Research 2, 033298 (2020) [<https://journals.aps.org/prresearch/abstract/10.1103/PhysRevResearch.2.033298>].

Career stage:

Postdoc

Incoherent Transport in the detection of Majorana bound states

Author: Oscar Eastman¹

¹ *University of Utrecht/QuMat*

Corresponding Author: o.a.eastman@uu.nl

Despite their existence having been proposed in 1937, Majorana fermion's status as either fundamental particles or emergent Majorana bound states (MBS) remains unproven experimentally. Due to its analytic tractability as a theoretical model, the one-dimensional Kitaev model has been at the forefront of this search. A key feature of a MBS is the presence of a zero-bias peak in the conductivity spectrum. A myriad of other states have also been shown to exhibit zero-bias peaks, making it difficult to definitively resolve the presence of MBSs. Recently, it has been proposed that spatially resolved measurements of (differential) shot noise, (differential) conductance and the current Fano factor allow for hallmarks of MBSs to be distinguished. We provide an effective model for long-time incoherent dynamics of a finite N-site Kitaev chain coupled to an STM tip. Using the Keldysh-Schwinger functional-integral formalism, the tip can be integrated out exactly, and by making use of a Hubbard-Stratonovich transformation the fields of the chain are found to obey a classical Langevin equation.

References:

Short bio (50 words) or link to website:

<https://www.uu.nl/staff/OAEastman>

Relevant publications (optional):

Career stage:

Student

Posters II / 58

Signatures of many-body localization of quasiparticles in a flat band superconductor

Authors: Koushik Swaminathan¹; Poula Tadros²; Sebastiano Peotta¹

¹ *Aalto University*

² *University of Turku*

Corresponding Author: koushik.swaminathan@aalto.fi

We construct a class of exact eigenstates of the Hamiltonian obtained by projecting the Hubbard interaction term onto the flat band subspace of a generic lattice model. These exact eigenstates are many-body states in which an arbitrary number of localized fermionic particles coexist with a sea of mobile Cooper pairs with zero momentum. By considering the dice lattice as an example, we provide evidence that these exact eigenstates are, in fact, a manifestation of local integrals of motions of the projected Hamiltonian. In particular, the spin and particle densities retain memory of the initial state for a very long time if localized unpaired particles are present at the beginning of the time evolution. This shows that many-body localization of quasiparticles and superfluidity can coexist even in generic two-dimensional lattice models with flat bands, for which it is not known how to construct local conserved quantities. Our results open new perspectives on the old condensed matter problem of the interplay between superconductivity and localization.

References:

Signatures of many-body localization of quasiparticles in a flat band superconductor, Koushik Swaminathan, Poula Tadros, and Sebastiano Peotta, *Phys. Rev. Research* 5, 043215

Short bio (50 words) or link to website:

<https://fi.linkedin.com/in/koushikswaminathan>

Relevant publications (optional):

Signatures of many-body localization of quasiparticles in a flat band superconductor, Koushik Swaminathan, Poula Tadros, and Sebastiano Peotta, *Phys. Rev. Research* 5, 043215

Career stage:

Student

59

Phase transitions and nonequilibrium dynamics in driven quantum matter

Author: David Weld¹

¹ *UC Santa Barbara*

Corresponding Author: weld@ucsb.edu

Subjecting a quantum system to a time-dependent Hamiltonian can generate a rich array of dynamics and phases of matter. I will discuss results from a sequence of recent cold-atom experiments on kicked and driven quantum matter, highlighting data on anomalous transport, nonequilibrium phase diagrams, quantum thermodynamics, and the interplay between dynamical localization and Anderson localization.

References:

Short bio (50 words) or link to website:

<https://web.physics.ucsb.edu/~weld/>

Relevant publications (optional):

Measuring a localization phase diagram controlled by the interplay of disorder and driving. P. Dotti, Y. Bai, T. Shimasaki, A.R. Dardia, and D.M. Weld, *arXiv:2406.00214* (2024)

Anomalous localization in a kicked quasicrystal. T. Shimasaki, M. Prichard, H.E. Kondakci, J.E. Pagett, Y. Bai, P. Dotti, A. Cao, T.-C. Lu, T. Grover, and D.M. Weld, *Nature Physics* 20, 409–414 (2024).

Interaction-driven breakdown of dynamical localization in a kicked quantum gas. A. Cao, R. Sajjad, H. Mas, E.Q. Simmons, J.L. Tanlimco, E. Nolasco-Martinez, T. Shimasaki, H.E. Kondakci, V. Galitski, and D.M. Weld. *Nature Physics* 18, 1302–1306 (2022).

Career stage:

Professor

60

Exact Results of Fermi Polarons with Ultracold Atoms

Authors: Hui Hu^{None}; Jia Wang^{None}; Xia-Ji Liu^{None}

Corresponding Authors: hhu@swin.edu.au, xiajiliu@swin.edu.au, jiaawang@swin.edu.au

The behaviour of an impurity immersed in a many-body Fermi sea –the so-called Fermi polaron problem –is a long-standing challenge in condensed matter physics and many-body physics. Over the last two decades, there are numerous efforts from ultracold atom community to quantitatively understand the Fermi polaron physics. To date, the ground state of the attractive Fermi polaron has been theoretically predicted and experimentally measured to a great accuracy. However, describing the excited states of Fermi polarons proves to be notably difficult and current theoretical works fail to explain the latest spectral measurement at finite temperature.

In this talk, we present two exact results for the finite-temperature spectral function of Fermi polarons. On the one hand, we propose an exactly solvable model in the immobile heavy polaron limit, which exactly establishes various salient quasiparticle features in the spectral function. On the other hand, we derive an exact set of equations of the spectral function for mobile Fermi polarons, by using the diagrammatic theory and by including particle-hole excitations of the Fermi sea shake-up to arbitrarily high orders. This provides a very rare case that a quantum many-body system can be exactly solved by working out the complete Feynman diagrams. Our exact results of Fermi polarons might be used to better understand the intriguing polaron dynamical responses in two or three dimensions, whether in free space or within lattices.

References:

Short bio (50 words) or link to website:

<https://www.swinburne.edu.au/research/our-research/access-our-research/find-a-researcher-or-supervisor/researcher-profile/?id=hhu>

Relevant publications (optional):

Career stage:

Professor

61

Non-Hermitian band geometry and dynamics of exciton-polaritons

Author: Eliezer Estrecho¹

¹ *The Australian National University*

Corresponding Author: eliezer.estrecho@anu.edu.au

Losses are ubiquitous in exciton-polariton systems, resulting in a short lifetime compared to thermalisation, which makes polariton condensates a good platform for investigating non-equilibrium physics. Additionally, losses can lead to intriguing non-Hermitian effects in systems with non-Hermitian effective Hamiltonians. In this talk, I will present the rich features that can arise in exciton-polariton systems when the losses depend on polarization and momentum [1].

A novel topological winding number can exist in momentum space around exceptional points, which are degeneracies where the eigenstates coalesce. This has direct consequences for the band geometry, resulting in the generalization of the quantum geometric tensor. We propose a method to experimentally measure the non-Hermitian tensor using the polarization of exciton polaritons [2]. The non-Hermiticity also leads to nontrivial dynamics in space, momentum, and pseudospin. For example, a wavepacket can split in both momentum and real space and exhibit self-acceleration without any external potential. Pseudospin defects can also form in momentum space along arcs

where the imaginary parts of the energy eigenvalues cross [3]. The pseudospin dynamics are further modified due to an imbalance in loss rates, which directly impacts the zitterbewegung effect for exciton-polaritons.

References:

- [1] R. Su, E. Estrecho, et al., *Sci. Adv.* 7, eabj8905 (2021).
- [2] Y.-M. R. Hu, E. A. Ostrovskaya, and E. Estrecho, *Phys. Rev. B* 108, 115404 (2023).
- [3] Y.-M. R. Hu, E. A. Ostrovskaya, and E. Estrecho, *Opt. Mater. Express* 14, 664 (2024).

Short bio (50 words) or link to website:

Eliezer Estrecho is an ARC Discovery Early Career Researcher Award Fellow specialising in exciton polaritons across various material and photonic platforms. His research encompasses precise measurements of polariton condensates, such as the interaction strength and the laser linewidth, and exploration of novel effects emerging from non-Hermitian physics.

Relevant publications (optional):

Career stage:

Postdoc

62

Galactic-Scale Superfluidity: True Macroscopic Condensation with Long-Range Interactions?

Author: Nick Proukakis¹

Co-authors: Alex Soto¹; Gary Liu¹; Gerasimos Rigopoulos¹; Milos Indjin¹

¹ *Newcastle University*

Corresponding Authors: alex.soto@newcastle.ac.uk, nikolaos.proukakis@newcastle.ac.uk, m.indjin@newcastle.ac.uk, gerasimos.rigopoulos@ncl.ac.uk, i-kang.liu1@ncl.ac.uk

Dark matter, an integral component of the perceived mass-energy content of the Universe, is usually modelled as a collection of collisionless particles through the established cold dark model (CDM). Despite its impressive success in reproducing large-scale features, increasing evidence is indicating potential shortfalls on shorter (< galactic) scales. An alternative model, “Fuzzy Dark Matter”(FDM), has been gaining increasing attention in the cosmological community: this model postulates the existence of an ultralight bosonic particle exhibiting galactic-size de Broglie wavelengths, facilitating a wave description: central to this model is the suppression of small-scale gravitational collapse due to quantum pressure, which leads to galaxies containing “solitonic cores”. Here I will outline the links between such a cosmological model, laboratory condensates and astrophysical observations, critically discussing implications and open questions.

Using established tools from finite-temperature non-equilibrium condensates [1], I will present a picture of a coherent self-bound galactic-scale solitonic condensate (balancing gravitational attraction against quantum pressure), surrounded by a halo of partially-incoherent particles resembling a quasi-condensate state with spatiotemporally-localised regions of enhanced coherence and a quasi-equilibrium turbulent vortex tangle [2]. Drawing on the standard two-fluid model and atomic bimodal distributions, I will present an extended theory which allows both incoherent and coherent degrees of freedom to be fully self-consistently coupled, in a manner incorporating both the cosmological CDM and FDM models, and also existing cold-atom kinetic and stochastic models [3]. Moreover, by contrasting our findings to astrophysical observations, I will critically analyse the viability of such models [4,5].

Funding: Leverhulme Trust, Horizon 2020.

References:

- [1] N P Proukakis et al. (Eds), *Quantum Gases: Finite Temperature and Non-Equilibrium Dynamics* (Imperial College Press, 2013)
- [2] I-K Liu, N P Proukakis, G Rigopoulos, Coherent and Incoherent Structures in Fuzzy Dark Matter Haloes, *Monthly Notices of the Royal Astronomical Society* 521, 3625 (2023).
- [3] N P Proukakis, G Rigopoulos, A Soto, Unified Description of Corpuscular and Fuzzy Bosonic Dark Matter I and II, *Phys. Rev. D* 108, 083513 (2023) [and follow-on preprint].
- [4] M Indjin, I-K Liu, N P Proukakis, G Rigopoulos, Virialized Profiles and Oscillations of Self-Interacting Fuzzy Dark Matter Solitons, *Phys. Rev. D* 109, 103518 (2024).
- [5] N P Proukakis, G Rigopoulos, A Soto, Hybrid Model of Condensate and Particle Dark Matter: Linear Perturbations in the Hydrodynamic Limit, *Phys. Rev. D* (In Press) (2024).

Short bio (50 words) or link to website:

<https://orcid.org/0000-0003-0126-5820>

https://scholar.google.com/citations?user=YTPCd_AAAAAJ&hl=en

<https://www.ncl.ac.uk/math-physics/people/profile/nikolaosroukakis.html>

Relevant publications (optional):

NP Proukakis, Universality of Bose-Einstein Condensation and Quenched Formation Dynamics, *Encyclopedia of Condensed Matter Physics* (Elsevier, 2nd Edition), arXiv:2304.09541

Career stage:

Professor

63

Observation of Shapiro Steps in an Atomic Superfluid

Authors: Erik Bernhart¹; Herwig Ott¹; Marvin Röhrle¹

¹ *RPTU Kaiserslautern-Landau*

Corresponding Authors: ebernar@rhrk.uni-kl.de, herwig.ott@rptu.de, mroehrle@rhrk.uni-kl.de

Shapiro steps occur in the reverse AC Josephson effect, which is one of the three fundamental effects in superconducting Josephson junctions. When a DC and an AC current are applied simultaneously to a Josephson junction, finite voltage steps are generated across the junction. The voltage is directly linked to the applied frequency via $V = h/(2e) \times f$, where f is the frequency of the alternating current. The series connection of several such junctions in one device corresponds to the current voltage standard.

We have observed Shapiro steps in a Bose-Einstein condensate of rubidium atoms. Following the protocol proposed by Singh et al [1], we move a narrow barrier through the superfluid at a constant velocity, which corresponds to a DC particle current through the barrier. At the same time, we perform a sinusoidal modulation of the barrier velocity with frequency f , which corresponds to an additional AC current through the barrier. When the instantaneous velocity of the barrier exceeds the critical velocity of the superfluid, a finite particle imbalance occurs between the two sides of the barrier. We find that the corresponding chemical potential difference takes on discrete values corresponding to Shapiro steps. We characterize the Shapiro steps and investigate their microscopic dynamics.

- [1] V. P. Singh, J. Polo, L. Mathey, and L. Amico. arXiv:2307.08743 (2023)

References:

- [1] V. P. Singh, J. Polo, L. Mathey, and L. Amico. arXiv:2307.08743 (2023)

Short bio (50 words) or link to website:

<https://physik.rptu.de/ags/ott>

Relevant publications (optional):**Career stage:**

Professor

64

Reynolds similitude of a pure superfluid at low temperatures**Author:** Hiromitsu Takeuchi¹¹ *Osaka Metropolitan University***Corresponding Author:** takeuchi@omu.ac.jp

The Reynolds similitude, a key concept in hydrodynamics, states that two phenomena of different length scales with a similar geometry are physically identical. Flow properties are universally determined in a unified way in terms of the Reynolds number $calR$ (dimensionless, ratio of inertial to viscous forces in incompressible fluids). For example, the drag coefficient c_D of objects with similar shapes moving in fluids is expressed by a universal function of $calR$.

Certain studies introduced similar dimensionless numbers, that is, the superfluid Reynolds number $calR_s$, to characterize turbulent flows in superfluids. However, the applicability of the similitude to inviscid quantum fluids is nontrivial as the original theory is applicable to viscous fluids. This study proposes a method to verify the similitude using current experimental techniques in quantum liquid He-II. A highly precise relation between c_D and $calR_s$ was obtained in terms of the terminal speed of a macroscopic body falling in He-II at finite temperatures across the Knudsen (ballistic) and hydrodynamic regimes of thermal excitations. The Reynolds similitude in superfluids proves the quantum viscosity of a pure superfluid and can facilitate a unified mutual development of classical and quantum hydrodynamics; the concept of quantum viscosity provides a practical correspondence between classical and quantum turbulence as a dissipative phenomenon.

References:

[1] Hiromitsu Takeuchi, Phys. Rev. B 109, L020502 (2024)

Short bio (50 words) or link to website:<http://hiromitsu-takeuchi.appspot.com>**Relevant publications (optional):****Career stage:**

Professor

Posters II / 65**Universal description of massive vortices in superfluids****Author:** Akihiro Kanjo¹**Co-author:** Hiromitsu Takeuchi¹¹ *Osaka Metropolitan University*

Corresponding Author: takeuchi@omu.ac.jp

The point vortex model in hydrodynamics is an effective theory for describing the motion of quantum vortices in superfluids. Regarding the vortex core as a cylinder immersed in the fluid, it behaves like a cylinder with circulation in a perfect fluid. This model has been traditionally used to describe vortex dynamics in superfluid ^4He , where the inertia of the effective cylinder or the vortex mass is neglected as the core size is typically much smaller than the characteristic length scales of the considered system. In contrast, in micro-scale superfluids of ultra-cold atoms, the vortex mass may affect vortex dynamics as was pointed out by several researchers. This work formulates the dynamics of massive point vortices in a unified manner applicable to different superfluid systems and reveals how this effect can be observably enhanced in a uniform superfluid.

References:

Short bio (50 words) or link to website:

<http://hiromitsu-takeuchi.appspot.com>

Relevant publications (optional):

Career stage:

Professor

66

(Non equilibrium) thermodynamics of classical Integrable models in their thermodynamic limit

Author: Leticia Cugliandolo^{None}

Corresponding Author: leticia@lpthe.jussieu.fr

Motivated by recent experimental developments in atomic physics, a large theoretical effort has been devoted to the analysis of the dynamics of quantum isolated systems after a sudden quench. In this talk I will describe the evolution of a family of classical many-body integrable (Neumann) models after instantaneous quenches of the same kind. The asymptotic dynamics of these models can be fully elucidated, and the stationary properties (in the thermodynamic limit) compared to the ones obtained exactly using a Generalised Gibbs Ensemble. The latter can not only be built but also used to evaluate analytically all relevant observables, a quite remarkable fact for an interacting integrable system with a non-trivial phase diagram.

References:

Generalised Gibbs Ensemble for spherically constrained harmonic models
Damien Barbier, Leticia F. Cugliandolo, Gustavo S. Lozano, Nicolás Nessi
arXiv:2204.03081 SciPost Physics 13, 048 (2022)

Quenched dynamics of classical isolated systems: the spherical spin model with two-body random interactions or the Neumann integrable model
Leticia F. Cugliandolo, Gustavo S. Lozano, Nicolás Nessi, Marco Picco and Alessandro Tartaglia
arXiv:1712.07688 J. Stat. Mech. P063206 (2018)

Short bio (50 words) or link to website:

www.lpthe.jussieu.fr/~leticia

Relevant publications (optional):

Career stage:

Professor

Posters I / 67

Active matter in two dimensions

Author: Leticia Cugliandolo^{None}

Corresponding Author: leticia@lpthe.jussieu.fr

Active matter is a new kind of soft matter relevant to describe numerous biological problems with manifold realizations in two dimensions. I will discuss several intriguing aspects of its phase behavior including the melting of an active solide (with special emphasis on the role of dislocations and disclinations) and the mechanisms leading to motility induced phase separation.

References:

Dynamics of Motility-Induced clusters: coarsening beyond Ostwald ripening
Claudio Caporusso, Leticia F. Cugliandolo, Pasquale Digregorio, Giuseppe Gonnella, Demian Levis and Antonio Suma
arXiv:2211.12361 Phys. Rev. Lett. 131, 068201 (2023)

Unified analysis of Topological Defects in 2D systems of Active and Passive disks
Pasquale Digregorio, Demian Levis, Leticia F. Cugliandolo, Giuseppe Gonnella, Ignacio Pagonabarraga
arXiv:2106.03454 Soft Matter 18, 566 (2022)

Motility-Induced Microphase and Macrophase Separation in a two-dimensional Active Brownian Particle system
Claudio B. Caporusso, Pasquale Digregorio, Demian Levis, Leticia F. Cugliandolo, Giuseppe Gonnella
arXiv:2005.06893 Phys. Rev. Lett. 125, 178004 (2020)

Full phase diagram of active Brownian disks: from melting to motility-induced phase separation
Pasquale Digregorio, Demian Levis, Antonio Suma, Leticia F. Cugliandolo, Giuseppe Gonnella and Ignacio Pagonabarraga
arXiv:1805.12484 Phys. Rev. Lett. 121, 098003 (2018)

Short bio (50 words) or link to website:

www.lpthe.jussieu.fr/~leticia

Relevant publications (optional):

Career stage:

Professor

68

Medium-enhanced repulsion between polaron quasiparticles in a quantum gas

Author: Meera Parish^{None}

Corresponding Author: meera.parish@monash.edu

The problem of mobile quantum impurities immersed in a quantum gas [1] has attracted much attention recently owing to its clean realisation in cold atomic gases, as well as its relevance to a variety of

systems spanning a range of energy scales, from semiconductors to neutron stars. Of particular interest is the interactions between polaron quasiparticles—impurities that are dressed by excitations of the surrounding gas—since this has ramifications for the phases of matter that emerge from such impurities. However, there is currently much debate about the nature of the interactions, with the latest experiments even disagreeing on the sign of the interaction strength [2,3].

In this talk, I will focus on bosonic impurities in a Bose-Einstein condensate and I will reveal that the medium actually enhances the existing repulsive interactions between the bosonic impurities. Furthermore, one can show that this is the *dominant* effect at zero temperature in the regime of weak interactions, in contrast to the prevailing wisdom in the field.

References:

- [1] M.M. Parish and J. Levinsen, “Fermi polarons and beyond –Varenna lectures.” arXiv:2306.01215
- [2] C. Baroni et al., “Mediated interactions between Fermi polarons and the role of impurity quantum statistics.” *Nature Physics* 20, 68–73 (2024).
- [3] L. Tan et al., “Bose Polaron Interactions in a Cavity-Coupled Monolayer Semiconductor.” *Phys. Rev. X* 13, 031036 (2023).

Short bio (50 words) or link to website:

Meera Parish is a Professor in theoretical physics and ARC Future Fellow at Monash University. Since obtaining her Ph.D. from the University of Cambridge, she has been a PCTS postdoctoral fellow at Princeton University, and a Lecturer and EPSRC research fellow at University College London.

Relevant publications (optional):

Career stage:

Professor

69

Tweezers, trapped ions, and Rydbergs: a quantum simulation zoo

Author: Arghavan Safavi-Naini¹

¹ *University of Amsterdam*

Corresponding Author: a.safavinaini@uva.nl

Trapped-ions are one of the most mature platforms for quantum computation and quantum simulation. In this talk I will show that by adding other ingredients, such as optical tweezers, neutral atoms, and Rydberg atoms, we can engineer more flexible quantum simulation platforms, as well as new quantum computation architectures.

In trapped-ion quantum simulators the spin-spin interactions mediated by the collective motion of the ions in the crystal (phonons) are tunable range power law interactions. I will show that additional optical tweezer potentials can be used to engineer the phonon spectrum, and thus tune the interactions and connectivity of the ion qubits beyond the power-law interactions accessible in current setups.

Next, I will show that optical tweezers delivering qubit state-dependent local potentials allow us to create a new scalable architecture for trapped-ion quantum computing. Finally, I will discuss how adding one more ingredient, a gas of neutral atoms, allows us to explore ultracold chemistry and exploit ion mediated interactions for infinite range Rydberg blockade and facilitation.

References:

Short bio (50 words) or link to website:

Arghavan Safavi-Naini is an associate professor at the university of Amsterdam and a member of QuSoft at the Centrum Wiskunde and Informatica. She runs a joint experimental-theory group (hyqs.nl) focused on hybrid quantum simulation platforms, quantum algorithms, and numerical methods for open quantum systems.

Relevant publications (optional):

Career stage:

Professor

Posters II / 70

Non-perturbative corrections to the weakly interacting two component Fermi gas

Authors: Carlos Sá de Melo^{None}; Nikolai Kaschewski¹; Axel Pelster¹

¹ RPTU Kaiserslautern-Landau

Corresponding Authors: carlos.sademelo@physics.gatech.edu, kaschews@rhrk.uni-kl.de, axel.pelster@physik.uni-kl.de

A simplified mean-field description of fermionic systems relies on the Hartree-Fock-Bogoliubov (HFB) approach, where the two-particle interaction is decomposed into three distinct channels. A major issue with this method is that the separation between the channels is somewhat arbitrary. Depending on the physical situation to be described, different channels turn out to be important.

In this poster, we present a self-consistently generalized mean-field theory, which is based on introducing a separate weighting factor for each channel. This ansatz removes the arbitrariness of the channel separation by providing an extremization principle for their optimal partitioning.

The power of our technique is illustrated by considering the example of two unpolarized fermionic species with contact interaction. In this case the Fock contribution vanishes and we obtain a coupling between the Hartree and the Bogoliubov channel. This results non only in first beyond mean-field corrections[1,2] already at the mean-field but also decreases the critical temperature in qualitative agreement to particle-hole fluctuations [3]. Due to the non-perturbative nature of the channel coupling we also obtain results which are not captured by any fluctuation theory in one channel alone. This requires the introduction of an effective interaction range as a new length scale and should become relevant for large enough densities. With this our formalism builds a natural theoretical bridge between fermionic superfluidity in ultracold atomic gases and superconductivity in condensed matter physics as well as the realm of nuclei and neutron matter.strong text

References:

- [1] C. A. R. Sá de Melo, M. Randeria, J. R. Engelbrecht, Phys. Rev. Lett. 71, 3202 (1993)
- [2] J. R. Engelbrecht, M. Randeria and C. A. R. Sá de Melo, Phys. Rev. B 55, 15153 (1997)
- [3] L. Gor'Kov and T. Melik-Barkhudarov, Sov. Phys. JETP 13, 1018 (1961)

Short bio (50 words) or link to website:

<https://www-user.rhrk.uni-kl.de/~apelster>

Relevant publications (optional):

Career stage:

Postdoc

Posters I / 71

Bogoliubov theory of 1D anyons in a lattice

Authors: Axel Pelster¹; Bin-Han Tang¹; Martin Bonkhoff^{None}

¹ *RPTU Kaiserslautern-Landau*

Corresponding Authors: tangb@rhrk.uni-kl.de, axel.pelster@physik.uni-kl.de, martinbonkhoff@gmx.de

In a one-dimensional lattice anyons can be defined via generalized commutation relations containing a statistical parameter, which interpolates between the boson limit and the pseudo-fermion limit. The corresponding anyon-Hubbard model is mapped to a Bose-Hubbard model via a fractional Jordan-Wigner transformation, yielding a complex hopping term with a density-dependent Peierls phase. Here we work out a corresponding Bogoliubov theory. To this end we start with the underlying mean-field theory, where we allow for the condensate a finite momentum and determine it from extremizing the mean-field energy. With this we calculate various physical properties and discuss their dependence on the statistical parameter and the lattice size. Among them are both the condensate and the superfluid density as well as the equation of state and the compressibility. Based on the mean-field theory we then analyse the resulting dispersion of the Bogoliubov quasi-particles, which turns out to be in accordance with the Goldstone theorem. In particular, this leads to two different sound velocities for wave propagations to the left and the right, which originates from parity breaking.

References:

Short bio (50 words) or link to website:

<https://www-user.rhrk.uni-kl.de/~apelster>

Relevant publications (optional):

Career stage:

Postdoc

Posters I / 72

Unravelling Interaction and Temperature Contributions in Unpolarized Trapped Fermionic Atoms in the BCS Regime

Authors: Artur Widera¹; Axel Pelster¹; Felix Lang¹; Jennifer Koch¹; Sejung Yong¹; Sian Barbosa¹

¹ *RPTU Kaiserslautern-Landau*

Corresponding Authors: langf@rhrk.uni-kl.de, yong@rhrk.uni-kl.de, jekoch@rhrk.uni-kl.de, axel.pelster@physik.uni-kl.de, widera@physik.uni-kl.de, cbarbosa@physik.uni-kl.de

In the BCS limit density profiles for unpolarized trapped fermionic clouds of atoms are largely featureless. Therefore, it is a delicate task to analyze them in order to quantify their respective interaction and temperature contributions. Temperature measure-

ments have so far been mostly considered in an indirect way, where one sweeps isentropically from the BCS to the BEC limit. Instead we suggest here a direct thermometry, which relies on measuring the line density and comparing the obtained data with a Hartree-Bogoliubov mean-field theory combined with a local density approximation. In case of an attractive interaction between two-components of 6 Li atoms trapped in a tri-axial harmonic confinement we show that minimizing the error within such an experiment-theory collaboration turns out to be a reasonable criterion for analyzing in detail measured densities and, thus, for ultimately determining the sample temperatures. The findings are discussed in view of various possible sources of errors.

References:

Short bio (50 words) or link to website:

<https://www-user.rhrk.uni-kl.de/~apelster>

Relevant publications (optional):

Career stage:

Postdoc

73

Shear flow and vortex array instabilities in annular strongly-correlated atomic superfluids

Author: Giacomo Roati¹

¹ *CNR-INO*

Corresponding Author: giacomo.roati@cnr.it

At the interface between two fluid layers in relative motion, infinitesimal fluctuations can grow exponentially, generating vorticity and causing the laminar flow to break down. Here, we study this scenario by creating two counter-rotating flows in annular atomic Fermi superfluids across the BEC-BCS crossover [1]. Due to the continuity of the superfluid wavefunction and the quantisation of circulation, the superfluids cannot maintain a continuous vortex sheet. Instead, we observe the formation of a regular array of quantised vortices forming along the shear layer. This vortex array is unstable. We connect its dynamics to the instability of the counter-propagating flows, establishing a clear link between shear flow and vortex instabilities. Our work opens the prospects for exploring out-of-equilibrium phenomena such as vortex matter phase transitions and the spontaneous emergence and decay of two-dimensional quantum turbulence in strongly-correlated superfluids.

[1] D. Hernández-Rajkov et al., *Nature Physics* 20, 939 (2024)

References:

Short bio (50 words) or link to website:

My scientific interests regard quantum simulation with strongly-correlated atomic Fermi systems. As Research Director, I have however the ambition of fostering the cooperation of the different scientific areas of CNR-INO, supporting and conducting cross-fertilisation actions and projects.

Relevant publications (optional):

Career stage:

Professor

77

Prescod-Weinstein

78

Driven-dissipative spinor superfluids: a compact Kardar-Parisi-Zhang dynamics of the phase**Author:** Marzena Szymanska¹¹ *University College London***Corresponding Author:** m.szymanska@ucl.ac.uk

Driven-dissipative quantum fluids can differ substantially from their equilibrium counterparts. The long-wavelength phase dynamics of a polariton/photon condensate has been shown to obey Kardar-Parisi-Zhang (KPZ) equation. Since the phase is a compact variable, vortices in 2D and phase slips in 1D can proliferate destroying the KPZ scaling. The interplay between KPZ physics and topological defects is currently subject of great interest, especially in polariton context [1,2,3]. Here, we consider multicomponent system relevant to polariton condensate with different polarisations. The effective theory for Z₂ degenerate coupled condensates with U(1) \times U(1) symmetries maps onto coupled multicomponent KPZ equations. We perform dynamical renormalisation group analysis as well as exact numerical simulations to place polariton condensates in the subspace of a generally rich flow diagram.

Career stage:**Relevant publications (optional):****Short bio (50 words) or link to website:****References:**

- [1] A. Ferrier, A. Zamora, G. Dagvadorj, and M.H. Szymańska Phys. Rev. B 105, 205301 (2022);
- [2] A. Zamora, N. Lad, and M. H. Szymanska Phys. Rev. Lett. 125, 265701 (2020);
- [3] A. Zamora, L. M. Sieberer, K. Dunnett, S. Diehl, and M. H. Szymańska Phys. Rev. X 7, 041006 (2017);
- [4] H. Weinberger, P. Comaron and M. Szymanska “Multicomponent KPZ in Degenerate Coupled Condensate” in preparation.

79

Quartet superfluid in mass-imbalanced ultracold Fermi mixtures

In this talk, I will introduce our recent works on universal few-body clusters and the resulted high-order fermion superfluid in mass-imbalanced Fermi mixtures. First, we exactly solve the (N+1) problems in 2D with N=3 and 4, where a light atom interacts with N heavy fermions via contact potentials. It is found that the critical heavy-light mass ratios to support a (3+1) tetramer and a (4+1) pentamer

are sufficient low to be accessible by a number of mass-imbalanced Fermi mixtures now available in cold atoms laboratories. Further, we study the associated few-body correlations in modifying the Fermi polaron properties and fermion superfluidity of a many-body heavy-light system. In particular, we identify a new fermion superfluidity, called the quartet superfluid (QSF), well beyond the conventional pairing framework in such simple two-component Fermi system. This superfluid phase corresponds to the condensation of quartet (or tetramer) clusters and thus features high-order correlations, as manifested in the momentum-space crystallization of pairing field and the density distribution of heavy fermions. Finally, we explore the dimensional crossover of various universal clusters from 3D to 2D, and show that these clusters are very robust against the presence of an axial confinement and a finite effective range. This suggests the detectability of few-body clusters and quartet superfluid in realistic quasi-2D ultracold Fermi mixtures.

Career stage:

Relevant publications (optional):

Short bio (50 words) or link to website:

References:

80

Wave breaking and multisoliton fission in a chip-scale superfluid waveflume

Corresponding Author: c.baker3@uq.edu.au

In this talk I will present research interfacing cavity optomechanics and superfluid physics for the study of nonlinear wave phenomena.

Building upon our previous work in superfluid optomechanics [1], I will present a novel sensor architecture formed by covering nanofabricated silicon photonic crystal beams with a thin superfluid helium-4 film. This creates an optically addressable quasi-one-dimensional wave tank containing a few femtoliters of superfluid helium, upon which waves can be generated, propagate and be read-out.

Superfluid helium's characteristics present a unique opportunity for the study of nonlinear wave propagation. Indeed, thanks to superfluid helium's vanishing viscosity, the depth of the film h can readily be made as small as a few nanometers without wave attenuation—something impossible to do with classical fluids. Our platform thus enables us to generate waves whose aspect ratio (defined as the wavelength over depth λ/h) exceeds 10,000:1, two orders of magnitude larger than that achievable in the world's largest wave tanks and exceeding that of the most extreme terrestrial phenomena such as tsunamis. This, combined with our recently developed ability to engineer strong fountain pressure forces [2], now allows us to combine within a single device high spatial and temporal resolution along with strong actuation capabilities.

Leveraging these unique characteristics, I will show how our superfluid wave tank enables us to generate and measure (within a sub-millimetre-sized device in a laboratory setting) a rich variety of superfluid nonlinear wave phenomena for the first time, including wavebreaking, multisoliton fission and optomechanical dissipative solitons [3]—opening up the way for the study of extreme regimes of nonlinear hydrodynamics on a chip.

Career stage:

Relevant publications (optional):

Short bio (50 words) or link to website:

References:

- [1] X. He et al., *Nature Physics* 16, 4 (2020); Y. P. Sachkou et al., *Science* 366, 1480 (2019), G. Harris et al, *Nature Physics* 12, 8 (2016); W. W. Wasserman et al., *Opt. Express* 30, 30822 (2022); C. Baker et al., *New J. Phys* 18, 123025 (2016).
- [2] A. Sawadsky et al., *Science Advances* 9, eade3591 (2023)
- [3] J. Zhang et al., *Optomechanical Dissipative Solitons*, *Nature* 600, 75 (2021)