


# Gauge invariance and thermodynamic stability of rotating magnetized systems

Tokyo Univ. of Science  
Kazuya Mameda


K. Fukushima, K. Hattori and KM, arXiv:2407.\*\*\*\*\* [hep-ph]

# QCD matter under rotation



- ✓ elementary particles affected by  $\vec{\Omega}$  (= source of angular momentum)
- ✓  $\vec{\Omega} \parallel \vec{B}$  is more crucial than either  $\vec{\Omega}$  or  $\vec{B}$

# Early attempt : Thermodynamics




Landau-Lifshitz (1958) Vilenkin (1979)

$$Z = \text{tr} \exp[-\beta(H - \Omega \mathcal{J})] \longleftrightarrow H - \mu N$$

Chen-Fukushima-Huang-Mameda (2016)

$$Z = \det \left[ \mathcal{D}_0 - \gamma^0 \Omega (L + S) \right]$$

inverse magnetic catalysis





# Early attempt : Transport



---

Hattori-Yin (2016)  
Kubo formula

$$\rho = \frac{eB\Omega}{4\pi^2}$$



Zeeman coupling



spin-rotation coupling



# Early attempt : Transport


---

Hattori-Yin (2016)  
Kubo formula

$$\rho = \frac{eB\Omega}{4\pi^2}$$

$$\vec{B} \parallel \vec{\Omega}$$

# of   $>$  # of 



# Puzzle on magneto-vortical charge

---

Kubo formula

$$\rho = \frac{eB\Omega}{4\pi^2} \quad \text{Hattori-Yin (2016)}$$

free energy

$$\rho = \frac{eB\Omega}{4\pi^2} + (\text{divergence w.r.t. AM})$$

Chen-Fukushima-Huang-Mameda (2016)

Ebihara-Fukushima-Mameda (2017)

# Answer (as of June)

---

Kubo formula

$$\rho = \frac{eB\Omega}{4\pi^2} \quad \text{Hattori-Yin (2016)}$$

free energy

$$\rho = \frac{eB\Omega}{4\pi^2} + \text{(divergence w.r.t. AM)}$$

# Final answer

---

Fukushima-Hattori-Mameda (in prep.)

correct Kubo formula

$$\rho = -\frac{eB\Omega}{4\pi^2}$$

correct free energy

$$\rho = -\frac{eB\Omega}{4\pi^2}$$

I will convince you!



# Choice of angular momenta

---

$$Z = \det \left[ \mathcal{D}_0 - \gamma^0 \Omega (L + S) \right]$$

Chen-Fukushima-Huang-Mameda (2016)


$$L_{\text{can}} = -i(x\partial_y - y\partial_x) \quad \text{conserved AM}$$

this work

$$L_{\text{kin}} = -i(xD_y - yD_x) \quad \text{gauge invariant AM}$$


# Classical interpretation

---




# Classical interpretation

---



$$H - \Omega L \quad \text{unstable}$$

# Classical interpretation




$$\begin{aligned}
 e\vec{E} &= -eB\Omega\vec{r} \\
 &= -\vec{\nabla} [\Omega(L - L_{\text{kin}})] \\
 &\text{for symmetric gauge}
 \end{aligned}$$

$$H + \Omega(L - L_{\text{kin}}) - \Omega L = H - \Omega L_{\text{kin}} \quad \text{stable}$$

gauge invariance  $\longleftrightarrow$  thermodynamic stability

# Quantum mechanics



Landau level basis  $|n, m\rangle \propto (a^\dagger)^n (b^\dagger)^m |0,0\rangle$

kinetic energy

$$\vec{\Pi}^2 = eB(2a^\dagger a + 1)$$

distance from origin

$$\vec{X}^2 = (2b^\dagger b + 1)/eB$$


# Angular momenta

---

$$L_{\text{kin}} = x\Pi_y - y\Pi_x = \Lambda + \Delta$$


$$\Lambda = (x - X)\Pi_y - (y - Y)\Pi_x$$

$$\Delta = X\Pi_y - Y\Pi_x$$



cyclotron motion

$$= -(2a^\dagger a + 1)$$





circular motion of guiding center

$$= i(a^\dagger b^\dagger - ab)$$

# Lowest Landau Level (LLL)

---

$$\langle J_{\text{kin}} \rangle_{\text{LLL}} = \langle \Lambda + \Delta + S \rangle_{\text{LLL}} = -1/2$$





# Lowest Landau Level

---

$$\langle J_{\text{kin}} \rangle_{\text{LLL}} = \langle \Lambda + \Delta + S \rangle_{\text{LLL}} = -1/2$$

$$\vec{B} \parallel \vec{\Omega}$$

# of  < # of 

This suggests

$$\rho = -\frac{eB\Omega}{4\pi^2}$$



# LLL Pressure

---

Fukushima-Hattori-KM (in prep.)

$$Z = \det \left[ \mathcal{D}_0 - \gamma^0 \underline{\Omega(\Lambda + \Delta + S)} \right]$$

$\nu = -\Omega/2$  (LLL)

$$P_{\text{LLL}} = \frac{eB}{2\pi} \int \frac{dp_z}{2\pi} \left[ \ln \left( 1 + e^{-\beta(\epsilon - \nu)} \right) + \ln \left( 1 + e^{-\beta(\epsilon + \nu)} \right) \right]$$

massless limit

$$\rho = \frac{\partial P_{\text{LLL}}}{\partial \nu} = -\frac{eB\Omega}{4\pi^2}$$



# Anatomy

---

for any  $B$       only for strong  $B$

$$\rho = \frac{\partial P_{LLL}}{\partial \mu} = \frac{eB\Omega}{4\pi^2} - \frac{eB\Omega}{2\pi^2} + \frac{eB\mu}{2\pi^2}$$

spin orbit

$$J = \frac{\partial P_{LLL}}{\partial \Omega} = \frac{eB\mu}{4\pi^2} - \frac{eB\mu}{2\pi^2} + \frac{eB\Omega}{8\pi^2}$$

$\sim J_{\text{CSE}}^5/2$

Only **the spin part** is due to chiral anomaly Yang-Yamamoto (2021)

# Summary

---

- ✓ reformulate gauge-invariant and stable thermodynamics
- ✓ discover cyclotron contribution to magneto-vortical charge
- ✓ identify the anomaly-related part as the spin contribution
- ✓ kinder to lattice than pure rotation : no boundary effect, homogeneity
- ✓ applicability to various fields

HIC : spin polarization under strong B

neutron stars, electron systems, cold atoms, quantum optics