

# Exotic Higgs decay $h \rightarrow Za$ in the final state of two muons and two $\tau$ leptons

Monday 21 October 2024 17:00 (20 minutes)

Exotic Higgs decays are among the most promising areas to be explored at the High Luminosity Large Hadron Collider, given the unprecedentedly large amount of the Standard Model-like Higgs bosons that will be produced. In this context, we introduce a new search channel where a Higgs boson with a mass of 125 GeV decays to a Z boson and a light pseudoscalar (a), in the final state with two muons and two  $\tau$  leptons ( $h \rightarrow Za \rightarrow \mu\mu\tau\tau$ ). In the preliminary analysis, performed at a center-of-mass energy of  $\sqrt{s} = 13$  TeV, masses of the pseudoscalar boson between 3.5 and 30 GeV are probed, focusing on the kinematic region where the  $h \rightarrow Za$  decay is on-shell. We analyze several collider variables to propose a suitable signal region and consider the different decay channels for the di-tau system, as sensitivity is strongly influenced by the  $\tau\tau$  reconstruction efficiency. Model-independent limits are set at the 95% confidence level as a function of  $m_a$ . Upper limits are also established within the Axion-Like Particle (ALP) framework, based on a specific model motivated by the anomalous  $g - 2$  measurement.

**Authors:** RAMOS, Camila; NO, Jose Miguel (IFT-UAM/CSIC); Dr ZURITA, José Francisco (IFIC - Univ. of Valencia and CSIC (ES)); Dr SANDÁ SEOANE, Rosa María (Instituto de Física Teórica UAM-CSIC)

**Presenter:** Dr SANDÁ SEOANE, Rosa María (Instituto de Física Teórica UAM-CSIC)

**Session Classification:** Contributed Talks