

Sustainability in the subatomic sciences

Véronique Boisvert (b. 329.7 CO₂ ppm)
@VBoisvertRHULPP



ROYAL
HOLLOWAY
UNIVERSITY
OF LONDON

8th April 2024 (426.7 CO₂ ppm – 4.0 ppm (0.95%) 1 year change!)


Outline

- The climate emergency
- CO₂e emissions & solutions from:
 - Accelerators (construction/operation)
 - Detectors
 - Computing
 - Rest (travel, conferences, buildings, etc.)
- Possible recommendations
- **Disclaimer:**
 - I'm not a climate/energy scientist!
 - My research is on ATLAS, so energy frontier bias!





Climate Change: an emergency

- UK parliament first to approve a motion to declare an “environment and climate emergency” on 1st May 2019
- Of the top 10 GHG emitters, only Japan, Canada and the EU have **legally** binding target of “net zero emissions by 2050 (2045)”
 - **The pandemic was a blip (lessons)**
- IPCC 2015 Paris agreement: aim to stay “below 2°C” so focus on 1.5 °C
 - NDC: Countries make pledges for how to achieve this (and then increase those pledges over time)
 - Climate Action Tracker: “With all target pledges, including those made in Glasgow, global greenhouse gas emissions in 2030 will still be around twice as high as necessary for the 1.5 °C limit”



IPCC AR6






Ice ages: ~ -5°C
 +4°C: civilization breakdown...

Emissions from accelerators: operations

- CERN now releases [Environment reports](#) (1st: 2017-18, 2nd: 2019-20, 3rd: 2021-22)
- CERN peak power: ~180 MW (~ 1/3 of Geneva)
- Per year: ~ 1.2 TWh (~ 2% of Switzerland, 0.03% of Europe)
- LHC: ~55% of CERN's E consumption
- Electricity mainly comes from France: 90% carbon free (2022)

Electrical power distribution 2018



Emissions from FNAL

- In the US, DOE requirements to report yearly on environmental impacts including emissions
- REC: Renewable Energy Certificates

[2019 FNAL
Environment
Report](#)

2019:
CERN used **4**
times more
electrical E
than FNAL
and yet had **4**
times less
Scope 2
emissions

Scope 1 & 2 Greenhouse Gas Emissions

Goal: Reduce direct GHG emissions by 50 percent by FY 2025 relative to FY 2008 baseline
Interim Target (FY 2019): -31.0%

Current Performance: -62.5%

| | FY 2008 | FY 2019 | % Change |
|----------------------------------|------------------|------------------|---------------|
| Facility Energy | 343,366.8 | 161,122.7 | -53.1% |
| Non-Fleet V&E Fuel | 142.6 | 186.6 | 30.9% |
| Fleet Fuel | 691.6 | 0.0 | -100.0% |
| Fugitive Emissions | 40,165.1 | 139.1 | -99.7% |
| On-Site Landfills | 0.0 | 0.0 | N/A% |
| On-Site WWT | 0.0 | 0.0 | N/A% |
| Renewables | 0.0 | 0.0 | N/A% |
| RECs | 0.0 | -17,435.4 | N/A |
| Total (MtCO₂e) | 384,366.1 | 144,013.0 | -62.5% |



Scope 3 Greenhouse Gas Emissions

Goal: Reduce indirect GHG emissions by 25 percent by FY 2025 relative to FY 2008 baseline
Interim Target (FY 2019): -13.0%



Current Performance: -51.0%

| | FY 2008 | FY 2019 | % Change |
|----------------------------------|-----------------|-----------------|---------------|
| T&D Losses* | 22,287.8 | 7,306.8 | -67.2% |
| T&D RECs Credit | 0.0 | -1,148.5 | N/A |
| Air Travel | 2,215.8 | 2,530.1 | 14.2% |
| Ground Travel | 168.9 | 128.5 | -23.9% |
| Commute | 4,633.3 | 5,392.5 | 16.4% |
| Off-Site MSW | 191.8 | 247.7 | 29.1% |
| Off-Site WWT | 4.8 | 11.0 | 129.2% |
| Total (MtCO₂e) | 29,502.4 | 14,468.1 | -51.0% |

* Includes T&D losses for purchased renewable electricity


Emissions from accelerators: construction

- Potential future of energy frontier: [FCC](#) (ee then hh)
 - ~100 km tunnel, caverns, buildings, roads, etc.
- Concrete needed for the tunnel, which means (Portland) cement!
- Half of emissions from Portland clinker ([ref](#))
- Ken Bloom and my rough calculation:
 - ~260k tonnes of CO₂ emissions
- [Paper](#) on emissions from road tunnels:
 - Lowest estimate: ~500k tonnes CO₂ emissions
- Comparison: Using [report](#) for CO₂e for construction of buildings: = building 8 London Shards!
- 1.4% of CH CO₂e emissions (2016)
- **Plant 6 million trees!**



Emissions from accelerators: solutions

- District heating:
 - Hot water from LHC cooling at Point 8 ready to heat 8000 homes in Ferney-Voltaire, CERN also looking at Point 2 and 5, and Point 1 could heat CERN building on Meyrin site
- Since 2011 series of workshops: **Energy for Sustainable Science at Research Infrastructures**, 7th one: September 2024 in Madrid
- Long-standing R&D in lowering **accelerator power** requirements
 - Eg Energy-Recovery in a Laser-Driven Plasma Wakefield Acceleration





Physics Today **62**, 3, 44 (2009);
<https://doi.org/10.1063/1.3099645>

Emissions from detectors

- Dominant CO₂e emissions from CERN: gases used in experiments!
- Scope 1: direct emissions from organization/vehicles etc.
- Scope 2: indirect emissions from electricity generation, heating, etc.
- Scope 3: all other indirect emissions, upstream and downstream (business travel, personnel commutes, catering, etc.)



| GROUP | GASES | tCO ₂ e 2021 | tCO ₂ e 2022 |
|---------------------------------|---|-------------------------|-------------------------|
| Perfluorocarbons (PFCs) | CF ₄ , C ₂ F ₆ , C ₃ F ₈ , C ₄ F ₁₀ , C ₆ F ₁₄ | 55 921 | 68 989 |
| Hydrochlorofluorocarbons (HFCs) | HFC-23 (CHF ₃) HFC-32 (CH ₂ F ₂) HFC-134a (C ₂ H ₂ F ₄) HFC-404a HFC-407c HFC-410a HFC-507 | 36 557 | 86 211 |
| Other F-gases | SF ₆ , NF ₃ | 16 838 | 18 355 |
| Hydrofluoroolefins (HFO)/HFCs | R-449 R1234ze NOVEC 649 | 86 | 199 |
| | CO ₂ | 13 771 | 10 419 |
| Total Scope 1 | | 123 174 | 184 173 |



Emissions from detectors

- SF6, HFCs and PFCs: particle detection
- HFCs and PFCs: detector cooling
- HFCs: air conditioning systems
- SF6: also used for electrical insulation in power supply systems


Particle detection



Gas recirculation is 90%

Emissions from detectors: solutions



- 2020: CERN launched a working group on managing F-gases, with representatives from the departments concerned and the large LHC experiments. The group looked at issues such as the implementation of a centralised F-gas procurement policy, leak detection, replacement alternatives, training courses for personnel handling F-gases, and improving traceability and reporting.



Emissions from detectors: solutions

- Crucial to do R&D in finding replacements (eco-gases) and ensure 100% leak-free and 100% recirculation
 - CERN has tested NOVEC 649:
Equivalent radiation stability to C_6F_{14} used as a liquid coolant in all LHC experiments

G. D. Hallewell



The “green” use of fluorocarbons in Cherenkov detectors and silicon tracker cooling systems: challenges and opportunities in an unfolding era of alternatives

Embedded emissions from accelerators & detectors

HECAP+ 2023

- Future projects need to compute the full **life cycle analysis** of emissions of all accelerator and detector components

Best Practice 6.1: Life cycle data for a silicon wafer

The ecological impacts of a 1 cm² silicon wafer (thickness 775 μm, diameter 300 mm, weight 0.128 kg) as identified in 2000, are summarised in



Table 6.1 [194].

| Inputs | Quantity | Outputs | Quantity |
|---|----------------------------|--|----------------------------|
| Hydrogen chloride HCl (hydrochloric acid) | 0.000675 kg | Co-products: Si in other co-products | 0.000286 kg |
| Graphite (as electrode material) | 0.000163 kg | Co-products: Silicon tetrachloride | 0.00415 kg |
| Wood chips | 0.00183 kg | Co-products: Si residues for solar cells | 65.2 × 10 ⁻⁶ |
| Petroleum coke | 0.000597 kg | Polished silicon wafer | 1 cm ² |
| Quartz | 0.00486 kg | | |
| Electricity | 0.385 kWh | | |
| Dry wood | 0.00398 kg | | |
| Air emissions | Quantity | Discharge to Water | Quantity |
| CH ₄ | 68.8 × 10 ⁻⁶ kg | Metal chlorides | 0.000787 kg |
| CO | 0.000167 kg | | |
| CO ₂ | 0.00833 kg | Waste | Quantity |
| Ethane | 29 × 10 ⁻⁶ kg | SiO ₂ | 16.3 × 10 ⁻⁶ kg |
| H ₂ O | 0.00188 kg | | |
| Methanol | 85.1 × 10 ⁻⁶ kg | | |
| NO _x | 13.8 × 10 ⁻⁶ kg | | |
| Particulate matter | 0.000201 kg | | |
| SO ₂ | 34.4 × 10 ⁻⁶ kg | | |
| Hydrogen | 0.000125 kg | | |

Table 6.1: Inputs, outputs and emissions of silicon wafer production [194].

Emissions from Computing

- Global IT sector could be 2-6% of global CO₂e emissions, growing to 20% by 2030
- 70% from data centres and communication networks
- HEP uses Grid centres all over the world, yet emissions from electricity vary wildly



Solutions:

- Choose sites with green electricity...
- [Green500 list](#)
- Optimize your code ;-)


Far future (2040):

- All OECD electricity grids will be emissions free...
- But huge demand for electricity

Emissions from Computing


- **Embedded emissions...**
 - 326 (620) kg CO₂e 13' (16') MacBook Pro, 128 GB (1 TB) storage
- **... far outnumber running emissions (80-85% of lifetime emissions)**
 - 2g (3g) CO₂e/h MacBook Pro
 - 10g CO₂e/h average-efficient laptop
 - 50g CO₂e/h desktop with screen
 - + 22g CO₂e/h for servers, networks
- Replacing farms less often can help a lot
- In general ASIC/FPGA/GPU/TPU use less power than CPUs, but exact numbers depend on software/architecture

Numbers
from Mike
Berners-Lee



Emissions from Travel

- Commuting, conferences, etc.
- A nearly carbon-neutral conference model
- Although aviation is 2.4% (2018) of global emissions (more than Australia or Italy or France!), rate of growth is large and carbon neutral flights long way off (CO₂ emissions increased by 32% from 2013-2018)
- Environmental groups calling for frequent flyers levy since eg in 2015 only 12% of people in England took 3 flights or more!
- Carbon offsetting as short term mitigation? controversial





| | AGU Fall Meeting 2019 | ICHEP Melbourne 2012 | ICHEP Valencia 2014 | ICHEP Chicago 2016 | ICHEP Seoul 2018 | ICHEP Prague 2020 (virtual) |
|--|-----------------------|----------------------|---------------------|--------------------|------------------|-----------------------------|
| Number of participants | 24,009 | 764 | 966 | 1,120 | 1,178 | 2,877 |
| GHG emissions per participant [kg CO ₂ e] | 2,883 | 8,432 | 1,902 | 2,699 | 2,648 | 0 |

Table 5.3: Total number of participants of recent ICHEP conferences and the GHG emissions per participant. The corresponding numbers for the American Geophysical Union (AGU) Fall Meeting [147] are shown for reference.

Emissions from food & Total

- [IPCC report in August 2019 on Land Usage](#)
- How about migrating our PP catering (meetings, conf, workshops) in that direction?



Mike Berners-Lee: average UK: 13 tonnes of CO₂e

Remaining carbon budget:

(50% chance of staying < 1.5°C) 460 GtCO₂

Per year per person: 2.2t

POSSIBLE RECOMMENDATIONS

European Strategy Update 2020

7



Environmental and societal impact

A. The energy efficiency of present and future accelerators, and of computing facilities, is and should remain an area requiring constant attention. Travel also represents an environmental challenge, due to the international nature of the field. *The environmental impact of particle physics activities should continue to be carefully studied and minimised. A detailed plan for the minimisation of environmental impact and for the saving and re-use of energy should be part of the approval process for any major project. Alternatives to travel should be explored and encouraged.*

B. Particle physics, with its fundamental questions and technological innovations, attracts bright young minds. Their education and training are crucial for the needs of the field and of society at large. *For early-career researchers to thrive, the particle physics community should place strong emphasis on their supervision and training. Additional measures should be taken in large collaborations to increase the recognition of individuals developing and maintaining experiments, computing and software. The particle physics community commits to placing the principles of equality, diversity and inclusion at the heart of all its activities.*

C. Particle physics has contributed to advances in many fields that have brought great benefits to society. Awareness of knowledge and technology transfer and the associated societal impact is important at all phases of particle physics projects. *Particle physics research centres should promote knowledge and technology transfer and support their researchers in enabling it. The particle physics community should engage with industry to facilitate knowledge transfer and technological development.*

D. Exploring the fundamental properties of nature inspires and excites. It is part of the duty of researchers to share the excitement of scientific achievements with all stakeholders and the public. The concepts of the Standard Model, a well-established theory for elementary particles, are an integral part of culture. *Public engagement, education and communication in particle physics should continue to be recognised as important components of the scientific activity and receive adequate support. Particle physicists should work with the broad community of scientists to intensify engagement between scientific disciplines. The particle physics community should work with educators and relevant authorities to explore the adoption of basic knowledge of elementary particles and their interactions in the regular school curriculum.*

a) The energy efficiency of present and future accelerators, and of computing facilities, is and should remain an area requiring constant attention. Travel also represents an environmental challenge, due to the international nature of the field. *The environmental impact of particle physics activities should continue to be carefully studied and minimised. A detailed plan for the minimisation of environmental impact and for the saving and re-use of energy should be part of the approval process for any major project. Alternatives to travel should be explored and encouraged.*


Snowmass white paper: [\[2203.12389\]](https://arxiv.org/abs/2203.12389) [Climate impacts of particle physics \(arxiv.org\)](https://arxiv.org/abs/2203.12389)

Recommendations

- New experiments and facility construction projects should **report on their planned emissions and energy usage** as part of their environmental assessment
 - Eg LHCb TDR for Phase II, CLIC LCA, ISIS-II LCA (H. Wakeling)
- Review across all international laboratories to ascertain whether emissions are **reported clearly and in a standardized way**
- Take steps to mitigate impact on climate change by setting **concrete reduction goals and defining pathways to reaching them**
 - spend a portion of research time on directly tackling challenges related to climate change

Sustainable Concrete Construction

Knowledge and technology for the environment - Highlights



□ + Others



CERN to partner with industry on innovation to reduce environmental impact of large-scale facilities

In its commitment to minimising its environmental impact and developing technologies that can help society towards a better planet, CERN has formed an innovation partnership with ABB, with the aim of reducing the Laboratory's energy consumption

Environment | 14 June, 2022



CERN and Airbus partnership on future clean aviation

CERN and Airbus UpNext sign a collaboration agreement to assess the use of superconducting technologies for future low-emission aeroplanes.

AerospaceEnvironment | 01 December, 2022



CERN Innovation Programme on Environmental Applications

Recommendations

- Minimize the **travel** emissions of users
- Long-term projects should consider the **evolving social and economic context**
- Actively engage in **learning** about the climate emergency and about the climate impact of particle-physics research
 - See next slide!
- **Promote and publicize their actions** surrounding the climate emergency to the general public and other scientific communities
- **Engage with the broader international community** to collectively reduce emissions

Learn More – Sustainability at STFC

STFC's sustainability learning programme created with The University of Oxford is open for registration.

- The 'Creating a sustainable STFC' on-line course is specifically for STFC staff.
- Participants come away with a personalised action plan that will help them embed sustainability into their work.
- Find out about the course and how to register for one of the upcoming intakes visit the sustainability pages on The Source.

More information about sustainability including STFC's environmental data and targets can be found on the sustainability pages on The Source.



What about your own university/Lab?

Recommendations

- Eg ATLAS Sustainability Forum!
 - atlas-sustainability-forum@cern.ch
- Detailed Recommendations in each area listed in HECAP+ report

Version 1.0, 5 June 2023

Environmental sustainability in basic research

Recommendations – Mobility



Individual actions:

- Re-assess business travel needs, using remote technologies wherever practicable.
- Choose environmentally sustainable means of transport for daily commutes as well as unavoidable business travel, amalgamating long-distance trips where possible.



Further group actions:

- Define mobility requirements and travel policies that minimise emissions, while accounting for the differing needs of particular groups, such as early-career researchers or those who are geographically isolated.
- Re-assess needs for in-person meetings, and prioritise formats that minimise travel emissions and diversify participation by making use of hybrid, virtual or local hub participation, and optimising the meeting location(s).



Further institutional actions:


- Support environmentally sustainable commuting by improving on-site bicycle infrastructure, subsidising public transport and providing shuttle services.
- Disincentivise car travel where viable alternatives exist, facilitate car pooling, and provide on-site charging stations.
- Incentivise the reduction of business travel, e.g., by implementing carbon budgets with appropriate concessions.
- Ensure unavoidable travel is made via environmentally sustainable means through flexible travel policies and budgets, and the use of travel agents that offer multi-modal itineraries. Employ carbon offsetting only as a last resort.
- Remove any requirement on past mobility as an indication of quality in hiring decisions.
- Lobby for improved and environmentally sustainable local and regional transport infrastructure.

Sustainable HEP 2024 -
3rd edition: 10th-12th June
2024: free, online-only,
time zone friendly

INDICO.CERN.CH/E/SUSTHEP24

SUSTAINABLE HEP 2024

3rd International Workshop on Sustainable High Energy Physics
10 – 12 June 2024



The 3rd edition of the Sustainable High Energy Physics (HEP) workshop, will take place Monday 10th through Wednesday 12th June from 14:00 to 17:00 CET. Within three half-days, this free, online-only workshop aims to present the intersection of HEP and the climate crisis, to highlight the sustainable initiatives ongoing in HEP, and to workshop with attendees on positive tangible outcomes. The program will consist of invited talks, panel discussions, workshops and submitted talks accompanied by a discussion forum on Mattermost.

ORGANIZING COMMITTEE

- SHREYASI ACHARYA (INFN BARI)
- JULIETTE ALIMENA (DESY)
- DANIEL BRITZGER (MPP)
- BRENDON BULLARD (SLAC)
- SHREYA SAHA (MCGILL)
- HANNAH WAKELING (OXFORD)

ADVISORY COMMITTEE

- SHANKHA BANJEREE (CHENNAI)
- NIKLAS BEISERT (ETHZ)
- VALERIE DOMCKE (CERN)
- VALERIE LANG (FREIBURG)
- PETER MILLINGTON (MANCHESTER)
- AYAN PAUL (NORTHEASTERN)

CONTACT
indico.cern.ch/e/susthep24
susthep24@physics.ox.ac.uk

10 – 12 JUNE 2024
14:00–17:00 CET
ONLINE VIA ZOOM

SCAN ME



An aerial photograph of a coastal landscape featuring sand dunes, a bay, and distant hills under a clear sky. The text is overlaid on this image.

THE CLIMATE EMERGENCY: CAN
PARTICLE PHYSICS EVER BE
SUSTAINABLE?


DISCUSSION/QUESTIONS

BACK UP


Emissions from detectors: solutions

Presented at DPF 2019

- Alternative gas example: replace CF_4
- CF_4 prevents ageing, improves timing resolution and is a scintillator
- CMS CSC: currently 10% CF_4
 - Reduce concentration to 5%
 - Replace with CF_3I or HFO_{1234ze}
- LHCb RICH studies:
 - CF_4 or C_4F_{10} used for good refractive index
 - Could replace C_4F_{10} with C_4H_{10} but flammable
 - Replace CF_4 with CO_2 : under study
 - Use of SiPM to reduce the chromatic error and increase the yield




S. Easo and O. Ullaland



Gas Recuperation systems at LHC experiments

Sometimes it is not possible to recirculate 100% of the gas mixture due to detector constrains

- Air permeability, max recirculation fraction, impurities, etc.
- A fraction of gas has to be renewed
 - Some gas is sent to the atmosphere
- This fraction of gas mixture can be sent to a recuperation plant where the GHG is extracted, stored and re-used
- Challenges: R&D, custom development, operation and recuperated gas quality
- Gas recuperation also to empty/fill the detectors during LS



LHCb RICH1 and RICH2

During LS2 need to empty the detectors for maintenance/upgrade:

RICH1: 4 m³ of C₄F₁₀

RICH2: 100 m³ of CF₄

Gas recuperation system for empty/filling and cleaning from Air contamination

New gas recuperation system developed for the empty and filling of the detector

CMS CSC

Small Permeability to Air: accumulation of N₂

Need to inject fresh gas continuously to keep N₂ stable and guarantee detector performance

~80 l/h of CF₄ would be lost in exhaust without gas recuperation system


The R134a recuperation system for RPCs

ATLAS and CMS RPC Gas Systems


- Detector volume $\sim 15 \text{ m}^3$
- Gas mixture: $\sim 95\% \text{ C}_2\text{H}_2\text{F}_4$, $\sim 5\% \text{ iC}_4\text{H}_{10}$, $0.3\% \text{ SF}_6$
- Gas recirculation: $\sim 90\%$
- Maximum recirculation validated for RPC detectors
- **Fundamental to repair detector leaks**
- To have the gas at the exhaust of the gas system

R134a and iC₄H₁₀ form an azeotrope

A mixture of liquids whose proportions cannot be altered or changed by simple distillation



$\text{C}_2\text{H}_2\text{F}_4$ recuperation prototype system under study in CMS Experiment



Recuperation efficiency $\sim 80\%$

First $\text{C}_2\text{H}_2\text{F}_4$ recuperation system under construction: installation foreseen beginning of 2023 in CMS experiment

Gas disposal

*Abatement plants are employed when GHGs are polluted
and therefore are not reusable*

In case all studies on recuperation will not bring to efficient recuperation plants,
industrial system able to destroy GHGs avoiding their emission into the atmosphere
have been considered

Quite heavy infrastructure required:


- CH₄/city gas + O₂ supply + N₂ supply
- Waste water treatment
- PFC/HFC are converted in CO₂ + HF acid dissolved in water
- disposal of remaining waste/mud
- To have the gas at the exhaust (600-1000 l/h)



Found also companies available to take PFC/HFC based mixture for disposal:
but extremely expensive


Carbon footprint of PP researchers

- Flying!
 - 15% of E consumption for an **average** UK citizen
 - Particle Physicists fly a lot (esp. seniors)! Let's say, per year:
 - 8 European trips (eg use from London to Zurich): $8 \times 148 \text{ Kg CO}_2$: **1184 Kg CO₂**
 - 1 overseas trip (eg use from London to NYC): **986 Kg CO₂**
 - Total: **2170 Kg CO₂**: ~87 countries where the average citizen **emits less CO₂ in a year** (incl. India, **Morocco**, Peru, **Colombia**)!
 - Using: [Guardian calculator](#)
 - [A nearly carbon-neutral conference model](#)
 - Best calculator: <https://www.atmosfair.de/en/offset/fix/>



Top Annual CO₂ Emitting countries, 2020


(from fossil fuels)



CO₂ emissions per capita vs. share of electricity generation from renewables, 2022

Our World in Data

Carbon dioxide (CO₂) emissions are measured in tonnes per person.




Data source: Global Carbon Budget (2023) and other sources
 OurWorldInData.org/co2-and-greenhouse-gas-emissions | CC BY

List of top CO₂ emitters


Forbes

| Country | 2018 CO2 Emissions in Billion Metric Tons | Global Share | Change Since Kyoto Protocol |
|--------------|---|--------------|-----------------------------|
| China | 9.43 | 27.8% | 54.6% |
| U.S. | 5.15 | 15.2% | -12.1% |
| India | 2.48 | 7.3% | 105.8% |
| Russia | 1.55 | 4.6% | 5.7% |
| Japan | 1.15 | 3.4% | -10.1% |
| Germany | 0.73 | 2.1% | -11.7% |
| South Korea | 0.70 | 2.1% | 34.1% |
| Iran | 0.66 | 1.9% | 57.7% |
| Saudi Arabia | 0.57 | 1.7% | 59.9% |
| Canada | 0.55 | 1.6% | 1.6% |

Economicshelp.org



Emissions pathway



Sustainable HEP

28-30 June 2021
Zoom
Europe/Zurich timezone

Overview
Timetable
Call for Abstracts
Contribution List
Speaker List
Book of Abstracts
Registration
Participant List
Talk Recordings
Closing Statement

Closing Statement

Workshop "Sustainable HEP"

Closing Statement (status: 14th July 2021, 403 signatures)

On 28th–30th June 2021, the [workshop "Sustainable High Energy Physics"](#) took place by videoconferencing means with more than 350 registered participants from around 45 countries and five continents. The aim of this workshop was to initiate a community discussion on how to align the scientific operations within this particular subfield of physics with requirements of climate sustainability. Achieving the latter is a most pressing global issue for the present decade (as evidenced by the [IPCC reports](#)). The main focus of the workshop was on the scientific travel culture and the virtualisation of scientific exchange. The following topics were highlighted at this occasion:

- characteristics of the climate crisis
- best practice examples on the virtualisation of scientific meetings
- challenges for research institutions to improve their climate sustainability
- improvement of global inclusiveness in scientific exchange through virtualisation
- domains of action for large scale experiments to improve their climate sustainability

We are organisers and participants of the workshop as well as members of the High Energy Physics community or related fields of physics. We understand that the climate impact of certain aspects of our field of research is a cause of concern and we assert that there is a need for determined action to align these with the goals of the Paris climate agreement and, more generally, with the needs of a sustainable society. Our aim is to trigger a discussion on how HEP can live up to its responsibility in the global transition to a sustainable and climate-neutral world, while maintaining the high quality of research and international scientific exchange. In this context, we highlight increased inclusiveness as a crucial co-benefit of online formats.

We thus encourage members of our community to discuss and enable suitable implementations of sustainable development for our field. We stress that this is a call to develop a balanced and deliberated approach that brings together the needs of a global HEP community with the needs of climate sustainability. We call on research and funding institutions to adjust the general framework for research accordingly and to facilitate a transformation towards sustainable means. Consequently, we invite the formation of working groups to continue the discussions initiated at the workshop and to conduct further installations of the workshop on related topics of sustainability that deserve discussions in a broader setting.

Signatures

The following persons have signed the statement as individuals on their own behalf. Please note that institutions are mentioned merely to identify the signatories' current scientific affiliations. This statement does not (necessarily) reflect the opinions of these institutions.

[sign here](#)


workshop organisers:

Niklas Beisert (ETH Zürich)
Valerie Domcke (CERN/EPFL)
Astrid Eichhorn (CP3-Origins, University of Southern Denmark)
Kai Schmitz (CERN)

workshop participants:

Also: white paper for Australian
Astronomy: "The imperative to reduce
carbon emissions in astronomy"

Figure 4: Territorial UK greenhouse gas emissions by NC sector, 2019 (%)



Source: Table 1.2, Final UK greenhouse gas emissions national statistics 1990-2019 Excel data tables

Note: LULUCF is land use, land use change and forestry.

World Emissions Clock

Green electricity grids by 2035

- Germany's target updated in 2022
 - The US, Canada and UK have already committed to a similar goal [100% renewable electricity grid by 2035]. Denmark is already aiming for more than 100% renewable power by 2027, Austria 100% by 2030 and Portugal and the Netherlands are well on track with recent plans to expand renewable capacities till 2030.”
- US pledge
- UK CCC plan: