

Indications for New Higgs Bosons in Associated Di-Photon Production

Wednesday, November 6, 2024 9:20 AM (20 minutes)

Statistically significant excesses exist at around 152 GeV in associated di-photon production contained in the sidebands of SM Higgs analyses. They are most pronounced in the $\gamma\gamma + \tau$, $\gamma\gamma + \text{MET}$, $\gamma\gamma + \geq 1\ell + \geq 1b$, $\gamma\gamma + 4j$ signal regions, and can be explained by the Drell-Yan production of new Higgs bosons, i.e. $pp \rightarrow W \rightarrow H^\pm \rightarrow \tau\nu, tb, WZ$. We then specialize our analysis for the real Higgs triplet and two-Higgs doublet models, resulting in a combined significance of $\approx 4\sigma$ and $\approx 4.3\sigma$, respectively.

Primary track

BSM Higgs physics

Is the speaker a PhD student or post-doc?

Yes - My participation will be fully supported by my research group

Primary authors: CRIVELLIN, Andreas (University of Zurich (CH)); MELLADO GARCIA, Bruce (University of Wisconsin); COLORETTI, Guglielmo (University of Zurich (UZH) / Paul Scherrer Institute (PSI)); Dr ASHANU-JJAMAN, Saiyad (SGTB Khalsa College); MAHARATHY, Siddharth Prasad (Institute of Physics); BANIK, Sumit (University of Zurich & PSI)

Presenter: BANIK, Sumit (University of Zurich & PSI)

Session Classification: BSM Higgs physics 3 - sal IX

Track Classification: BSM Higgs physics