

Higgs-boson production in weak-boson fusion and $H \rightarrow b\bar{b}$ decay at NNLO with realistic event selection criteria

Based on [Phys.Rev.D 110 \(2024\) 5, 054017](#)

Ivan Novikov



HIGGS 2024

2024-11-06


In collaboration with Konstantin Asteriadis, Arnd Behring, Kirill Melnikov, and Raoul Röntsch

b -quark Yukawa coupling y_b


[CMS Collaboration, Nature 607, 60–68 (2022)]




The b -quark Yukawa coupling y_b can be measured in $H \rightarrow b\bar{b}$ decay



The $H \rightarrow b\bar{b}$ decay is difficult to measure
due to large number of b -jets from QCD backgrounds




On the other hand, Higgs-boson production in weak-boson fusion (WBF) can be separated from QCD backgrounds by its distinct signature of two back-to-back jets.




We look for events with **two light nearly-back-to-back jets with a high invariant mass** and **two b -tagged jets**.

$$|p_{\perp,b}| > 65 \text{ GeV} > \frac{m_H}{2}$$



These event selection criteria are rather strict and require production of a boosted Higgs boson



$$d\sigma = \text{Br}_{H \rightarrow b\bar{b}} d\sigma_{\text{WBF}} \frac{d\Gamma_{H \rightarrow b\bar{b}}}{\Gamma_{H \rightarrow b\bar{b}}}$$

$$\text{Br}_{H \rightarrow b\bar{b}} = \frac{\Gamma_{H \rightarrow b\bar{b}}}{\Gamma_{H \rightarrow \text{anything}}}$$

$$\int_{\text{inclusive}} d\sigma_{\text{WBF}} d\Gamma_{H \rightarrow b\bar{b}} = \int_{\text{inclusive}} d\sigma_{\text{WBF}} \times \int_{\text{inclusive}} d\Gamma_{H \rightarrow b\bar{b}}$$


$$\int_{\text{fiducial}} d\sigma_{\text{WBF}} d\Gamma_{H \rightarrow b\bar{b}} \neq \int_{\text{fiducial}} d\sigma_{\text{WBF}} \times \int_{\text{fiducial}} d\Gamma_{H \rightarrow b\bar{b}}$$

The event selection criteria introduce a correlation between the production and the decay subprocesses, even in the narrow-width approximation.

- ▶ Weak-boson fusion $pp \rightarrow Hjj$ up to NNLO QCD
[\[Cacciari, Dreyer, Karlberg, Salam, Zanderighi \(2015\)\]](#)
[\[Cruz-Martinez, Gehrmann, Glover, Huss \(2018\)\]](#)

$$\sigma_{\text{fiducial}}^{\text{WBF}}/\text{fb} \approx \underset{\text{LO}}{971} \quad - \underset{\Delta\text{NLO}}{81} \quad - \underset{\Delta\text{NNLO}}{31} + \dots$$

(-8%)
(-3%)




- ▶ Electroweak corrections and interference effects in WBF $pp \rightarrow Hjj$ up to NLO EW ($\sim -5\%$) [\[Ciccolini, Denner, Dittmaier \(2007\)\]](#)
- ▶ Nonfactorizable corrections to WBF $pp \rightarrow Hjj$ at NNLO QCD ($\sim -0.3\%$)
[\[Liu, Melnikov, Penin \(2019\)\]](#) [\[Asteriadis, Brønnum-Hansen, Melnikov \(2023\)\]](#)

NNLO QCD corrections to weak-boson fusion are of order $\sim -3\%$


- ▶ $H \rightarrow b\bar{b}$ with massless b quarks up to N³LO [[Mondini, Schiavi, Williams \(2019\)](#)]
- ▶ $H \rightarrow b\bar{b}$ with massive b quarks up to NNLO [[Behring, Bizoń \(2020\)](#)]

$$\Gamma_{H \rightarrow b\bar{b}} / \text{MeV} \approx \underbrace{1.926}_{\text{LO}} + \underbrace{0.400}_{\Delta\text{NLO}} + \underbrace{0.106}_{\Delta\text{NNLO}} + \dots$$

$(\mu = m_H)$
(+21%)
(+6%)




NNLO QCD corrections to $H \rightarrow b\bar{b}$ decay are of order $\sim +6\%$



With *massless* b quarks b -jet tagging is potentially IRC-unsafe, because a **soft** gluon can split into a $b\bar{b}$ pair, which end up in different jets and change their flavor.

In the $H \rightarrow b\bar{b}$ calculation this **soft singularity** is regulated by a finite b -quark mass.



The available weak-boson-fusion calculations neglect the b -quark mass.

To ensure IRC-safety, we do not tag b jets originating from WBF.

As a result, we can use the standard anti- k_{\perp} jet clustering algorithm.

- ▶ Combined $pp \rightarrow H(\rightarrow b\bar{b})jj$ with NNLO production and LO decay with massless b quarks
[\[Asteriadis, Caola, Melnikov, Röntsch \(2022\)\]](#)

$$\sigma_{\text{fiducial}}/\text{fb} = \underset{\text{LO}}{75.9} \quad - \underset{\Delta\text{NLO}}{5.0} \quad - \underset{\Delta\text{NNLO}}{1.5} + \dots$$



(-7%)
(-2%)

- ▶ New result: $pp \rightarrow H(\rightarrow b\bar{b})jj$ with massive b quarks up to NNLO QCD

$$\sigma_{\text{fiducial}}/\text{fb} = \underset{\text{LO}}{75.6} \quad - \underset{\Delta\text{NLO}}{23.2} \quad - \underset{\Delta\text{NNLO}}{7.8} + \dots$$

(-31%)
(-10%)

There are large negative corrections to the fiducial cross-section: -41% compared to LO!



The impact of scale variation in the decay $H \rightarrow b\bar{b}$ is comparable to that in the WBF production, and does not capture the observed large corrections either

$$d\sigma = \text{Br}_{H \rightarrow b\bar{b}} d\sigma_{\text{WBF}} \frac{d\Gamma_{H \rightarrow b\bar{b}}}{\Gamma_{H \rightarrow b\bar{b}}}$$

$d\Gamma_{H \rightarrow b\bar{b}}$

→

$$\sigma_{\text{fiducial}}/\text{fb} = 75.6 \quad - 5.3 \quad - 5.0 \quad + \dots$$

LO	ΔNLO decay	ΔNNLO decay	
	(-7%)	(-7%)	



$\Gamma_{H \rightarrow b\bar{b}}$

→



$$\Gamma_{H \rightarrow b\bar{b}}/\text{MeV} = 1.926 + 0.400 + 0.106 + \dots$$

$(\mu = m_H)$	LO	ΔNLO	ΔNNLO	
		(+21%)	(+6%)	

Corrections to the total $H \rightarrow b\bar{b}$ decay width $\Gamma_{H \rightarrow b\bar{b}}$ are *positive*, but they are large and *negative* with the used event selection criteria





QCD radiation in the $H \rightarrow b\bar{b}$ decay tends to reduce the **transverse momentum** $p_{\perp,b}$ of the **b -jet**, lowering the probability that they pass the b -jet selection criteria.


$p_{\perp b_2}$ thresholdcumulative
→

With the chosen $p_{\perp b_2}$ threshold the decay corrections do not seem to converge.
Relaxing this threshold seems to improve perturbative convergence,
but might degrade purity of event selection

$p_{\perp b}$ distributions






The K-factor $d\sigma/d\sigma^{\text{LO}}$ is more-or-less flat for distributions of transverse momenta $p_{\perp b}$ for leading ($p_{\perp b_1}$) and subleading ($p_{\perp b_2}$) b jets



The distribution of the transverse momentum $p_{\perp,b\bar{b}}$ of the reconstructed Higgs boson shows stronger suppression at small transverse momentum.

m_{bb} distribution



(b jets originating in WBF are not included)

QCD radiation in the $H \rightarrow b\bar{b}$ decay reduces the invariant mass m_{bb} of the reconstructed Higgs boson. Rarely, QCD radiation from weak-boson fusion can increase this invariant mass.

- ▶ We provide, for the first time, an NNLO-QCD-accurate fully-differential description of the combined WBF process $pp \rightarrow H(\rightarrow b\bar{b})jj$.
- ▶ b -jets originating in WBF are not tagged, a calculation of WBF with *massive* b -quarks would be necessary to account for them.
- ▶ There are large negative corrections, the NNLO fiducial cross-section is $\sim 40\%$ smaller than the LO cross-section.
- ▶ QCD radiation in the $H \rightarrow b\bar{b}$ decay makes a large impact because of stringent restrictions on b -jet momenta.

In the future, it would be interesting to try to resum these fixed-order results and/or match them to a parton shower.

- ▶ We provide, for the first time, an NNLO-QCD-accurate fully-differential description of the combined WBF process $pp \rightarrow H(\rightarrow b\bar{b})jj$.
- ▶ b -jets originating in WBF are not tagged, a calculation of WBF with *massive* b -quarks would be necessary to account for them.
- ▶ There are large negative corrections, the NNLO fiducial cross-section is $\sim 40\%$ smaller than the LO cross-section.
- ▶ QCD radiation in the $H \rightarrow b\bar{b}$ decay makes a large impact because of stringent restrictions on b -jet momenta.

In the future, it would be interesting to try to resum these fixed-order results and/or match them to a parton shower.

Thank you for your attention!

Backup

	$\sigma_{\text{inclusive}}^{\text{WBF}}/\text{fb}$	$\sigma_{\text{fiducial}}^{\text{WBF}}/\text{fb}$	$\sigma_{\text{fiducial}}^{\text{LO } H \rightarrow b\bar{b}}/\text{fb}$	$\Gamma_{H \rightarrow b\bar{b}}/\text{keV}$
LO	4099^{+51}_{-67}	971^{-61}_{+69}	$75.9^{-5.6}_{+6.5}$	1926^{+243}_{-194}
NLO	3970^{+25}_{-23}	890^{+8}_{-18}	$70.9^{+0.2}_{-1.2}$	2327^{+110}_{-112}
NNLO	3932^{+15}_{-10}	859^{+8}_{-10}	$69.4^{+0.5}_{-0.2}$	2432^{+31}_{-47}

$$d\sigma_{\text{WBF}} = d\sigma_{\text{WBF}}^{(0)} + d\sigma_{\text{WBF}}^{(1)} + d\sigma_{\text{WBF}}^{(2)} + \dots$$

$$d\sigma^{\text{N}^n\text{LO}} = \text{Br}_{H \rightarrow b\bar{b}} \sum_{k=0}^n d\sigma_{\text{WBF}}^{(n-k)} \frac{d\Gamma_{H \rightarrow b\bar{b}}^{\text{N}^k\text{LO}}}{\Gamma_{H \rightarrow b\bar{b}}^{\text{N}^k\text{LO}}}$$

$$\implies \int_{\text{inclusive}} d\sigma^{\text{N}^n\text{LO}} = \sigma_{\text{inclusive}}^{\text{N}^n\text{LO}}$$

The N^nLO cross-section is defined such that upon integration over all events the inclusive cross-section at the same order is exactly reproduced

$$\sigma^{(1)} = \Delta_{\text{prod}}^{(1,0)} + \Delta_{\text{dec}}^{(0,1)} + \Delta_{\text{exp}}^{(0,1)} \quad \sigma^{(2)} = \Delta_{\text{prod}}^{(2,0)} + \Delta_{\text{dec}}^{(1,1)} + \Delta_{\text{dec}}^{(0,2)} + \Delta_{\text{exp}}^{(1,1)} + \Delta_{\text{exp}}^{(0,2)}$$

$$d\Gamma_{H \rightarrow b\bar{b}} = d\Gamma^{(0)} + d\Gamma^{(1)} + d\Gamma^{(2)} + \dots$$

$$\Delta_{\text{prod}}^{(i,0)} = \frac{\text{Br}_{H \rightarrow b\bar{b}}}{\Gamma^{\text{LO}}} \int d\sigma_{\text{WBF}}^{(i)} d\Gamma^{(0)}$$

$$\Delta_{\text{dec}}^{(i,j)} = \frac{\text{Br}_{H \rightarrow b\bar{b}}}{\Gamma^{\text{N}^j\text{LO}}} \int d\sigma_{\text{WBF}}^{(i)} d\Gamma^{(j)}$$

$$\Delta_{\text{exp}}^{(i,1)} = -\frac{\text{Br}_{H \rightarrow b\bar{b}} \Gamma^{(1)}}{\Gamma^{\text{LO}} \Gamma^{\text{NLO}}} \int d\sigma_{\text{WBF}}^{(i)} d\Gamma^{(0)}$$

$$\Delta_{\text{exp}}^{(0,2)} = -\frac{\text{Br}_{H \rightarrow b\bar{b}} \Gamma^{(2)}}{\Gamma^{\text{NLO}} \Gamma^{\text{NNLO}}} \int d\sigma_{\text{WBF}}^{(0)} d\Gamma^{\text{NLO}}$$

We split perturbative corrections into *production*, *decay*, and *expansion* corrections


Sources of corrections

$$d\sigma = \underbrace{\text{Br}_{H \rightarrow b\bar{b}}}_{\text{constant}} \underbrace{d\sigma_{\text{WBF}}}_{\text{blue box}} \underbrace{\frac{d\Gamma_{H \rightarrow b\bar{b}}}{\Gamma_{H \rightarrow b\bar{b}}}}_{\text{orange box}}$$


The diagram shows the decomposition of the differential cross-section $d\sigma$ into three parts: a constant branching ratio $\text{Br}_{H \rightarrow b\bar{b}}$, a WBF production term $d\sigma_{\text{WBF}}$ (highlighted in a blue box), and a decay width ratio $\frac{d\Gamma_{H \rightarrow b\bar{b}}}{\Gamma_{H \rightarrow b\bar{b}}}$ (highlighted in an orange box). Dotted arrows connect these components to their respective correction terms on the right. The production term is corrected by $\Delta_{\text{prod}}^{(1,0)}$ and $\Delta_{\text{prod}}^{(2,0)}$. The decay width ratio is corrected by $\Delta_{\text{dec}}^{(0,1)}$, $\Delta_{\text{dec}}^{(0,2)}$, $\Delta_{\text{dec}}^{(1,1)}$, $\Delta_{\text{exp}}^{(0,1)}$, $\Delta_{\text{exp}}^{(0,2)}$, and $\Delta_{\text{exp}}^{(1,1)}$. The total LO cross-section is $\sigma_{\text{LO}} = 75.6 \text{ fb}$.

$\Delta_{\text{prod}}^{(1,0)} = -4.9 \text{ fb}$	$\sigma_{\text{LO}} = 75.6 \text{ fb}$
$\Delta_{\text{prod}}^{(2,0)} = -1.5 \text{ fb}$	
$\Delta_{\text{dec}}^{(0,1)} = -5.3 \text{ fb}$	
$\Delta_{\text{dec}}^{(0,2)} = -5.0 \text{ fb}$	
$\Delta_{\text{dec}}^{(1,1)} = +0.4 \text{ fb}$	
$\Delta_{\text{exp}}^{(0,1)} = -13.0 \text{ fb}$	
$\Delta_{\text{exp}}^{(0,2)} = -2.5 \text{ fb}$	
$\Delta_{\text{exp}}^{(1,1)} = +0.8 \text{ fb}$	

This large effect is a sum of corrections to the Higgs production in WBF (-8%), to $H \rightarrow b\bar{b}$ decay (-14%), and the positive corrections to the total $H \rightarrow b\bar{b}$ width $\Gamma_{H \rightarrow b\bar{b}}$ (-19%)

$$S_i = \lim_{E_i \rightarrow 0} S_2$$


$$S_2 = L_S(p_2) \otimes S_1$$

$$C_{ij} = \lim_{\theta_{ij} \rightarrow 0} C_{12}$$


$$C_{12} = L_C(p_2) \otimes S_1$$

Amplitudes with **soft** and/or **collinear** emissions factorize into amplitudes of lower multiplicity and some universal limit factors.

Nested soft-collinear subtraction scheme

$$\int d^d p_2 \left| \text{diagram} \right|^2 = \int d^4 p_2 (1 - S_2)(1 - C_{12}) \left| \text{diagram} \right|^2 \Bigg] \text{finite}$$

$$\left[\begin{aligned}
 &+ \int d^d p_2 (1 - S_2) C_{12} \left| \text{diagram} \right|^2 \\
 &+ \int d^d p_2 S_2 \left| \text{diagram} \right|^2
 \end{aligned} \right]$$

factorize and integrate analytically

$$\left[\begin{aligned}
 &= \text{finite part} + \frac{\#}{\epsilon_{\text{IR}}^n} \left| \text{diagram} \right|^2 \\
 &= \text{finite part} - \frac{\#}{\epsilon_{\text{IR}}^n} \left| \text{diagram} \right|^2
 \end{aligned} \right] \text{infrared divergences cancel}$$

Catani's formula

We use nested soft-collinear subtraction scheme to cancel infrared divergences between real and virtual corrections.