

# Associated production of Higgses in 2HDMs

S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION

*Guglielmo Coloretti*

*University of Zurich and Paul Scherrer Institut*

*6.11.2024*

# Outline


---

1. (Asymmetric-) associated production of Higgses at the LHC
2. Drell-Yan production of Higgses in 2HDMs
3. The flavored aligned 2HDM (A-2HDM)
4. Correlating  $\text{Br}[A \rightarrow \gamma\gamma]$  with EDMs
5. The 152 GeV and 95 GeV di-photon excesses

# Scalar sector


$$\mathcal{L} = +\mu^2 |\Phi|^2 - \lambda |\Phi|^4$$

- **Minimality of the scalar sector of the SM not guaranteed theoretically**
- Scalar extensions common to multiple NP models
- Run-3 data (on their way) and HL-LHC stage will provide a unique opportunity to inspect non-trivial signatures




# (Asymmetric-) associated production

- Provides a yet unexplored window on new physics [ATLAS and CMS review]  
[Talk by Maggie]  
[Talk by Davide]
- Additional particles required in the signal regions (on top of the decays of the NP candidate)
- **Reduced SM background and enhanced NP sensitivity**



# (Asymmetric-) associated production

- Provides a yet unexplored window on new physics [ATLAS and CMS review]  
[Talk by Maggie]  
[Talk by Davide]
- Additional particles required in the signal regions (on top of the decays of the NP candidate)
- **Reduced SM background and enhanced NP sensitivity**




# (Asymmetric-) associated production


[ATLAS and CMS review]

- Provides a yet unexplored window on new physics
- Additional particles required in the signal regions (on top of the decays of the NP candidate)
- **Reduced SM background and enhanced NP sensitivity**

[Talk by Maggie]  
[Talk by Davide]



Gluon-fusion initiated via a heavy scalar



Drell-Yan production

**Future lepton colliders!**

# Drell-Yan production

---

**New Higgses mostly produced via Drell-Yan  
at the LHC must have specific properties**

# Drell-Yan production

---

**New Higgses mostly produced via Drell-Yan  
at the LHC must have specific properties**

Transform non-trivially  
under  $SU(2)_L$

Gauge interaction with SM  
fields



# Drell-Yan production

---

**New Higgses mostly produced via Drell-Yan at the LHC must have specific properties**

Transform non-trivially under  $SU(2)_L$

No direct (or tiny) Yukawa couplings

Gauge interaction with SM fields

Suppressed gluon-fusion production

# Drell-Yan production

---

**New Higgses mostly produced via Drell-Yan at the LHC must have specific properties**

Transform non-trivially under  $SU(2)_L$

No direct (or tiny) Yukawa couplings

Have small vacuum expectation value

Gauge interaction with SM fields

Suppressed gluon-fusion production

Suppressed VBF and VH production

# Drell-Yan production

**New Higgses mostly produced via Drell-Yan at the LHC must have specific properties**

Transform non-trivially under  $SU(2)_L$

No direct (or tiny) Yukawa couplings

Have small vacuum expectation value

Small mixing with the SM Higgs boson

Gauge interaction with SM fields

Suppressed gluon-fusion production

Suppressed VBF and VH production

Alignment limit

# 2HDMs and Drell-Yan


---

2HDMs strongly motivated:  
MSSM, GUT, PQ (strong CP), custodial symmetry

# 2HDMs and Drell-Yan

2HDMs strongly motivated:  
MSSM, GUT, PQ (strong CP), custodial symmetry

$Y_{\Phi_{NP}} \ll 1$   
 $\langle \Phi_{SM} \rangle \approx v, \langle \Phi_{NP} \rangle \ll 1$   
Small mixing



Suppressed GF  
Suppressed VBF/VH  
Higgs/Flavor bounds


# 2HDMs and Drell-Yan

2HDMs strongly motivated:  
MSSM, GUT, PQ (strong CP), custodial symmetry

$$Y_{\Phi_{NP}} \ll 1$$

$$\langle \Phi_{SM} \rangle \approx v, \langle \Phi_{NP} \rangle \ll 1$$

Small mixing




Suppressed GF

Suppressed VBF/VH


Higgs/Flavor bounds

Drell-Yan is the main  
production mechanism



# 2HDMs and Drell-Yan

- Difficult to obtain sizable  $\text{Br}[H/A \rightarrow \gamma\gamma]$  with  $Z_2$  symmetries for masses around the EW scale




Alignment limit ( $\beta - \alpha \approx 0$ )

	Type-I	Type-II
$\kappa_u$	$-1 / \tan(\beta)$	$-1 / \tan(\beta)$
$\kappa_d$	$1 / \tan(\beta)$	$\tan(\beta)$

# 2HDMs and Drell-Yan

- Difficult to obtain sizable  $\text{Br}[H/A \rightarrow \gamma\gamma]$  with  $Z_2$  symmetries for masses around the EW scale



Alignment limit ( $\beta - \alpha \approx 0$ )


	Type-I	Type-II
$\kappa_u$	$-1/\tan(\beta)$	$-1/\tan(\beta)$
$\kappa_d$	$1/\tan(\beta)$	$\tan(\beta)$

$$\kappa_u \gg 1 \Rightarrow \kappa_d \gg 1$$



# 2HDMs and Drell-Yan

- Difficult to obtain sizable  $\text{Br}[H/A \rightarrow \gamma\gamma]$  with  $Z_2$  symmetries for masses around the EW scale



Alignment limit  $(\beta - \alpha) \approx 0$

	Type-I	Type-II
$\kappa_u$	$-1/\tan(\beta)$	$-1/\tan(\beta)$
$\kappa_d$	$1/\tan(\beta)$	$\tan(\beta)$

$$\kappa_u \gg 1 \Rightarrow \kappa_d \gg 1$$


$$\kappa_u \gg 1, \kappa_d \ll 1$$

Flavor bounds:  $m_{H^\pm} > 600 \text{ GeV}$

[J. Haller, A. Hoecker et al.]

# 2HDMs and Drell-Yan

- Difficult to obtain sizable  $\text{Br}[H/A \rightarrow \gamma\gamma]$  with  $Z_2$  symmetries for masses around the EW scale



Alignment limit  $(\beta - \alpha) \approx 0$

	Type-I	Type-II
$\kappa_u$	$-1/\tan(\beta)$	$-1/\tan(\beta)$
$\kappa_d$	$1/\tan(\beta)$	$\tan(\beta)$

$$\kappa_u \gg 1 \Rightarrow \kappa_d \gg 1$$

$$\kappa_u \gg 1, \kappa_d \ll 1$$

Flavor bounds:  $m_{H^\pm} > 600 \text{ GeV}$

[J. Haller, A. Hoecker et al.]

- Composite Higgs? **Relaxing  $Z_2$  symmetries?**

# A-2HDM

[P. Tuzon, A. Pich]

- Yukawa's of  $\phi_1 \propto$  Yukawa's of  $\phi_2 \implies$  **NO FCNC (tree)**

$$\mathcal{L}_Y = -\bar{Q}_L Y_d (\phi_1 + \zeta_d \phi_2) d_R - \bar{Q}_L Y_u (\tilde{\phi}_1 + \zeta_u^* \tilde{\phi}_2) u_R \\ - \bar{L}_L Y_\ell (\phi_1 + \zeta_\ell \phi_2) \ell_R + \text{h.c.}$$

$\zeta_u \ll 1 \rightarrow$  **suppressed GF (small mixing)**

# A-2HDM

[P. Tuzon, A. Pich]

- Yukawa's of  $\phi_1 \propto$  Yukawa's of  $\phi_2 \Rightarrow$  **NO FCNC (tree)**

$$\mathcal{L}_Y = -\bar{Q}_L Y_d (\phi_1 + \zeta_d \phi_2) d_R - \bar{Q}_L Y_u (\tilde{\phi}_1 + \zeta_u^* \tilde{\phi}_2) u_R \\ - \bar{L}_L Y_\ell (\phi_1 + \zeta_\ell \phi_2) \ell_R + \text{h.c.}$$

$\zeta_u \ll 1 \rightarrow$  **suppressed GF**  
**(small mixing)**

- No  $Z_2$  symmetry imposed  $\Rightarrow$   **$\lambda_6$  and  $\lambda_7$  terms allowed**

$$\mathcal{V} = \mathcal{V}_{Z_2} + [\lambda_6(\phi_1^\dagger \phi_1) + \lambda_7(\phi_2^\dagger \phi_2)] \phi_1^\dagger \phi_2 + \text{h.c.}$$

# A-2HDM

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]


- Yukawa's of  $\phi_1 \propto$  Yukawa's of  $\phi_2 \Rightarrow$  **NO FCNC (tree)**

$$\mathcal{L}_Y = -\bar{Q}_L Y_d (\phi_1 + \zeta_d \phi_2) d_R - \bar{Q}_L Y_u (\tilde{\phi}_1 + \zeta_u^* \tilde{\phi}_2) u_R - \bar{L}_L Y_\ell (\phi_1 + \zeta_\ell \phi_2) \ell_R + \text{h.c.}$$

$\zeta_u \ll 1 \rightarrow$  **suppressed GF (small mixing)**

- No  $Z_2$  symmetry imposed  $\Rightarrow$   **$\lambda_6$  and  $\lambda_7$  terms allowed**

$$\mathcal{V} = \mathcal{V}_{Z_2} + [\lambda_6 (\phi_1^\dagger \phi_1) + \lambda_7 (\phi_2^\dagger \phi_2)] \phi_1^\dagger \phi_2 + \text{h.c.}$$



**Sizable Br[ $H/A \rightarrow \gamma\gamma$ ] through  $H^\pm$  loop**

# A-2HDM: CP-violation

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]

- CP-violation of the model (Baryogenesis?)

[[K. Enomoto](#), [S. Kanemura](#), [Y. Mura](#)]

Yukawa sector:  $\zeta_u, \zeta_d, \zeta_\ell$

Scalar potential:  $\lambda_5, \lambda_6, \lambda_7$

# A-2HDM: CP-violation

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]

- CP-violation of the model (Baryogenesis?)

[[K. Enomoto](#), [S. Kanemura](#), [Y. Mura](#)]

Yukawa sector:  $\zeta_u, \zeta_d, \zeta_\ell$

Scalar potential:  $\lambda_5, \lambda_6, \lambda_7$

- $\Re[\lambda_6]/\Im[\lambda_6]$  drives  $\text{Br}[H/A \rightarrow \gamma\gamma]$

# A-2HDM: CP-violation

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]

- CP-violation of the model (Baryogenesis?)


[K. Enomoto, S. Kanemura, Y. Mura]

Yukawa sector:  $\zeta_u, \zeta_d, \zeta_\ell$

Scalar potential:  $\lambda_5, \lambda_6, \lambda_7$

- $\Re[\lambda_6]/\Im[\lambda_6]$  drives  $\text{Br}[H/A \rightarrow \gamma\gamma]$
- **Correlating  $\text{Br}[A \rightarrow \gamma\gamma]$  with EDMs?**

$$i d_f \bar{f} \sigma^{\mu\nu} q_\mu \gamma_5 f \subset$$





# Higgs basis

[S. Davidson, H. Haber]

- Higgs-flavor symmetry:  $U(2)_{HF}^{ab} \Phi_a = \Phi_b$
- $\langle \Phi_1 \rangle = v, \langle \Phi_2 \rangle = 0$  ;  $\lambda_{1,2,3,4,5} \iff Z_{1,2,3,4,5}$   
**Suppressed VBF / VH (small mixing)**  $\lambda_{6,7} \iff Z_{7,6}$

# Higgs basis

[S. Davidson, H. Haber]

- Higgs-flavor symmetry:  $U(2)_{HF}^{ab} \Phi_a = \Phi_b$
- $\langle \Phi_1 \rangle = v, \langle \Phi_2 \rangle = 0$  ;  $\lambda_{1,2,3,4,5} \iff Z_{1,2,3,4,5}$   
Suppressed VBF / VH (small mixing)  $\lambda_{6,7} \iff Z_{7,6}$

- **Explicit treatment of CP-violation**

$$\Im(Z_5^* Z_6^2) = \Im(Z_5^* Z_7^2) = \Im(Z_6^* Z_7) = 0$$

# Higgs basis

[S. Davidson, H. Haber]

- Higgs-flavor symmetry:  $U(2)_{HF}^{ab} \Phi_a = \Phi_b$
- $\langle \Phi_1 \rangle = v, \langle \Phi_2 \rangle = 0$  ;  $\lambda_{1,2,3,4,5} \iff Z_{1,2,3,4,5}$   
 $\lambda_{6,7} \iff Z_{7,6}$   
 Suppressed VBF / VH (small mixing)

- **Explicit treatment of CP-violation**

$$\Im(Z_5^* Z_6^2) = \Im(Z_5^* Z_7^2) = \Im(Z_6^* Z_7) = 0$$

$$\mathcal{M}_{hHA}^2 = v^2 \begin{pmatrix} Z_1 & \Re(Z_6) & -\Im(Z_6) \\ \Re(Z_6) & \frac{1}{2}[Z_{34} + \Re(Z_5)] & -\frac{1}{2}\Im(Z_5) \\ -\Im(Z_6) & -\frac{1}{2}\Im(Z_5) & \frac{1}{2}[Z_{34} - \Re(Z_5)] \end{pmatrix}$$

# Higgs basis

[S. Davidson, H. Haber]

- Higgs-flavor symmetry:  $U(2)_{HF}^{ab} \Phi_a = \Phi_b$
- $\langle \Phi_1 \rangle = v, \langle \Phi_2 \rangle = 0$  ;  $\lambda_{1,2,3,4,5} \iff Z_{1,2,3,4,5}$   
 $\lambda_{6,7} \iff Z_{7,6}$   
 Suppressed VBF / VH (small mixing)

- **Explicit treatment of CP-violation**

$$\Im(Z_5^* Z_6^2) = \Im(Z_5^* Z_7^2) = \Im(Z_6^* Z_7) = 0$$

SM-Higgs mass

$$\mathcal{M}_{hHA}^2 = v^2 \begin{pmatrix} Z_1 & \Re(Z_6) & -\Im(Z_6) \\ \Re(Z_6) & \frac{1}{2}[Z_{34} + \Re(Z_5)] & -\frac{1}{2}\Im(Z_5) \\ -\Im(Z_6) & -\frac{1}{2}\Im(Z_5) & \frac{1}{2}[Z_{34} - \Re(Z_5)] \end{pmatrix}$$

# Higgs basis

[S. Davidson, H. Haber]

- Higgs-flavor symmetry:  $U(2)_{HF}^{ab} \Phi_a = \Phi_b$
- $\langle \Phi_1 \rangle = v, \langle \Phi_2 \rangle = 0$  ;  $\lambda_{1,2,3,4,5} \iff Z_{1,2,3,4,5}$   
 $\lambda_{6,7} \iff Z_{7,6}$   
 Suppressed VBF / VH (small mixing)

- **Explicit treatment of CP-violation**

$$\Im(Z_5^* Z_6^2) = \Im(Z_5^* Z_7^2) = \Im(Z_6^* Z_7) = 0$$

SM-Higgs mass

Alignment

$$\mathcal{M}_{hHA}^2 = v^2 \begin{pmatrix} Z_1 & \Re(Z_6) & -\Im(Z_6) \\ \Re(Z_6) & \frac{1}{2}[Z_{34} + \Re(Z_5)] & -\frac{1}{2}\Im(Z_5) \\ -\Im(Z_6) & -\frac{1}{2}\Im(Z_5) & \frac{1}{2}[Z_{34} - \Re(Z_5)] \end{pmatrix}$$

# Higgs basis

[S. Davidson, H. Haber]

- Higgs-flavor symmetry:  $U(2)_{HF}^{ab} \Phi_a = \Phi_b$
- $\langle \Phi_1 \rangle = v, \langle \Phi_2 \rangle = 0$  ;  $\lambda_{1,2,3,4,5} \iff Z_{1,2,3,4,5}$   
 $\lambda_{6,7} \iff Z_{7,6}$   
 Suppressed VBF / VH (small mixing)

- **Explicit treatment of CP-violation**

$$\Im(Z_5^* Z_6^2) = \Im(Z_5^* Z_7^2) = \Im(Z_6^* Z_7) = 0$$

SM-Higgs mass

Alignment

H/A mixing

$$\mathcal{M}_{hHA}^2 = v^2 \begin{pmatrix} Z_1 & \Re(Z_6) & -\Im(Z_6) \\ \Re(Z_6) & \frac{1}{2}[Z_{34} + \Re(Z_5)] & -\frac{1}{2}\Im(Z_5) \\ -\Im(Z_6) & -\frac{1}{2}\Im(Z_5) & \frac{1}{2}[Z_{34} - \Re(Z_5)] \end{pmatrix}$$

# Higgs basis

[S. Davidson, H. Haber]

- Higgs-flavor symmetry:  $U(2)_{HF}^{ab} \Phi_a = \Phi_b$
- $\langle \Phi_1 \rangle = v, \langle \Phi_2 \rangle = 0$  ;  $\lambda_{1,2,3,4,5} \iff Z_{1,2,3,4,5}$   
 $\lambda_{6,7} \iff Z_{7,6}$   
 Suppressed VBF / VH (small mixing)

- **Explicit treatment of CP-violation**

$$\Im(Z_5^* Z_6^2) = \Im(Z_5^* Z_7^2) = \Im(Z_6^* Z_7) = 0$$

SM-Higgs mass

Alignment

H/A mixing

$$\mathcal{M}_{hHA}^2 = v^2 \begin{pmatrix} Z_1 & \Re(Z_6) & -\Im(Z_6) \\ \Re(Z_6) & \frac{1}{2}[Z_{34} + \Re(Z_5)] & -\frac{1}{2}\Im(Z_5) \\ -\Im(Z_6) & -\frac{1}{2}\Im(Z_5) & \frac{1}{2}[Z_{34} - \Re(Z_5)] \end{pmatrix}$$

**$Z_7$  independent of mixing angles**

# Higgs basis

[S. Davidson, H. Haber]

- Higgs-flavor symmetry:  $U(2)_{HF}^{ab} \Phi_a = \Phi_b$
- $\langle \Phi_1 \rangle = v, \langle \Phi_2 \rangle = 0$  ;  $\lambda_{1,2,3,4,5} \iff Z_{1,2,3,4,5}$   
 $\lambda_{6,7} \iff Z_{7,6}$   
 Suppressed VBF / VH (small mixing)

- **Explicit treatment of CP-violation**

$$\Im(Z_5^* Z_6^2) = \Im(Z_5^* Z_7^2) = \Im(Z_6^* Z_7) = 0$$

SM-Higgs mass

Alignment

H/A mixing

$$\mathcal{M}_{hHA}^2 = v^2 \begin{pmatrix} Z_1 & \Re(Z_6) & -\Im(Z_6) \\ \Re(Z_6) & \frac{1}{2}[Z_{34} + \Re(Z_5)] & -\frac{1}{2}\Im(Z_5) \\ -\Im(Z_6) & -\frac{1}{2}\Im(Z_5) & \frac{1}{2}[Z_{34} - \Re(Z_5)] \end{pmatrix}$$

Br[H/A → γγ]  
sizable!

**Z<sub>7</sub> independent of mixing angles**




- Electron gives stringent bounds ( $10^{-30}$  e cm $^{-1}$ )
- Projections for neutron/proton ( $10^{-28}/10^{-29}$  e cm $^{-1}$ )

$$d_n = + (0.78 \pm 0.03)d_d - (0.20 \pm 0.01)d_u - e(1.1 \pm 0.55)\tilde{d}_d - e(0.55 \pm 0.28)\tilde{d}_u + e(50 \pm 40) \text{ MeV } d_G$$

- RGE improved chromo-magnetic contributions

- Analytic results

[W. Altmannshofer, S. Gori, N. Hamer, H. Patel]



- Electron gives stringent bounds ( $10^{-30}$  e cm $^{-1}$ )
- Projections for neutron/proton ( $10^{-28}/10^{-29}$  e cm $^{-1}$ )


$$d_n = + (0.78 \pm 0.03)d_d - (0.20 \pm 0.01)d_u - e(1.1 \pm 0.55)\tilde{d}_d - e(0.55 \pm 0.28)\tilde{d}_u + e(50 \pm 40) \text{ MeV } d_G$$

- RGE improved chromo-magnetic contributions

- Analytic results


[W. Altmannshofer, S. Gori, N. Hamer, H. Patel]

- $H^\pm$  contribution  $\propto \Im[\zeta_u^* \zeta_d]$




# Hints for New Physics at 152 GeV

No significant excess in **inclusive  $\gamma\gamma$**  searches




Interesting excesses in  $\gamma\gamma + X$   
 ( $X$  represents additional particles in the signal regions)



**Associated production mechanism**


# A-2HDM: $A_{152} \rightarrow \gamma\gamma$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]



# A-2HDM: $A_{152} \rightarrow \gamma\gamma$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]




$m$	$h$	$H$	$A$	$H^\pm$
[GeV]	125	200	152	130

[ATLAS]

# A-2HDM: $A_{152} \rightarrow \gamma\gamma$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]



$m$	$h$	$H$	$A$	$H^\pm$
[GeV]	125	200	152	130


[ATLAS]

- $\zeta_u = \zeta_d = \zeta_\ell = \zeta_f \in \mathbb{R}$

# A-2HDM: $A_{152} \rightarrow \gamma\gamma$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]

[ATLAS]




$m$	$h$	$H$	$A$	$H^\pm$
[GeV]	125	200	152	130

- $\zeta_u = \zeta_d = \zeta_\ell = \zeta_f \in \mathbb{R}$
- $\theta_{12} = 10^{-3}$   
 $\theta_{13} = \theta_{23} = 10^{-2}$

# A-2HDM: $A_{152} \rightarrow \gamma\gamma$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]



$m$	$h$	$H$	$A$	$H^\pm$
[GeV]	125	200	152	130


[ATLAS]

- $\zeta_u = \zeta_d = \zeta_\ell = \zeta_f \in \mathbb{R}$
- $\theta_{12} = 10^{-3}$   
 $\theta_{13} = \theta_{23} = 10^{-2}$
- $Z_2 = -Z_3 = 0.2$
- $\Re[Z_7] = 0.1$
- HiggsTools, perturbativity, vacuum stability



# A-2HDM: $A_{152} \rightarrow \gamma\gamma$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]



$m$	$h$	$H$	$A$	$H^\pm$
[GeV]	125	200	152	130


[ATLAS]

- $\zeta_u = \zeta_d = \zeta_\ell = \zeta_f \in \mathbb{R}$
- $\theta_{12} = 10^{-3}$   
 $\theta_{13} = \theta_{23} = 10^{-2}$
- $Z_2 = -Z_3 = 0.2$
- $\Re[Z_7] = 0.1$
- HiggsTools, perturbativity, vacuum stability

# Hints at 95/98 GeV

## Inclusive searches

- LEP:  $Z + b\bar{b}$  ( $2.3 \sigma$ )
- ATLAS:  $\gamma\gamma$  ( $1.7 \sigma$ )
- CMS:  $\gamma\gamma$  ( $2.9 \sigma$ )
- CMS:  $\bar{\tau}\tau$  ( $2.4 \sigma$ )  
(but not seen by ATLAS)






[U. Haisch, A. Malinauskas]

[T. Biekotter, S. Heinemeyer, C. Munoz]


[T. Biekotter, S. Heinemeyer, G. Weiglein et al.]

[T. Biekotter, S. Heinemeyer, G. Weiglein]




# A-2HDM: $A_{95} \rightarrow \gamma\gamma, H_{98} \rightarrow b\bar{b}$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]



# A-2HDM: $A_{95} \rightarrow \gamma\gamma, H_{98} \rightarrow b\bar{b}$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]




$m$	$h$	$H$	$A$	$H^\pm$
[GeV]	125	98	95	130

[ATLAS]

# A-2HDM: $A_{95} \rightarrow \gamma\gamma, H_{98} \rightarrow b\bar{b}$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]




$m$	$h$	$H$	$A$	$H^\pm$
[GeV]	125	98	95	130

[ATLAS]

- Gluon fusion contribution:  
 $\zeta_d = \zeta_\ell = 0, \zeta_u \in \mathbb{R}$

# A-2HDM: $A_{95} \rightarrow \gamma\gamma, H_{98} \rightarrow b\bar{b}$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]




$m$	$h$	$H$	$A$	$H^\pm$
[GeV]	125	98	95	130

[ATLAS]

- Gluon fusion contribution:  
 $\zeta_d = \zeta_\ell = 0, \zeta_u \in \mathbb{R}$
- $\theta_{12} = 0.25$  ( $\mu_{b\bar{b},98}^{\text{LEP}}$ )  
 $\theta_{13} = 10^{-2}, \theta_{23} = 3 \times 10^{-2}$

# A-2HDM: $A_{95} \rightarrow \gamma\gamma, H_{98} \rightarrow b\bar{b}$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]




$m$	$h$	$H$	$A$	$H^\pm$
[GeV]	125	98	95	130

[ATLAS]

- Gluon fusion contribution:  $\zeta_d = \zeta_\ell = 0, \zeta_u \in \mathbb{R}$
- $\theta_{12} = 0.25$  ( $\mu_{b\bar{b},98}^{\text{LEP}}$ )  
 $\theta_{13} = 10^{-2}, \theta_{23} = 3 \times 10^{-2}$
- $Z_2 = -Z_3 = 0.2$
- $\Re[Z_7] = 0.1$
- HiggsTools, perturbativity, vacuum stability

# A-2HDM: $A_{95} \rightarrow \gamma\gamma, H_{98} \rightarrow b\bar{b}$

[S. Banik, GC, A. Crivellin, H. Haber - IN PREPARATION]



$m$	$h$	$H$	$A$	$H^\pm$
[GeV]	125	98	95	130

[ATLAS]

- Gluon fusion contribution:  $\zeta_d = \zeta_\ell = 0, \zeta_u \in \mathbb{R}$
- $\theta_{12} = 0.25$  ( $\mu_{b\bar{b},98}^{\text{LEP}}$ )  
 $\theta_{13} = 10^{-2}, \theta_{23} = 3 \times 10^{-2}$
- $Z_2 = -Z_3 = 0.2$
- $\Re[Z_7] = 0.1$
- HiggsTools, perturbativity, vacuum stability



# Conclusions and Outlook

---

- Asymmetric associated production of scalars is a prominent signature to look for NP at the LHC
- A-2HDM offers sizable  $\text{Br}[H/A \rightarrow \gamma\gamma]$  which can be correlated to CP-violating effects such as EDMs
- A-2HDM provides an independent explanation of the diphoton excesses at 95 GeV and at 152 GeV
- Outlook: correlate  $\text{Br}[H/A \rightarrow \gamma\gamma]$  to Baryogenesis, SFOPT and the effects in the trilinear SM-Higgs coupling

# Conclusions and Outlook

---

- Asymmetric associated production of scalars is a prominent signature to look for NP at the LHC
- A-2HDM offers sizable  $\text{Br}[H/A \rightarrow \gamma\gamma]$  which can be correlated to CP-violating effects such as EDMs
- A-2HDM provides an independent explanation of the diphoton excesses at 95 GeV and at 152 GeV
- Outlook: correlate  $\text{Br}[H/A \rightarrow \gamma\gamma]$  to Baryogenesis, SFOPT and the effects in the trilinear SM-Higgs coupling

**THANK YOU FOR THE ATTENTION!**





---

# BACK UP SLIDES

---




# 2HDMs: flavor bounds

[J. Haller, A. Hoecker et al.]




# A-2HDM: flavor bounds

[A. Karan, V. Miralles, A. Pich]



# 2HDMs: EW precision

[J. Haller, A. Hoecker et al.]






# A-2HDM: EW precision

[A. Karan, V. Miralles, A. Pich]



ST PDG- $M_W$  95%    ST CDF- $M_W$  95%




# FCC-ee prospects

Courtesy of Rebeca Gonzalez Suarez

- Scalars produced in associated production via DY are a prominent candidate for FCC-ee

## FCC-ee

- 16 years, 4 IPs
- Flexibility in the run scenario: in order and operation periods.
  - Additional runs, e.g. 125GeV possible
- Stringent experimental requirements



FCC feasibility Mid-term report - Deliverable #8, physics and Experiments

integrated luminosity per year summed over 4 IPs corresponding to 185 days of physics per year and 75% efficiency

Working point	Z, years 1-2	Z, later	WW, years 1-2	WW, later	ZH	$t\bar{t}$
$\sqrt{s}$ (GeV)	88, 91, 94		157, 163		240	340-350
Lumi/IP ( $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ )	70	140	10	20	5.0	0.75
Lumi/year ( $\text{ab}^{-1}$ )	34	68	4.8	9.6	2.4	0.36
Run time (year)	2	2	2	-	3	1
Number of events	$6 \times 10^{12}$ Z		$2.4 \times 10^8$ WW		$1.45 \times 10^6$ ZH + 45k WW $\rightarrow$ H	$1.9 \times 10^6$ $t\bar{t}$ +330k ZH +80k WW $\rightarrow$ H

all the data of LEP1 in minutes




# FCC-ee prospects

Courtesy of Rebeca Gonzalez Suarez

- Scalars produced in associated production via DY are a prominent candidate for FCC-ee

## FCC-ee

- 16 years, 4 IPs
- Flexibility in the run scenario: in order and operation periods.
  - Additional runs, e.g. 125GeV possible
- Stringent experimental requirements




integrated luminosity per year summed over 4 IPs corresponding to 185 days of physics per year and 75% efficiency

Working point	Z, years 1-2	Z, later	WW, years 1-2	WW, later	ZH	t $\bar{t}$	
$\sqrt{s}$ (GeV)	88, 91, 94		157, 163		240	340-350	365
Lumi/IP ( $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ )	70	140	10	20	5.0	0.75	1.20
Lumi/year ( $\text{ab}^{-1}$ )	34	68	4.8	9.6	2.4	0.36	0.58
Run time (year)	2	2	2	-	3	1	4
Number of events	$6 \times 10^{12}$ Z		$2.4 \times 10^8$ WW		$1.45 \times 10^6$ ZH + 45k WW $\rightarrow$ H	$1.9 \times 10^9$ t $\bar{t}$ +330k ZH +80k WW $\rightarrow$ H	

all the data of LEP1 in minutes

# Real triplet at the FCC-ee

- Only  $Z^*/\gamma^*$  s-channel (suppressed  $\Delta^0\Delta^0$  production)
- Pair production of the charged components



	$SU(2)_L$	$U(1)_Y$
$\Delta$	3	0

Parameters  $\rightarrow \langle \Delta \rangle = v_\Delta, \alpha_\Delta$   
 Fields  $\rightarrow$  neutral  $\Delta^0$ , charged  $\Delta^\pm$

# $6\ell + 2\nu$ at the FCC-ee

- The decay  $\Delta^\pm \rightarrow W^\pm Z$  leads to a  $6\ell(+\text{MET})$  signature


Events expected in the SM model

$$e^+e^- \rightarrow 6\ell(+\text{MET}) \approx 1$$


Events expected in the  $\Delta\text{SM}$  model

$$e^+e^- \rightarrow \Delta^\pm \Delta^\mp \rightarrow 6\ell + \text{MET} \approx 46$$

- Assuming the integrated luminosity at the  $t\bar{t}$  working point (4 years run)
- Log-Likely-hood ratio yields  $\chi^2 \approx 80$
- $\sigma(e^+e^- \rightarrow \Delta^\pm \Delta^\mp)$  could be measured at  $\approx 9\sigma$




# $t\bar{t}$ distributions as a probe for NP



$e^+e^-$  future colliders have a dedicated  $t\bar{t}$  run: is there any new physics testable scenario?

*“No model can describe all measured distributions within their uncertainties.”* [ATLAS]



- Higher order corrections? Toponium?
- New Physics pollution of this SM measurement?


[F. Maltoni, D. Pagani et al.]

[F. Maltoni, C. Severi et al.]


[Bagnschi, Corcella et al.]

# 95 GeV and 152 GeV excesses?

[S. Banik, GC, A. Crivellin, B. Mellado]



- $S_{95}$ : SM singlet mostly decaying to  $b\bar{b}$
- $\Delta^0$ : real Higgs triplet mostly decaying to  $WW$



**Consistent with the 95 GeV  $\gamma\gamma$  signal strength and a mass for  $\Delta^0$  of 152 GeV**