


Anomalous Couplings in Higgs plus Jet Production

Higgs (2024) Uppsala

Benjamin Campillo, Gudrun Heinrich, Matthias Kerner, Lucas Kunz | 06.11.2024

Motivation - Higgs Physics

- Run2 measured couplings of Higgs to heavy fermions
- Run3 and HL-LHC are expected to give even more precise measurements
- Precise theoretical predictions needed
- Want to distinguish BSM effects from higher-order uncertainties in Higgs p_T spectrum



[ATLAS '22]

History

Standard Model

- NLO study with full top mass [Bonciani et al. 2023; Jones, Kerner, and Luisoni 2018; Neumann 2018]
- several NNLO calculations in HTL [Boughezal, Caola, et al. 2013; Chen, Gehrmann, Glover, and Jaquier 2015; Boughezal, Caola, et al. 2015; Boughezal, Focke, et al. 2015; Caola, Melnikov, and Schulze 2015; Chen, Gehrmann, Glover, and Huss 2019; Campbell, Ellis, and Seth 2019]

Effective Field Theory

- LO Study in [Grojean et al. 2014; Schlaffer et al. 2014]
- NLL study [Battaglia et al. 2021]
- LO + RGE running study [Maltoni, Ventura, and Vryonidou 2024]
- 2-loop RGE running [Di Noi, Gröber, and Mandal 2024]
- NLO in HTL [Grazzini et al. 2017]

but not at NLO with full top mass dependence!

Effective Field Theory - Formulae

Linear EFT - Standard Model Effective Field Theory (SMEFT)

- SM symmetries - Poincaré and gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$
- Higgs part of $SU(2)_L$ doublet
- ordering via canonical mass dimension

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \sum_i \frac{c_i^{(5)}}{\Lambda} \mathcal{O}_i^{(5)} + \sum_i \frac{c_i^{(6)}}{\Lambda^2} \mathcal{O}_i^{(6)} + \sum_i \frac{c_i^{(7)}}{\Lambda^3} \mathcal{O}_i^{(7)} + \sum_i \frac{c_i^{(8)}}{\Lambda^4} \mathcal{O}_i^{(8)} + \mathcal{O}\left(\frac{1}{\Lambda^5}\right)$$

Non-linear EFT - Higgs Effective Field Theory (HEFT)

- EW symmetry breaking is realised non-linearly
- Scalar doublet splits into EW singlet h and Goldstone Bosons, which satisfy $SU(2)$
- counting via chiral dimension: $d_\chi(A^\mu, \phi, h, v) = 0$, $d_\chi(\partial^\mu, \bar{\Psi}\Psi, g, Y) = 1$,
- Weinbergs Power counting formula [Weinberg 1979]: $d_\chi = 2 + (D - 2)L + \sum_{k=1}^{\infty} 2(k - 1)V_{2k}$

Effective Operators for Higgs plus Jet

Effective operators up to $d_\chi = 4$:

- Rescaling of top Yukawa coupling (c_t): $O_t = -m_t \frac{\hbar}{v} \bar{t} t$
- Effective Higgs-gluon coupling (c_g): $O_g = \frac{\alpha_s}{8\pi} \frac{\hbar}{v} G_{\mu\nu}^a G^{a,\mu\nu}$

$$\kappa_g = 3/2 c_g : \quad \frac{\sigma_{\text{incl}}(c_t, c_g)}{\sigma_{\text{incl}}^{\text{SM}}} = (c_t + \kappa_g)^2 + \mathcal{O}\left(\frac{\kappa_g}{c_t + \kappa_g} \frac{m_h^2}{4m_t^2}\right)^1$$


$c_t + \kappa_g \approx 1 \pm 10\%$ Measured in inclusive Higgs production in gluon fusion

¹[Grojean et al. 2014]

Calculation


Leading Order

- calculated analytically
- one-loop part based on SM calculation [Baur and Glover 1990]
- cross-checked via GoSam [Cullen et al. 2012; Cullen et al. 2014]



Real Radiation


- calculated with GoSam, ninja [Peraro 2014] and OneLOop [Hameren 2011]
- Implemented into POWHEG-BOX-V2 (does subtraction of IR divergences, PS integration) [Nason 2004; Frixione, Nason, and Oleari 2007; Alioli et al. 2010]



Calculation

Virtual Corrections

- one-loop part done with GoSam , ninja and OneL0op
- two-loop part based on SM calculation
[\[Jones, Kerner, and Luisoni 2018\]](#) using REDUZE [\[Manteuffel and Studerus 2012\]](#) and SECDEC-3 [\[Borowka et al. 2015\]](#)



Calculation

Two-Loop Virtuals

- two-loop form factors computed in [\[Jones, Kerner, and Luisoni 2018\]](#) for 2000 points

- Use: $d\sigma^V \sim 2\Re \left[\left(\mathcal{M}_{c_t}^{2L} + \mathcal{M}_{c_g}^{1L} \right) \cdot \left(\mathcal{M}_{c_t}^{1L} + \mathcal{M}_{c_g}^{0L} \right)^\dagger \right]$
 $= 2\Re \left[\mathcal{M}_{c_t}^{2L} \cdot \left(\mathcal{M}_{c_t}^{1L} + \mathcal{M}_{c_g}^{0L} \right)^\dagger + \mathcal{M}_{c_g}^{1L} \cdot \left(\mathcal{M}_{c_t}^{1L} + \mathcal{M}_{c_g}^{0L} \right)^\dagger \right]$

- $\mathcal{M}_{c_t} = c_t \mathcal{M}_{\text{SM}}$

- \Rightarrow Rescale [\[Jones, Kerner, and Luisoni 2018\]](#) by c_t and add Born c_g form factors

- Use neural network to interpolate $\left[\mathcal{M}_{c_t}^{2L} \cdot \left(\mathcal{M}_{c_t}^{1L} + \mathcal{M}_{c_g}^{0L} \right)^\dagger \right]$

- Calculate $\mathcal{M}_{c_g}^{1L} \cdot \left(\mathcal{M}_{c_t}^{1L} + \mathcal{M}_{c_g}^{0L} \right)^\dagger$ via GoSam and implement into POWHEG-BOX-V2

Results - Cross Section

$p_{T,H}^{\text{cut}}$ [GeV]	$\sigma_{\text{cut}}(c_t = 0.9, c_g = 1/15)$ [fb]		$\sigma_{\text{cut}}(\text{SM})$ [fb]	
	LO	NLO	LO	NLO
50	$(4.5^{+1.8}_{-1.2}) \cdot 10^3$	$(9.1^{+0.1}_{-1.6}) \cdot 10^3$	$(4.5^{+1.8}_{-1.2}) \cdot 10^3$	$(9.1^{+0.0}_{-1.4}) \cdot 10^3$
400	13^{+6}_{-4}	28^{+2}_{-6}	12^{+5}_{-3}	$25^{+1.6}_{-5}$
600	$1.6^{+0.7}_{-0.5}$	$3.3^{+0.3}_{-0.7}$	$1.3^{+0.6}_{-0.4}$	$2.7^{+0.2}_{-0.5}$
800	$0.29^{+0.14}_{-0.09}$	$0.60^{+0.07}_{-0.12}$	$0.21^{+0.10}_{-0.06}$	$0.43^{+0.04}_{-0.09}$


- Scale given by: $\mu_0 = \frac{H_T}{2} = \frac{1}{2} \left(\sqrt{m_H^2 + p_{T,H}^2} + \sum_i |p_{T,i}| \right)$, (sum i over final state partons)
- Difference only exceeds scale uncertainties for highly boosted Higgs bosons, $p_{T,H} > 600\text{GeV}$

Results - Cross Section

$p_{T,H}^{\text{cut}}$ [GeV]	$\sigma_{\text{cut,HEFT}}/\sigma_{\text{cut,SM}}$	
	LO	NLO
50	0.9966 ± 0.0006	1.00 ± 0.04
400	1.119 ± 0.007	1.10 ± 0.01
600	1.250 ± 0.012	1.22 ± 0.01
800	1.410 ± 0.016	1.37 ± 0.01

- Ratio increases for higher $p_{T,H}$ cuts
- Monte-Carlo uncertainties

Results - Transverse Momentum Distribution



- Only for boosted Higgs is BSM outside SM scale uncertainty
- Cannot use HTL!

Results - Fits

- $\sigma = c_t^2 A + 2c_t c_g B + c_g^2 C$, $\sigma_{\text{SM}} = A$

Results - Fits

$$\blacksquare \sigma = c_t^2 A + 2c_t c_g B + c_g^2 C \quad , \quad \sigma_{\text{SM}} = A$$



$p_{T,H}^{\text{cut}}$	A	B	C
50	9.13 ± 0.12	13.1 ± 0.3	18.9 ± 0.5
400	$(2.54 \pm 0.01) \cdot 10^{-2}$	$(5.65 \pm 0.08) \cdot 10^{-2}$	$(13.7 \pm 0.1) \cdot 10^{-2}$
600	$(2.73 \pm 0.01) \cdot 10^{-3}$	$(8.37 \pm 0.11) \cdot 10^{-3}$	$(28.9 \pm 0.2) \cdot 10^{-3}$
800	$(4.33 \pm 0.02) \cdot 10^{-4}$	$(17.5 \pm 0.2) \cdot 10^{-4}$	$(80.6 \pm 0.5) \cdot 10^{-4}$

$$\blacksquare C > B > A$$

- $\blacksquare C/A$ increases with cut; A scales as $1/p_{T,H}^2$, C as $1/p_{T,H}$ [Caola, Forte, et al. 2016]

- \blacksquare Uncertainties = Monte-Carlo + Fit

Results - Heat Maps



- Bounds inspired by [\[Celada et al. 2024\]](#)
- Higgs p_T cut of 400GeV
- $\sigma = c_t^2 A + 2c_t c_g B + c_g^2 C$, $C > B > A$

- Ratios can vary considerably
- Inside grey band: Negative c_g : Lower ratio, higher K-factors
Positive c_g : Higher ratio, lower K-factors

Conclusion and Outlook

Higgs plus jet at NLO QCD


- First study of NLO Higgs plus jet production with HEFT operators and full top mass dependence.
- Studied the total cross section and transverse momentum ($p_{T,H}$) differential distribution
 - Shown that high $p_{T,H}$ regime is sensitive to BSM effects
 - Still a lot hidden in scale uncertainties
- Studied heat maps of ratio to SM
 - Fit of coupling coefficients allows for fast evaluation of cross section
- Code will be available on the POWHEG-BOX-V2

Outlook

- Higher order HEFT operators
- Higher order QCD corrections
- Other SM uncertainties

Thank you for your attention!

Backup - Chromomagnetic Operator



(c)

$$d_\chi = 6$$

non-hermitian ($\bar{L}R$)

6: $\psi^2 XH + \text{h.c. [LG]}$			
$Q_{eW} (\bar{\ell}_p \sigma^{\mu\nu} e_r) \tau^I H W_{\mu\nu}^I$	$Q_{uG} (\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} G_{\mu\nu}^A$	$Q_{dG} (\bar{q}_p \sigma^{\mu\nu} T^A d_r) H G_{\mu\nu}^A$	
$Q_{eB} (\bar{\ell}_p \sigma^{\mu\nu} e_r) H B_{\mu\nu}$	$Q_{uW} (\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} W_{\mu\nu}^I$	$Q_{dW} (\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I H W_{\mu\nu}^I$	
	$Q_{uB} (\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{H} B_{\mu\nu}$	$Q_{dB} (\bar{q}_p \sigma^{\mu\nu} d_r) H B_{\mu\nu}$	

Table taken from [\[Isidori, Wilsch, and Wyler 2024\]](#)

LG = Loop generated

Backup - Form Factors for $g(p_1)g(p_2) \rightarrow g(p_3)h(p_4)$

$$M^{\mu\nu\tau} = F_a T_a^{\mu\nu\tau} + F_b T_b^{\mu\nu\tau} + F_c T_c^{\mu\nu\tau} + F_d T_d^{\mu\nu\tau}$$

$$T_a^{\mu\nu\tau} = (s_{12}g^{\mu\nu} - 2p_2^\mu p_1^\nu)(s_{23}p_1^\tau - s_{13}p_2^\tau) \frac{1}{2s_{13}}$$

$$T_b^{\mu\nu\tau} = (s_{23}g^{\nu\tau} - 2p_3^\nu p_2^\tau)(s_{13}p_2^\mu - s_{12}p_3^\mu) \frac{1}{2s_{12}}$$

$$T_c^{\mu\nu\tau} = (s_{13}g^{\tau\mu} - 2p_1^\tau p_3^\mu)(s_{12}p_3^\nu - s_{23}p_1^\nu) \frac{1}{2s_{23}}$$

$$T_d^{\mu\nu\tau} = [g^{\mu\nu}(s_{23}p_1^\tau - s_{13}p_2^\tau) + g^{\nu\tau}(s_{13}p_2^\mu - s_{12}p_3^\mu) + g^{\tau\mu}(s_{12}p_3^\nu - s_{23}p_1^\nu) + 2p_3^\mu p_1^\nu p_2^\tau - 2p_2^\mu p_3^\nu p_1^\tau] \frac{1}{2}$$

Backup - Form Factors for $g(p_1)g(p_2) \rightarrow g(p_3)h(p_4)$

$$P_{k, \mu\nu\tau} M^{\mu\nu\tau} = F_k$$

$$P_a^{\mu\nu\tau} = \frac{1}{(D-3) s_{23}} \left(-\frac{Ds_{13}}{s_{12}^2 s_{23}} T_a^{\mu\nu\tau} + \frac{D-4}{s_{23}^2} T_b^{\mu\nu\tau} + \frac{D-4}{s_{12} s_{13}} T_c^{\mu\nu\tau} + \frac{D-2}{s_{12} s_{23}} T_d^{\mu\nu\tau} \right)$$

$$P_b^{\mu\nu\tau} = \frac{1}{(D-3) s_{12}} \left(\frac{D-4}{s_{12} s_{23}} T_a^{\mu\nu\tau} - \frac{Ds_{12}}{s_{13} s_{23}^2} T_b^{\mu\nu\tau} + \frac{D-4}{s_{13}^2} T_c^{\mu\nu\tau} + \frac{D-2}{s_{13} s_{23}} T_d^{\mu\nu\tau} \right)$$

$$P_c^{\mu\nu\tau} = \frac{1}{(D-3) s_{13}} \left(\frac{D-4}{s_{12}^2} T_a^{\mu\nu\tau} + \frac{D-4}{s_{13} s_{23}} T_b^{\mu\nu\tau} - \frac{Ds_{23}}{s_{12} s_{13}^2} T_c^{\mu\nu\tau} + \frac{D-2}{s_{12} s_{13}} T_d^{\mu\nu\tau} \right)$$

$$P_d^{\mu\nu\tau} = \frac{D-2}{(D-3) s_{12} s_{13} s_{23}} \left(\frac{s_{13}}{s_{12}} T_a^{\mu\nu\tau} + \frac{s_{12}}{s_{23}} T_b^{\mu\nu\tau} + \frac{s_{23}}{s_{13}} T_c^{\mu\nu\tau} + \frac{D}{D-2} T_d^{\mu\nu\tau} \right)$$

Backup - Form Factors for $q(p_1)\bar{q}(p_2) \rightarrow g(p_3)h(p_4)$

$$M^{\rho} \epsilon_{\rho} = F_1 T_1 + F_2 T_2$$

$$T_1 = \bar{u}(p_1) \not{p}_3 v(p_2) (p_2 \cdot \epsilon_3) - \bar{u}(p_1) \not{\epsilon}_3 v(p_2) (p_2 \cdot p_3)$$

$$T_2 = \bar{u}(p_1) \not{p}_3 v(p_2) (p_1 \cdot \epsilon_3) - \bar{u}(p_1) \not{\epsilon}_3 v(p_2) (p_1 \cdot p_3)$$

$$\sum_{\text{spins}} P_i M^{\rho} \epsilon_{\rho} = F_i$$

$$P_1 = \frac{D-2}{2(D-3)s_{12}s_{13}^2} T_1^{\dagger} - \frac{D-4}{2(D-3)s_{12}s_{13}s_{23}} T_2^{\dagger}$$

$$P_2 = -\frac{D-4}{2(D-3)s_{12}s_{13}s_{23}} T_1^{\dagger} + \frac{D-2}{2(D-3)s_{12}s_{23}^2} T_2^{\dagger}$$

Backup - Scale dependency of couplings

- Why is there no scale dependency in the anomalous couplings?
- \Rightarrow Only at LL and without c_{tg} .

$$\begin{aligned}\partial_t c_t &= -4 \frac{m_t^2}{v^2} a_s c_{tg}, \\ \partial_t c_g &= \frac{m_t^2}{2v^2} c_{tg}, \\ \partial_t c_{tg} &= \frac{7a_s}{6} c_{tg}.\end{aligned}$$

Taken from [\[Battaglia et al. 2021\]](#) where $t = \log(Q^2/\mu^2)$

Backup - HEFT Lagrangian

$$\begin{aligned}
 \mathcal{L}_2 = & - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} - \frac{1}{4} W_{\mu\nu}^a W^{a,\mu\nu} - \frac{1}{4} G_{\mu\nu}^b G^{b,\mu\nu} + \sum_{\Psi=Q_L, L_L, U_R, D_R, E_R} \bar{\Psi} i \not{D} \Psi \\
 & + \frac{v^2}{4} \text{Tr} [D_\mu \Omega^\dagger D^\mu \Omega] (1 + F_\Omega(h)) + \frac{1}{2} (\partial_\mu h)(\partial^\mu h) - V(h) \\
 & - v \left[\bar{Q}_L \left(Y_u + \sum_{n=1}^{\infty} Y_u^{(n)} \left(\frac{h}{v} \right)^n \right) \Omega P_+ Q_R + \bar{Q}_L \left(Y_d + \sum_{n=1}^{\infty} Y_d^{(n)} \left(\frac{h}{v} \right)^n \right) \Omega P_- Q_R \right. \\
 & \quad \left. + \bar{L}_L \left(Y_e + \sum_{n=1}^{\infty} Y_e^{(n)} \left(\frac{h}{v} \right)^n \right) \Omega P_- L_R + h.c. \right]
 \end{aligned}$$

Backup - HEFT Lagrangian

Goldstone Bosons: $\Omega = \exp\left(2i \frac{\phi^a \sigma_a}{v}\right)$ with $D_\mu \Omega = \partial_\mu \Omega + igW_\mu \Omega - ig'B_\mu \Omega \sigma_3$

Right-handed Doublets: $Q_R = \begin{pmatrix} U_R \\ D_R \end{pmatrix}$, $L_R = \begin{pmatrix} 0 \\ E_R \end{pmatrix}$ with $P_+ = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $P_- = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$

Higgs Potential: $F_\Omega(h) = \sum_{n=1}^{\infty} f_{\Omega,n} \left(\frac{h}{v}\right)^n$, $V(h) = v^4 \sum_{n=2}^{\infty} f_{V,n} \left(\frac{h}{v}\right)^n$

Standard Model: $f_{\Omega,1} = 2$, $f_{\Omega,2} = 1$, $f_{V,2} = f_{V,3} = \frac{m_h^2}{2v^2}$, $f_{V,4} = \frac{m_h^2}{8v^2}$, $Y_f^{(1)} = Y_f$

Chiral Dimension: $d_\chi(A^\mu, \phi, h, v, f_{\Omega,n}) = 0$, $d_\chi(\partial^\mu, \bar{\Psi}\Psi, g, Y) = 1$, $d_\chi(f_{V,n}) = 2$