



ALICE-India Collaboration



# Quarkonium polarization in pp and Pb-Pb collisions with ALICE

Dushmanta Sahu  
Indian Institute of Technology Indore, India  
April 29, 2023

# Introduction:




## ➤ Why charmonia?

- Charm and anti-charm quarks produced early in the system's evolution : during the pre-equilibrium phase
- Affected by suppression and regeneration at LHC energies
- $J/\psi$  remains largely undiffused in the hadronic phase of a collision which makes it a better probe to study the deconfined phase
- Charmonium studies in hadronic collisions provide powerful tests of quantum chromodynamics (QCD)
- Charmonium production yield in Pb–Pb and p–Pb collisions can also be affected by the cold nuclear matter (CNM) effect (e.g. Shadowing effect)

## ➤ Polarization in pp collisions:

- Polarization is the measure of how much the spin of a particle is aligned in a given direction
- Gluon's polarization is preserved as the  $c\bar{c}$  pair evolves into a bound state of charmonium
- In two-body decays, the spin-alignment will be reflected in the angular distribution of the decay particles

# Introduction:



- $h = \frac{S \cdot p}{|p|} \rightarrow$  Helicity operator
- Vector ( $J^{PC} = 1^{--}$ ) quarkonia have the same charge-parity as an electron-positron pair and can be produced in electron-positron annihilation via an intermediate photon
- The states originating from this process are polarized, as a consequence of helicity conservation, a general property of QED (QCD) in the relativistic (massless) limit
- For our case, gluon fragmentation dominates the high  $p_T$  region, while Drell-Yan process dominates the low  $p_T$  region


# Introduction:

## $J/\psi$ polarization puzzle ?


- Measurements of polarization parameters from Tevatron, RHIC and LHC show almost no  $J/\psi$  polarization in hadronic collisions
- However, theoretical predictions based on the collinear factorized color singlet production channel at leading order (LO) and next-to-leading order (NLO) suggested substantially non-zero polarization at high  $p_T$
- Conflicting theoretical results from non-relativistic quantum chromodynamics (NRQCD) and Color Singlet Model

## Importance of $\psi(2S)$ polarization study :

- A small prompt  $J/\psi$  polarization can be interpreted as reflecting a mixture of directly produced mesons with those produced in the decays of heavier (P-wave) charmonium states
- $\psi(2S)$  is unaffected by feed-down decays from heavier charmonia



[Phys. Rev. Lett. 108, 172002 (2012)]





[ALICE Collaboration, Phys. Rev. Lett. 108, 082001 (2012)]


# Introduction:

## Importance of $\Upsilon(nS)$ polarization study :

- $b\bar{b}$  system satisfies the non relativistic calculations at high  $p_T$  much better than the  $c\bar{c}$
- Better probe for QCD
- Results from Tevatron show almost no (CDF) or longitudinal polarization for  $\Upsilon(1S)$  (D0)
- At lower energy and  $p_T$ , the E866 experiment has shown yet a different polarization pattern: the  $\Upsilon(2S)$  and  $\Upsilon(3S)$  states have maximal transverse polarization
- Unexpectedly, the  $\Upsilon(1S)$  found to be only weakly polarized



[D $\Phi$  Collaboration, Phys. Rev. Lett. 101, 182004 (2008)]



[NuSea Collaboration, Phys. Rev. Lett. 86, 2529 (2001)]

[CDF Collaboration, Phys. Rev. Lett. 88, 161802 (2002)]


# Introduction:

- The angular distribution in dilepton decay:

$$\frac{d^2 N}{d\cos\theta d\phi} = \frac{3}{4\pi(3 + \lambda_\theta)} (1 + \lambda_\theta \cos^2\theta + \lambda_\phi \sin^2\theta \cos 2\phi + \lambda_{\theta\phi} \sin 2\theta \cos\phi)$$

[P.Faccioli, et. al., Eur. Phys. J. C 69, 657 (2010)]


- $(\lambda_\theta, \lambda_\phi, \lambda_{\theta\phi}) = (1, 0, 0)$   $\longrightarrow$  Transverse polarization
- $(\lambda_\theta, \lambda_\phi, \lambda_{\theta\phi}) = (-1, 0, 0)$   $\longrightarrow$  Longitudinal polarization
- $(\lambda_\theta, \lambda_\phi, \lambda_{\theta\phi}) = (0, 0, 0)$   $\longrightarrow$  Unpolarized state



# Introduction:

## Frames of reference


- The helicity frame uses the  $\psi(2S)$  momentum as the quantization axis
- In the Collins–Soper frame, the quantization axis is chosen to be the bisector of the angle between the two incoming beams in the rest frame of the  $\psi(2S)$  meson
- We can define the frame-invariant variable  $\lambda_{inv}$



$$\lambda_{inv} = \frac{\lambda_{\theta} + 3\lambda_{\phi}}{1 - \lambda_{\phi}}$$

# Quarkonium polarization in Pb–Pb collisions:

- Large non-zero magnetic field in non-central heavy-ion collisions
- Production of vorticity due to large initial angular momentum
- Both the external magnetic field and the initial angular momentum produced in the non-central heavy-ion collisions may influence the quarkonium polarization
- Event Plane (EP) frame: direction of the polarization axis orthogonal to the event plane in the centre-of-mass of the colliding beams
- The studies in Collins–Soper and Helicity frames are also interesting in AA to study quarkonium suppression/regeneration in the QGP



## Magnetic field ( $\vec{B}$ ):

- Huge intensity ( $10^{14}$  T)
- Short lived ( $\tau = 1 \text{ fm}/c$ )

[Kharzeev et al., NPA 803 (2008)]


## Angular momentum ( $\vec{L}$ ):

- Largest in semicentral collisions
- Can affect the system evolution till freeze-out

[Becattini et al., PRC 77 (2008) 024906]




# ALICE detector (Run 2):



- Inclusive quarkonium measurements performed at forward rapidity in the dimuon decay channel

Muon spectrometer acceptance:  $-4.0 < \eta < -2.5$


[ALICE Muon spectrometer]



New measurements from Run 2 datasets

- $pp : \sqrt{s} = 13 \text{ TeV}$
- $Pb-Pb : \sqrt{s_{NN}} = 5.02 \text{ TeV}$

# Quarkonium polarization in Pb-Pb collisions:




- ALICE measurement of  $J/\psi$  polarization in Pb–Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV in Helicity (HE) and Collins-Soper (CS) reference frames
- $\lambda_\theta$  shows a  $2\sigma$  deviation from zero at low  $p_T$
- $3\sigma$  deviation from LHCb measurement in pp collisions in the Helicity frame
- Values compatible with ALICE results in pp collisions within uncertainties

ALICE Collaboration, Phys. Lett. B 815, 136146 (2021)

ALICE Collaboration, Eur. Phys. J. C 78, 562 (2018)


LHCb Collaboration, Eur. Phys. J. C 73, 2631 (2013)

# Quarkonium polarization in Pb-Pb collisions:



- ALICE measurement of  $J/\psi$  polarization in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV
- First measurement with respect to the Event Plane (EP)
- Small but significant polarisation ( $3.5\sigma$ ), particularly in the 40-60% centrality range
- Effect more pronounced at low transverse momentum ( $2 < p_T < 4$  GeV/c) in centrality 30-50%
- Qualitatively in agreement with spin alignment observed for light vector mesons [Phys. Rev. Lett. 125, 012301 (2022)]


ALI-PUB-521052



[ALICE Collaboration, Phys. Rev. Lett. 131, 042303 (2023)]

ALI-PUB-521057

# Quarkonium polarization in pp collisions:




- $J/\psi$  polarization measured in pp collisions in the CS and HE frames
- Dataset : ALICE  $\sqrt{s} = 7$  TeV (2010)  
ALICE  $\sqrt{s} = 8$  TeV (2012)  
LHCb  $\sqrt{s} = 7$  TeV (2011)
- No significant polarisation observed by ALICE and LHCb at forward rapidity
- Need for studies with higher center of mass energies
  - ✓ New ongoing analyses of  $J/\psi$  and  $\psi(2S)$  in pp collisions at  $\sqrt{s} = 13$  TeV

ALICE Collaboration, Phys. Rev. Lett. 108, 082001 (2012)

ALICE Collaboration, Eur. Phys. J. C 78, 562 (2018)

LHCb Collaboration, Eur. Phys. J. C 73, 2631 (2013)

# Quarkonium polarization in pp collisions:




- Recent preliminary measurement of  $\Upsilon(1S)$  polarization at  $\sqrt{s} = 13$  TeV from ALICE
- Results compatible with previous LHCb measurements at  $\sqrt{s} = 8$  TeV
- Polarization is evaluated down to  $p_T \sim 0$
- All values compatible with zero within uncertainties
- Large uncertainties due to limited statistical precision

LHCb Collaboration, JHEP 12, 110 (2017)


ALI-PREL-505590

# Quarkonium polarization in pp collisions:



Analysis Note:  $\psi(2S)$  polarization measurement in pp collisions at  $\sqrt{s} = 13$  TeV, <https://alice-notes.web.cern.ch/node/1472> (ALICE internal)

- Zero polarization within uncertainties for Collins-Soper frame
- Finite polarization in helicity frame for  $4 < p_T < 8$  GeV/c



## Conclusion and Outlook:

- ALICE has measured the polarization of several quarkonium states both in pp and Pb–Pb collisions
- No significant quarkonium polarization till now in pp collisions
- New  $J/\psi$  and  $\psi(2S)$  polarization analyses ongoing in pp collision at  $\sqrt{s} = 13$  TeV
- Results are compatible with other LHC measurements and recent model predictions
- Hint for non-zero polarization at low  $p_T$  in the HE and CS frames in Pb–Pb collisions
- From the results of EP frame analysis, possible correlation with  $\vec{B}$  and  $\vec{L}$  in the QGP formed in heavy-ion collision
- ALICE Run 3 with high luminosity will provide significant statistics for precision measurements

**THANK YOU!**