

Topics on Euclidean parton correlators

Yizhuang Liu, UJ

LaMET 2024

Outline

- Part I: Coordinate-space analyticity and time-ordering issue of TMD soft factors
- Part II: Bjorken limit of 2D large N Gross Neveu

General theory of coordinate space analyticity

- General theory of coordinate space analyticity
 1. Axioms: spectral-condition && micro-locality && temperedness of *Wightman-Distributions*.
 2. Consequence: existence of *analytic Wightman function* in the “permuted extended-tubes” and totally space-like region.
 3. Established in three steps.
 - First: Paley-Wiener type arguments for analyticity in the *forward tubes* $T_n^P = \{(z_n, z_{n-1}, \dots, z_1); \text{Im}(z_{P_k} - z_{P_{k-1}}) \in -V_+\}$ for $\mathcal{W}^P(z_n, \dots, z_1) = \langle \phi(z_{P_n}) \dots \phi(z_{P_1}) \rangle$. Spectral condition is crucial.
 - Second: apply proper complex Lorentz transforms to analytic-continue \mathcal{W}^P further into the *extended forward tubes*.

General theory

- Finally, local-commutativity implies that all the $n!$ analytic Wightman functions \mathcal{W}^P can be combined together to be a single-valued analytic Wightman function in the permuted extended-tubes: Union of all the extended tubes.
5. Important sub-regions of analyticity.
- Euclidean region $\mathcal{E}_n = \{(z_n, \dots, z_1); z_i^0 \in -iR, \text{Im}(\vec{z}_i) = 0, z_i \neq z_j\}$
 - Totally space-like real separations $\{(x_n, \dots, x_1); (x_i - x_j)^2 < 0\}$
7. $\mathcal{W}|_{\mathcal{E}_n}$ are called Schwinger functions && Euclidean correlation functions.

Properties of analytic Wightman functions

8. Properties of analytic Wightman functions
 - Covariance under proper **complex Lorentz transformations**.
 - Permutation symmetry && anti-symmetry.
 - Spin-statistics && CPT.
9. These properties can be non-trivial: for a complex scalar one has $\mathcal{W}_{\phi\phi^\dagger}(-iT e_t) = \mathcal{W}_{\phi\phi^\dagger}(iT e_t)$ ($T > 0$),
$$\langle \phi^\dagger(0) e^{-HT} \phi(0) \rangle = \langle \phi(0) e^{-HT} \phi^\dagger(0) \rangle.$$

This is an operator relation that **flips the operator ordering!**

Properties

10. Relation to real-time Wightman distributions. One can obtain real-time Wightman distribution as boundary-values of analytic Wightman functions.

11. The **Wightman-prescription**

$$\langle \phi(t_n) \dots \phi(t_1) \rangle = \lim_{\eta \rightarrow 0^+} \mathcal{W}(t_n - i\eta e_t, \dots, t_1 - i\eta e_t).$$

- One approaches the boundary point within the forward tube T_n .
- For invariant lengths, W-prescription means $-x_{ij}^2 + i\eta x_{ij}^0$.

12. The **Feynman-prescription**

$$\langle T\phi(t_n) \dots \phi(t_1) \rangle = \lim_{\eta \rightarrow 0^+} \mathcal{W}(t_n(1 - i\eta), \dots, t_1(1 - i\eta))$$

- For invariant lengths, F-prescription means $-x_{ij}^2 + i\eta$.

Schwinger functions from lattice models

- How to realize?
 1. Non-perturbative level. Osterwalder-Schrader reconstruction theorem. Distributions in the Euclidean regions \mathcal{E}_n that are rotational invariant, reflective-positive and grow moderately in n are Schwinger functions of a Wightman QFT and can be uniquely continued back to real time.
 2. Schwinger functions can be obtained as **scaling limits** of lattice models approaching critical points. $\langle \sigma(r\xi)\sigma(0) \rangle_{\xi \rightarrow \infty} \rightarrow Z(\xi) f(r)$. Many examples in 2D. Conjectured for QCD.
 3. Short distance limit: $f(r) \rightarrow \frac{1}{r^{2d}} (1 + r \ln r + \dots)$. **Perturbation to the UV-CFT**. UV of IR = IR of UV.
 4. CFTs are most “perfect” Wightman QFTs. Global (Hilbert space and operator algebra) from UV-asymptotics of local-correlators (OPE).

Analyticity in Perturbation theory

- Momentum space analyticity in DR perturbation theory.
 1. Analyticity in perturbation theory are again due to exponential decay in parametric representations.
 2. In momentum space, *below-threshold* quantities allow Schwinger-parametrization of the form $\int_0^\infty D\alpha F(\alpha) e^{Q^2 P(\alpha)}$ where $P(\alpha) > 0$ are rational functions.
 3. They can be continued to the region $Re(Q^2) < 0$.
- Similarly, in coordinate space for n -point function, one has parametric integrals of the forms $I = \int_0^\infty U(\alpha) e^{\sum_{i<j} x_{ij}^2 P_{ij}(\alpha)}$ for *Euclidean and totally-space-like separations*.

Analyticity in Perturbation theory

- Consider parametric integrals $I = \int_0^\infty U(\alpha) e^{\sum_{i<j} x_{ij}^2 P_{ij}(\alpha)}$.
- 1. The rational functions $P_{ij}(\alpha)$ are positive.
- 2. The $P_{ij}(\alpha)$ allows explicit representations through spanning trees and connected paths between i and j .
- 3. Only depends on invariant length-squares $x_{ij}^2 = (x_i - x_j)^2$.
- 4. Defines analytic function in **the below-threshold region** $\mathcal{E}'_n = \{(z_n, \dots, z_1); \operatorname{Re}(z_{ij}^2) < 0, \forall i \neq j\}$.
- 5. Agrees with spectral representation in $\mathcal{E}'_n \cap T_n^P$ for any P . This is because that $\mathcal{E}'_n \cap T_n^P$ is path-connected and contains $\mathcal{E}_n \cap T_n^P$.

Analyticity in PQCD

- For QCD perturbation theory in covariant gauges (Feynman gauge for example). Spectral condition and local commutativity are satisfied for gluon fields, to all orders.
- Below threshold representation for gluonic correlators exist in Euclidean and totally space-like real points.
- Thus, one has below threshold representation for gluonic Wightman functions in the below-threshold region $\mathcal{E}'_n = \{(z_n, \dots, z_1); \text{Re}(z_{ij}^2) < 0, \forall i \neq j\}$

DY TMD soft factor

- One application of the above is to establish the below-threshold representation for Drell-Yan TMD soft factor in the exponential regulator.
- $$S(b_{\perp}, \nu, \epsilon) = \frac{1}{N_c} \langle \text{Tr} \bar{T} U_{\bar{n}n}(\vec{b}_{\perp} - i\nu e_t) T U_{n\bar{n}}(0) \rangle$$
- $U_{n\bar{n}}(x)$ is a Wilson-line cusp at x , formed by past-pointing gauge-links in light-like directions $n = \frac{1}{\sqrt{2}}(e_t + e_z)$ and $\bar{n} = \frac{1}{\sqrt{2}}(e_t - e_z)$. $\nu > 0$ is the exponential regulator. \vec{b}_{\perp} is the transverse separation.
- The Wilson-loop can be expanded in terms of the gluonic Wightman functions picked-up from the Wilson-loops. Wightman prescriptions are used for the T and \bar{T} from analytic Wightman functions.

Checking invariant lengths

- To see if these analytic Wightman functions allow below-threshold representations, one needs to check the invariant length-squares. There are four types.
 1. Two points under the same T or \bar{T} , on different Wilson-line. $x_{A,ij}^2 = -2\lambda_i\lambda_j < 0$. Space-like.
 2. One point from T , another from \bar{T} , on same Wilson-line direction. $x_{B,ij}^2 = -v^2 - b_{\perp}^2 - \sqrt{2iv}(\lambda_i^L - \lambda_j^R)$. Below-threshold.
 3. One point from T , another from \bar{T} , on different Wilson-line directions. $x_{C,ij}^2 = -v^2 - b_{\perp}^2 - 2\lambda_i^L\lambda_j^R - \sqrt{2iv}(\lambda_i^L - \lambda_j^R)$. Below-threshold.

Null separation and below-threshold representation

4. Two points on the same Wilson-line. This is tricky since null-separation is encountered. But the $i\eta$ solves the problem.
5. $(\lambda_i n - \lambda_j n - i\eta e_t)^2 = -\eta^2 - \sqrt{2}i\eta(\lambda_i - \lambda_j)$. Approached within the blow-threshold region.
 - Thus, below-threshold representation exists.
 - Furthermore, the η s can be send to zero from the beginning for three reasons.
 1. The $-\eta^2$ regulates UV-light-cone divergences, which are regulated by the DR already.
 2. The $i\eta$ terms always have the same signs within the T and \bar{T} groups as the $i\nu$ terms. Thus, $i\eta$ s are replaced by the $i\nu$ s.
 3. The $-\eta^2$, $i\eta$ terms are added to terms with negative real parts that never vanish in the integration region.

Below-threshold representation

- Thus, we conclude that the DY TMD soft-factor allows below-threshold representations in terms of three invariant lengths:
 1. $x_{A,ij}^2 = -2\lambda_i\lambda_j$
 2. $x_{B,ij}^2 = -\nu^2 - b_{\perp}^2 - \sqrt{2i\nu}(\lambda_i^L - \lambda_j^R)$.
 3. $x_{C,ij}^2 = -\nu^2 - b_{\perp}^2 - 2\lambda_i^L\lambda_j^R - \sqrt{2i\nu}(\lambda_i^L - \lambda_j^R)$
- As far as $\nu \neq 0$ and $\epsilon \neq 0$, gluonic Wightman functions restricted to these separations are still covariant and permutation-symmetric.
- For $\nu = 0$, naïve invariant lengths for the DY-shape TMD soft factors. Can be used for the (non-gauge-invariant) δ regulator.

Relationship between soft factors: $S = S_t$

- The existence of below-threshold representation can be used to establish certain identities.
- Consider $S_t(\mathbf{b}_\perp, \nu, \epsilon) = \frac{1}{N_c} \langle \text{Tr} T \tilde{U}_{n\bar{n}}^\dagger(\vec{b}_\perp - \nu \mathbf{e}_z) \tilde{U}_{n\bar{n}}(0) \rangle$.
- 1. Here $\tilde{U}_{n\bar{n}}(0)$ is a Wilson-line cusp with future-pointing light-like link in \bar{n} directions.
- 2. Overall time-ordering.
- 3. A quark-anti-quark pair in n direction moving from past to $t = 0$, then transits to another pair in \bar{n} propagating to future. Space-like form factor.
- 4. $-\nu \mathbf{e}_z$ in the \mathbf{e}_z direction.

Relationship between soft factors: $S = S_t$

- One can show that $S = S_t$ based on Minkowskian parametric representations of S_t .
- The M-parametric representations of S_t , after **Wick-rotation**, become exactly the below-threshold representations for S .
- Thus, the DY-TMD soft factor can be represented as a **space-like form factor**.

Generalization: Analytic Wilson-loop averages.

- We can conjecture the following:
- For any closed complex-space-time valued oriented loop \mathcal{C} , if
 1. The loop is piece-wisely smooth with finite-numbers of cusp singularities with finite cusp angles.
 2. An arbitrary non-coinciding set of points picked up from \mathcal{C} always lives in the natural coordinate-space analyticity region (such as permuted – extended -tubes).
- Then the analytic Wilson-loop average $\langle W(\mathcal{C}) \rangle$ exists and behave like the analytic Wightman functions in the analyticity region.

Generalization: Analytic Wilson-loop averages.

- The analytic Wilson-loop average $\langle TrW(\mathcal{C}) \rangle$ depends only on the \mathcal{C} and the orientation.
- If $\mathcal{C} = \mathcal{C}_1 \cup \mathcal{C}_2 \cup \mathcal{C}_3 \dots \mathcal{C}_n$ with $\mathcal{C}_i \cap \mathcal{C}_j = \emptyset$. Then $\langle TrW(\mathcal{C}) \rangle = \langle TrW(\mathcal{C}^P) \rangle$ where $\mathcal{C}^P = \mathcal{C}_{P_1} \cup \mathcal{C}_{P_2} \cup \mathcal{C}_{P_3} \dots \mathcal{C}_{P_n}$. This plays the role of local-commutativity.
- For small Wilson-loop sizes, $\langle TrW(\mathcal{C}) \rangle$ allows perturbative expansion in terms of the perturbative gluonic Wightman functions.
- Analytic Wilson-loops leads to analytic Wightman functions of gauge-invariant operators such as $tr F^2$, if one performs small size OPE for the Wilson-loops.

Soft factor relations for three rapidity regulators.

- Given the above, one can define TMD soft factors that contains three regulators at once: off-light-cone, finite LF-length and exponential.
 1. $S_t(T_1, T_2, b_\perp, v, Y, \epsilon)$: still a “real-time” Wilson-loop with “transverse” gauge links in $\vec{b}_\perp - v e_z$ directions.
 - S_t is defined with time-like links with $v = n + e^{-Y} \bar{n}$ and $v' = \bar{n} + e^{-Y} n$. Resembles the heavy-quark form factor in the 2019 Ji-Liu-Liu paper.
 - Time-ordering: $T_1 = T_1(1 - i\eta)$ and $T_2 = T_2(1 - i\eta)$. Can be analytically continued smoothly to Euclidean times $T_1 = -iL^-$ and $T_2 = -iL^+$ where $L^\pm > 0$.
 2. $S(L^+, L^-, b_\perp, v, Y, \epsilon)$: a complex-valued Wilson-loop with “transverse” gauge-links in $\vec{b}_\perp - i v e_z$ direction.

Soft factor relations for three rapidity regulators.

- S is defined with space-like links in $n_Y = n - e^{-Y} \bar{n}$ and $\bar{n}_Y = \bar{n} - e^{-Y} n$. Resembles the Collins off-light-cone TMD-soft factor.
 - All underlying separations for $S(L^+, L^-, b_\perp, v, Y, \epsilon)$ are below-threshold. No null separations at all.
3. Complex Lorentz transform : $\Lambda(t, z) = (iz, it)$, or $\Lambda(e_t, e_z) = (ie_z, ie_t)$.
- Under Λ , $v \rightarrow in_Y$, $v' \rightarrow -i \bar{n}_Y$ and $-ve_z \rightarrow -ive_t$.
 - The Wilson-loop for $S_t(-iL^-, -iL^+, b_\perp, v, Y, \epsilon)$ maps exactly to the Wilson-loop for $S(L^+, L^-, b_\perp, v, Y, \epsilon)$ under the Λ .
4. Thus, one has the master equality $S_t(-iL^-, -iL^+, b_\perp, v, Y, \epsilon) = S(L^+, L^-, b_\perp, v, Y, \epsilon)$.

Comments

- The relations above implies that the rapidity evolution kernel for TMDPDFs and for LFWFs are same : S_t is the natural soft factor for LFWFs.
- The renormalization are multiplicative.
- Three standard orders of limits
 1. $Y \rightarrow \infty$ first, $L^\pm \rightarrow \infty$ second gives the exponential regulator.
 2. $Y \rightarrow \infty$ first, $\nu \rightarrow 0$ second gives the finite LF length regulator.
 3. $\nu \rightarrow 0, L^\pm \rightarrow \infty$ first at finite Y gives the off-light-cone regulator.
- Another possibility, keep L^\pm and ν finite, is it possible that $Y \rightarrow \infty$ and $\epsilon \rightarrow 0$ are related to each-other perturbatively ?

Outline

- Part I: Coordinate-space analyticity and time-ordering issue of TMD soft factors
- Part II: Bjorken limit of 2D large N Gross Neveu

Exact results for space-like structure function in 2D Gross-Neveu

- Consider the 2D Gross Neveu in large N . $\mathcal{L} = \bar{\psi} i \gamma \cdot \partial \psi - \sigma \bar{\psi} \psi - \frac{\sigma^2}{2g_0^2}$
 1. Large N expansion. Condensate $\sigma_0 = m$ (fermion mass). Running coupling $\frac{1}{g^2(\mu)} = \frac{N}{2\pi} \ln \frac{\mu^2}{m^2}$.
 2. Large N expansion can be performed systematically using effective coupling $g^2(k) = \frac{2\pi}{N} \int_0^\infty \int_{c-i\infty}^{c+i\infty} \frac{dt ds}{2\pi i} \frac{\Gamma(1-2s)\Gamma(s+t)}{\Gamma(1-s+t)} \left(\frac{m^2}{-k^2}\right)^{-s}$. The propagator for $\sigma = m$.
 3. Large k^2 expansion: shifts to $s = -n - t$. Borel-integrals at power $\left(\frac{m^2}{-k^2}\right)^n$. Marginality manifest.

Space-like structure function

- Define the “twist-three-type” correlator $\mathcal{E}(z^2 m^2, \lambda) \bar{u}(p) u(p) = \langle p, i | \bar{\psi}^i(x) \psi^i(x) | p, i \rangle - \langle p, i | \bar{\psi}^j(x) \psi^j(x) | p, i \rangle$.
- 1. $z^2 = -x^2 > 0$ space-like and $\lambda = -p \cdot x$. Analyticity in λ in whole complex plane.
- 2. We calculate $\mathcal{E}(z^2 m^2, \lambda)$ to NLO in $\frac{1}{N}$. One-bubble-chain diagrams.
- 3. $\mathcal{E}^{(1)}(z^2 m^2, \lambda) = \frac{2\pi}{N} e^{-i\lambda} (-F_1 + F_2 - F_3)(z^2 m^2, \lambda)$.
- 4. Bjorken limit $z^2 \rightarrow 0$ at fixed λ . Exact twist-expansion.

Twist-expansion.

- Hard functions and non-perturbative functions.

1.
$$F_1(z^2 m^2, \lambda) = \sum_{l=0}^{\infty} \left(\frac{z^2 m^2}{4}\right)^l \int_0^{\infty} dt q_1^{(l)}(t, \lambda, \mu) + \sum_{l=0}^{\infty} \left(\frac{z^2 m^2}{4}\right)^l \int_0^{\infty} dt \sum_{p=0}^{\infty} \left(\frac{z^2 m^2}{4}\right)^p \left(\left(\frac{z^2 m^2}{4}\right)^t \mathcal{H}_1^{l,p}(t, \lambda, \mu) + q_1^{l,p}(t, \lambda, \mu) \right)$$
2. Borel integrands $\mathcal{H}_1^{l,p}(t, \lambda, \mu)$ contains renormalon singularity at $t = n$ that cancels with the singularity of $q_1^{l,p+n}(t, \lambda, \mu)$.
3. The μ dependency cancels between $\mathcal{H}_1^{l,p}$ and $q_1^{l,p}$ for $p \geq 1$.
4. For $p = 0$, μ dependency cancels between $q_1^{(l)}(t, \lambda, \mu)$ and $\mathcal{H}_1^{l,0}$. $q_1^{l,0} \equiv 0$.
5. $q_1^{(l)}(t, \lambda, \mu)$ contains no Borel singularity at $t \geq 0$.

Operator content at LP

- Operator content. There are four quark operators even at LP.

- The "Hard function" at LP reads

$$\mathcal{H}^{(0)}(t, \alpha(z), \lambda) = \frac{1}{4\pi} \left(\frac{1}{t} {}_1\tilde{F}_1(2, 1, -\lambda) + \left(\frac{z^2 m^2}{4} \right)^t \Gamma(-t) {}_1\tilde{F}_1(2, 1+t, -\lambda) \right) \\ + \frac{\lambda}{2\pi} \left(\frac{1}{t} {}_1\tilde{F}_1(2, 2, -\lambda) + \left(\frac{z^2 m^2}{4} \right)^t \Gamma(-t) {}_1\tilde{F}_1(2, 2+t, -\lambda) \right).$$

- The first-line: explained by the operators $\sum_{n=0}^{\infty} \frac{1}{n!} \mathcal{H}_n(\alpha(z)) x_{\mu_1} \dots x_{\mu_n} \bar{\psi}_i \overleftrightarrow{\partial}^{\{\mu_1 \dots \mu_n\}} \psi_i$

- Second line: explained by the four-quark operators

$$\sum_{n=0}^{\infty} \frac{1}{n!} \tilde{\mathcal{H}}_n(\alpha(z)) x_{\mu_1} x_{\mu_2} \dots x_{\mu_{n+1}} \bar{\psi}_i \gamma^{\{\mu_1 \overleftrightarrow{\partial}^{\mu_2} \dots \overleftrightarrow{\partial}^{\mu_{n+1}}\}} \psi_i \bar{\psi} \psi$$

Operator content at LP : condensate contributes

4. Due to the fact that $g_0^2 \langle \bar{\psi} \psi \rangle = -m(1 + O(\frac{1}{N}))$. The contributions from

$$\sum_{n=0}^{\infty} \frac{1}{n!} \tilde{\mathcal{H}}_n(\alpha(z)) x_{\mu_1} x_{\mu_2} \dots x_{\mu_{n+1}} \bar{\psi}_i \gamma^{\{\mu_1 \overleftrightarrow{\partial}^{\mu_2} \dots \overleftrightarrow{\partial}^{\mu_{n+1}}\}} \psi_i \bar{\psi} \psi$$

are non-vanishing at the order $\frac{1}{N}$. Namely, $\tilde{\mathcal{H}}_n$ is of order $g_0^2 \frac{1}{N}$, while $\langle \bar{\psi} \psi \rangle$ is of order N .

- Thus, vacuum condensates start to contribute even at the leading power.
- At NLP ($O(z^2 m^2)$), there are up to eight quark operators $\bar{\psi} \gamma^+ (\partial^+)^n \psi$ $(\bar{\psi} \psi)^3$.
- Parton picture is non-longer convenient.

Twist-expansion and threshold expansion

- The small- z^2 expansion, in terms of the Borel-resummed hard and “collinear” functions, converges absolutely for any $z^2 < 0$.
- No instanton-like contributions in the coefficient functions.
- The threshold expansion $\lambda \rightarrow +i\infty$ can also be performed exactly.
 1. Threshold expansion in $\frac{1}{-i\lambda}$ commute with small z^2 expansion.
 2. Threshold expansion is asymptotic. Resurgence analysis can be performed.
 3. “Conspiracy” between Borel singularity of threshold expansion and branch-singularity of $\frac{1}{\ln x}$ for small- x expansion.

Conclusion

1. Introduction to coordinate-space analyticity in local QFT.
2. Relationships between TMD soft factors as an application.
3. Generalizable to three rapidity regulators implemented simultaneously.
4. Space-like structure function in 2D large N Gross-Neveu carefully investigated. Convergence of small z^2 expansion.
5. Vacuum condensate contribute even at LP.