




上海交通大学
SHANGHAI JIAO TONG UNIVERSITY

粒子与核物理研究所



Light-cone distribution amplitudes of a light baryon in LaMET

曾军 (Jun Zeng)

Institute of Nuclear and Particle Physics/INPAC,
Shanghai Jiao Tong University

Based on: JHEP 07 (2023) 191

Collaborators: Zhi-Fu Deng, Chao Han, Wei Wang, Jia-Lu Zhang (SJTU)

14 Aug 2024

2024 Meeting on Lattice Parton Physics from Large-Momentum
Effective Theory (LaMET 2024)

Outline



1

Introduction

2

Baryon LCDA

3

Matching Kernel

4


Expansion by regions

5


Summary and outlook

- Feynman proposed a parton model more than 50 years ago, and hadron structure information has been obtained by fitting a large number of high-energy experimental data.
- The hadron light-cone distribution amplitude (LCDAs) is a physical quantity that describes the momentum distribution of all parts of hadrons and reflects the internal structure of hadrons.
- The calculation of the hadron LCDAs by the basic theory of the strong interaction has been slow for a long time.


Especially the baryon LCDAs !



- **Describing Internal Structure:** LCDAs provide detailed information about the distribution of quark and gluons inside hadrons, especially in high-energy collisions.
- **Calculating Hadronic Decay Processes:** They help theoretical physicists predict experimental outcomes and compare them with actual observations, thereby validating or refining existing theories.
- **Non-Perturbative QCD Studies:** Through lattice QCD and other non-perturbative methods, LCDAs can be calculated from first principles, providing deep insights into strong interactions.
- **Exploring New Physics:** In the search for new physics beyond the Standard Model, LCDAs offer a crucial theoretical framework.



Inclusive: PDF



Exclusive: LCDAs

LCDAs is very very important !



Challenging:

- 1. Nonperturbative:** cannot be calculated directly from first principles; must be extracted from experimental data or nonperturbative QCD approaches, such as lattice QCD simulations or phenomenological models.
- 2. Large number of degrees of freedom:** The determination of LCDAs involves accounting for the contributions from all possible quark and gluon configurations within the baryon, leading to a large number of degrees of freedom and complicating the analysis.
- 3. Complexity of the operator product expansion (OPE):** the OPE involves three-quark operators and gluonic operators, leading to a more complex and computationally challenging analysis compared to mesons.
- 4. Limited experimental data:** limited experimental data for baryon structure observables, makes it challenging to constrain baryon light-cone distribution functions directly from the experiment.



Lattice QCD

1. Nucleon distribution amplitudes and proton decay matrix elements on the lattice(Vladimir M. Braun, 2009) **present a calculation of the first few moments of the leading-twist nucleon DA**
2. Light-cone distribution amplitudes of the nucleon and negative parity nucleon resonances from lattice QCD(Vladimir M. Braun, 2014) **calculate moments**
3. Light-cone distribution amplitudes of octet baryons from lattice QCD (RQCD Collaboration,2019)

The Baryon LCDA from Lattice QCD, See Jun Hua's slides!

Model

1. Modelling the Nucleon WF from Soft and hard processes(1996) **parameterization**
2. Nucleon distribution amplitude: The heterotic solution(1993)
amalgamates features of the Chernyak-Ogloblin-Zhitnitsky model with those of the Cari-Stefanis model
3. Nucleon WF and Form Factors in QCD(1983)
A model for the nucleon wave function is proposed based on a knowledge of these few first moments.
4. LCDA of the baryon(2021) **Chiral quark-soliton model**
5. Estimates of the isospin-violating $\Lambda b \rightarrow \Sigma^0 \varphi; \Sigma^0 J = \psi$ decays and the $\Sigma - \Lambda$ mixing(2023) **COZ model**





QCD sum rule

1. Nucleon WFs and QCD sum rules(1987)

2. Higher twist distribution amplitudes of the nucleon in QCD(2000)

present the first systematic study of higher-twist LCDAs of the nucleon in QCD.

Nonperturbative input parameters are estimated from QCD sum rules.

3. $\Lambda_b \rightarrow p$ transition form factors in perturbative QCD(2022)

Ligh-cone Sum Rule

1. Nucleon form factors and distribution amplitudes in QCD(2013)

extracted from the comparison with the experimental data on form factors

2. Wave functions of octet baryons(1989)


The model wave functions are proposed which fulfill the sum rules requirements.

3. Nucleon form factors in QCD(2006)






1. Introduction



octet baryon



decuplet baryon

The others baryon present in JHEP 07 (2024) 019 . See *Chao Han's slides!*





1. Introduction



At leading twist, the LCDAs of an octet baryon can be decomposed into three terms as


$$\begin{aligned}
& \langle 0 | f_\alpha(z_1 n) g_\beta(z_2 n) h_\gamma(z_3 n) | B(P_B, \lambda) \rangle \\
&= \frac{1}{4} f_V \left[(\not{P}_B C)_{\alpha\beta} (\gamma_5 u_B)_\gamma \underline{V^B(z_i n \cdot P_B)} + (\not{P}_B \gamma_5 C)_{\alpha\beta} (u_B)_\gamma \underline{A^B(z_i n \cdot P_B)} \right] \\
&+ \frac{1}{4} f_T (i\sigma_{\mu\nu} P_B^\nu C)_{\alpha\beta} (\gamma^\mu \gamma_5 u_B)_\gamma \underline{T^B(z_i n \cdot P_B)},
\end{aligned}$$

where $C \equiv i\gamma^2\gamma^0$ is the charge conjugation matrix, u_B stands for the spinor for the baryon and $\sigma_{\mu\nu} = \frac{i}{2} [\gamma_\mu, \gamma_\nu]$. $f_{V/A/T}$ is the corresponding decay constant for each LCDA. For proton and neutron, $f_T = f_V$ due to the isospin symmetry.

$$\langle 0 | f^T(z_1 n) (C\not{n})g(z_2 n) h(z_3 n) | B \rangle = -f_V V^B(z_i n \cdot P_B) P_B^+ \gamma_5 u_B,$$

$$\langle 0 | f^T(z_1 n) (C\gamma_5\not{n})g(z_2 n) h(z_3 n) | B \rangle = f_V A^B(z_i n \cdot P_B) P_B^+ u_B,$$

$$\langle 0 | f^T(z_1 n) (iC\sigma_{\mu\nu}n^\nu)g(z_2 n) \gamma^\mu h(z_3 n) | B \rangle = 2f_T T^B(z_i n \cdot P_B) P_B^+ \gamma_5 u_B,$$






1. Introduction



- **LCDAs** are defined as the correlation functions of light-cone operators inside a hadron, these quantities **can not be directly evaluated on the lattice**.
- It is highly indispensable to develop a method to calculate the full shape of baryon LCDAs from the first principle of QCD.
- A very inspiring approach was proposed to circumvent this problem and is now formulated as the **large-momentum effective theory** (LaMET).

- The LaMET provides a method to calculate the hadron LCDA from lattice QCD from first principles.
- The LCDA can be extracted by matching the quasi-DA to light cone direction by perturbation.





2. Baryon LCDA and quasi-DA



$$\Phi(x_1, x_2) f_{\Lambda} n \cdot p u_{\Lambda}(p) = \int \frac{dt_1 n \cdot p}{2\pi} \int \frac{dt_2 n \cdot p}{2\pi} e^{ix_1 n \cdot pt_1 + ix_2 n \cdot pt_2} \epsilon_{ijk} \langle 0 | U_i^T(t_1 n) \Gamma D_j(t_2 n) S_k(0) | \Lambda \rangle$$

where $\Gamma = C\gamma^5 \eta$


$$\tilde{\Phi}(x_1, x_2) \tilde{f}_{\Lambda} n_z \cdot p u_{\Lambda}(p) = \int \frac{dt_1 n_z \cdot p}{2\pi} \int \frac{dt_2 n_z \cdot p}{2\pi} e^{ix_1 n_z \cdot pt_1 + ix_2 n_z \cdot pt_2} \epsilon_{ijk} \langle 0 | U_i^T(t_1 n_z) \tilde{\Gamma} D_j(t_2 n_z) S_k(0) | \Lambda \rangle$$

$$Q_i(x) = \mathcal{W}_{ii'}(\infty, x) q_{i'}(x)$$

$$\tilde{\Gamma} = C\gamma^5 \eta_{\lambda} \quad (\lambda = t \text{ or } z)$$



Matching





2. Baryon LCDA and quasi-DA: Matching




$$\tilde{\Phi}(x_1, x_2, \mu) = \int dy_1 dy_2 \mathcal{C}(x_1, x_2, y_1, y_2, \mu) \Phi(y_1, y_2, \mu) + \mathcal{O}\left(\frac{1}{x_1 p^z}, \frac{1}{x_2 p^z}, \frac{1}{(1-x_1-x_2)p^z}\right)$$

Matching kernel is insensitive to the hadrons, in the calculation of LCDAs one can replace the hadron by the partonic state.

$$|\Lambda\rangle \rightarrow \frac{\epsilon_{abc}}{6} |u_a(k_1) d_b(k_2) s_c(k_3)\rangle$$

the quark state is chosen to have the same J^{PC} with the Λ .





2. Baryon LCDA and quasi-DA: partonic LCDA



$$\phi(x_1, x_2, \mu) S = \int \frac{dt_1 p^+}{2\pi} \int \frac{dt_2 p^+}{2\pi} e^{ix_1 p^+ t_1 + ix_1 p^+ t_2} \frac{\epsilon_{ijk} \epsilon_{abc}}{6} \langle 0 | U_i^T(t_1 n) \Gamma D_j(t_2 n) S_k(0) | u_a(k_1) d_b(k_2) s_c(k_3) \rangle$$










$$\tilde{\phi}(x_1, x_2, \mu) \tilde{S} = \int \frac{dt_1 p^z}{2\pi} \int \frac{dt_2 p^z}{2\pi} e^{ix_1 p^z t_1 + ix_1 p^z t_2} \frac{\epsilon_{ijk} \epsilon_{abc}}{6} \langle 0 | (U_i)^T(t_1 n_z) \tilde{\Gamma} D_j(t_2 n_z) S_k(0) | u_a(k_1) d_b(k_2) s_c(k_3) \rangle$$

Normalization factor:

$$S = \frac{\epsilon_{ijk} \epsilon_{abc}}{6} \langle 0 | (U_i)^T(0) \Gamma D_j(0) S_k(0) | u_a(k_1) d_b(k_2) s_c(k_3) \rangle$$

$$\tilde{S} = \frac{\epsilon_{ijk} \epsilon_{abc}}{6} \langle 0 | (U_i)^T(0) \tilde{\Gamma} D_j(0) S_k(0) | u_a(k_1) d_b(k_2) s_c(k_3) \rangle$$





$$f(x_1, x_2, x_{1,0}, x_{2,0}) =$$

$$\Phi(x_1, x_2) = \delta(x_1 - x_{1,0})\delta(x_2 - x_{2,0}) - \frac{\alpha_s C_F}{8\pi} \frac{1}{\epsilon_{IR}}$$

$$\times \left\{ \frac{\delta(x_1 - x_{1,0})\theta(x_1)\theta(\bar{x}_1)\theta(x_2)\theta(\bar{x}_2)}{\bar{x}_1(x_2 - x_{2,0})} \right.$$

$$\times \left[\frac{x_3(x_2 - x_{2,0} + 2\bar{x}_1)}{x_{3,0}}\theta(x_2 - x_{2,0}) \right.$$

$$\left. + \frac{x_2(x_2 - x_{2,0} - 2\bar{x}_1)}{x_{2,0}}\theta(x_{2,0} - x_2) \right]$$

$$+ \frac{\delta(x_3 - x_{3,0})2x_1\theta(x_1)\theta(\bar{x}_1)\theta(x_2)\theta(\bar{x}_2)}{x_{1,0}}$$

$$\times \left[\frac{\theta(x_2 - x_{2,0})}{x_1 + x_2} - \frac{\theta(x_{1,0} - x_1)}{x_1 - x_{1,0}} \right]$$

$$\left. + \{x_1 \leftrightarrow x_2, x_{1,0} \leftrightarrow x_{2,0}\} \right\} \oplus$$

Diagram of DA. Self-energies of external lines are not shown.



2. evolution of LCDA



$$\frac{d\Phi(x_1, x_2, \mu)}{d \ln \mu} = \frac{\alpha_s C_F}{4\pi} \int dy_1 \int dy_2 V \Phi(y_1, y_2, \mu).$$

At the one-loop, the evolution kernel V is

$$V = f(x_1, x_2, y_1, y_2).$$

Efremov-Radyushkin-Brodsky-Lepage (ERBL) evolution kernel.






2. LCDA and Quasi-DA



$$\begin{aligned} \tilde{\phi}(x_1, x_2, \mu) = & \delta(x_1 - x_{1,0})\delta(x_2 - x_{2,0}) + \frac{\alpha_s C_F}{8\pi} \\ & \times \left\{ \left[g_2 \delta(x_2 - x_{2,0}) + g_3 \delta(x_3 - x_{3,0}) \right. \right. \\ & \left. \left. + \{x_2 \leftrightarrow x_1, x_{2,0} \leftrightarrow x_{1,0}\} \right] \right\}_{\oplus}, \end{aligned}$$

where the \oplus denote

$$\begin{aligned} [g(x_1, x_2, x_{1,0}, x_{2,0})]_{\oplus} = & g(x_1, x_2, x_{1,0}, x_{2,0}) \\ & - \delta(x_1 - x_{1,0})\delta(x_2 - x_{2,0}) \\ & \times \int dy_1 dy_2 g(y_1, y_2, x_{1,0}, x_{2,0}) \end{aligned}$$





2. LCDA and Quasi DA



$$g_2 = \begin{cases} \frac{(x_{1,0} + x_1)(x_{3,0} + x_3) \ln\left(-\frac{x_{3,0} - x_3}{x_3}\right) - x_1(2x_{1,0} + x_{3,0} + x_3) \ln\left(-\frac{x_1}{x_3}\right)}{x_{1,0}x_{3,0}(x_{3,0} - x_3)} - \frac{x_1(2x_{1,0} + x_{3,0} + x_3) \ln\left(-\frac{x_1}{x_3}\right)}{x_{1,0}(x_{3,0} - x_3)(x_{1,0} + x_{3,0})}, & x_1 < 0 \\ \frac{x_1(-x_{1,0} + 2x_{2,0} + x_1 - 2)}{(x_1 - x_{1,0})x_{1,0}(x_{2,0} - 1)\epsilon\mathbb{R}} + \frac{2x_1 \ln\left(\frac{4x_1(x_3 - x_{3,0})(p^z)^2}{\mu^2}\right)}{x_{1,0}(x_3 - x_{3,0})} + \frac{x_1 \ln\left(\frac{4x_1x_3(p^z)^2}{\mu^2}\right)}{x_{1,0}(x_{1,0} + x_{3,0})} \\ + \frac{x_1(-3x_{1,0} - 2x_{3,0} + x_1)}{x_{1,0}(x_3 - x_{3,0})(x_{1,0} + x_{3,0})} - \frac{((x_3 - x_{3,0})^2 - 2x_3x_{1,0}) \ln\left(\frac{x_3 - x_{3,0}}{x_3}\right)}{x_{1,0}(x_3 - x_{3,0})x_{3,0}}, & 0 < x_1 < x_{1,0} \\ \frac{x_3(-x_{1,0} - 2x_{2,0} + x_1 + 2)}{(x_1 - x_{1,0})(x_{2,0} - 1)x_{3,0}\epsilon\mathbb{R}} + \frac{2x_3 \ln\left(\frac{4x_3(x_1 - x_{1,0})(p^z)^2}{\mu^2}\right)}{(x_1 - x_{1,0})x_{3,0}} + \frac{x_3 \ln\left(\frac{4x_1x_3(p^z)^2}{\mu^2}\right)}{x_{3,0}(x_{1,0} + x_{3,0})} \\ + \frac{x_3(-2x_{1,0} - 3x_{3,0} + x_3)}{(x_1 - x_{1,0})x_{3,0}(x_{1,0} + x_{3,0})} - \frac{((x_1 - x_{1,0})^2 - 2x_1x_{3,0}) \ln\left(\frac{x_1 - x_{1,0}}{x_1}\right)}{(x_1 - x_{1,0})x_{1,0}x_{3,0}}, & x_{1,0} < x_1 < x_{1,0} + x_{3,0} \\ \frac{(x_{1,0} + x_1)(x_{3,0} + x_3) \ln\left(-\frac{x_{3,0} - x_3}{x_3}\right) - x_1(2x_{1,0} + x_{3,0} + x_3) \ln\left(-\frac{x_1}{x_3}\right)}{x_{1,0}x_{3,0}(x_{3,0} - x_3)} - \frac{x_1(2x_{1,0} + x_{3,0} + x_3) \ln\left(-\frac{x_1}{x_3}\right)}{x_{1,0}(x_{3,0} - x_3)(x_{1,0} + x_{3,0})}, & x_1 > x_{1,0} + x_{3,0} \end{cases}$$





2. LCDA and Quasi DA



$$g_3 = \begin{cases} \frac{(x_{1,0}x_{2,0} + x_1x_2) \ln\left(\frac{x_2 - x_{2,0}}{x_2}\right)}{x_{1,0}(x_2 - x_{2,0})x_{2,0}} - \frac{x_1(x_{1,0} + x_2) \ln\left(-\frac{x_1}{x_2}\right)}{x_{1,0}(x_2 - x_{2,0})(x_{1,0} + x_{2,0})}, & x_1 < 0 \\ \frac{2x_1(x_{1,0} + x_2)}{(x_1 + x_2)(x_1 - x_{1,0})x_{1,0}\epsilon\text{IR}} + \frac{x_1 \ln\left(\frac{4x_1(x_2 - x_{2,0})(p^z)^2}{\mu^2}\right)}{x_{1,0}(x_2 - x_{2,0})} + \frac{x_1 \ln\left(\frac{4x_1x_2(p^z)^2}{\mu^2}\right)}{x_{1,0}(x_{1,0} + x_{2,0})} \\ + \frac{1}{x_1 - x_{1,0}} + \frac{2x_1 + x_2}{x_{1,0}(x_{1,0} + x_{2,0})} + \frac{(x_{1,0}(x_{2,0} + x_1) - x_1^2) \ln\left(\frac{x_2 - x_{2,0}}{x_2}\right)}{x_{1,0}(x_2 - x_{2,0})x_{2,0}}, & 0 < x_1 < x_{1,0} \\ \frac{2x_2(x_{2,0} + x_1)}{(x_1 + x_2)(x_2 - x_{2,0})x_{2,0}\epsilon\text{IR}} + \frac{x_2 \ln\left(\frac{4x_2(x_1 - x_{1,0})(p^z)^2}{\mu^2}\right)}{(x_1 - x_{1,0})x_{2,0}} + \frac{x_2 \ln\left(\frac{4x_1x_2(p^z)^2}{\mu^2}\right)}{x_{2,0}(x_{1,0} + x_{2,0})} \\ + \frac{1}{x_2 - x_{2,0}} + \frac{x_1 + 2x_2}{x_{2,0}(x_{1,0} + x_{2,0})} + \frac{((x_{1,0} + x_2)x_{2,0} - x_2^2) \ln\left(\frac{x_1 - x_{1,0}}{x_1}\right)}{(x_1 - x_{1,0})x_{1,0}x_{2,0}}, & x_{1,0} < x_1 < x_{1,0} + x_{2,0} \\ \frac{(x_{1,0}x_{2,0} + x_1x_2) \ln\left(\frac{x_1 - x_{1,0}}{x_1}\right)}{(x_1 - x_{1,0})x_{1,0}x_{2,0}} - \frac{x_2(x_{2,0} + x_1) \ln\left(-\frac{x_2}{x_1}\right)}{(x_1 - x_{1,0})x_{2,0}(x_{1,0} + x_{2,0})}, & x_1 > x_{1,0} + x_{2,0}. \end{cases}$$





3. Macting kernel



$$\tilde{\Phi}(x_1, x_2, \mu) = \int dy_1 dy_2 \mathcal{C}(x_1, x_2, y_1, y_2, \mu) \Phi(y_1, y_2, \mu) + \mathcal{O}\left(\frac{1}{x_1 p^z}, \frac{1}{x_2 p^z}, \frac{1}{(1-x_1-x_2)p^z}\right)$$

$$\mathcal{C}(x_1, x_2, y_1, y_2, \mu) = \delta(x_1 - y_1) \delta(x_2 - y_2) + \frac{\alpha_s C_F}{8\pi}$$

$$\times \left[C_2(x_1, x_2, y_1, y_2) \delta(x_2 - y_2) \right.$$

$$+ C_3(x_1, x_2, y_1, y_2) \delta(x_3 - y_3)$$

$$\left. + \{x_1 \leftrightarrow x_2, y_1 \leftrightarrow y_2\} \right], \oplus$$





3. Macting kernel



$$C_2(x_1, x_2, y_1, y_2) = \begin{cases} \frac{(x_1 + y_1)(x_3 + y_3) \ln \frac{y_1 - x_1}{-x_1}}{y_1(y_1 - x_1)y_3} - \frac{x_3(x_1 + y_1 + 2y_3) \ln \frac{x_3}{-x_1}}{(y_1 - x_1)y_3(y_1 + y_3)}, & x_1 < 0 \\ \frac{(x_1 - 3y_1 - 2y_3)x_1}{y_1(x_3 - y_3)(y_1 + y_3)} - \frac{[(x_3 - y_3)^2 - 2x_3y_1] \ln \frac{x_3 - y_3}{x_3}}{y_1(x_3 - y_3)y_3} + \frac{2x_1 \ln \frac{4x_1(x_3 - y_3)p_z^2}{\mu^2}}{y_1(x_3 - y_3)} + \frac{x_1 \ln \frac{4x_1x_3p_z^2}{\mu^2}}{y_1(y_1 + y_3)}, & 0 < x_1 < y_1 \\ \frac{(x_3 - 2y_1 - 3y_3)x_3}{y_3(x_1 - y_1)(y_1 + y_3)} - \frac{[(x_1 - y_1)^2 - 2x_1y_3] \ln \frac{x_1 - y_1}{x_1}}{(x_1 - y_1)y_1y_3} + \frac{2x_3 \ln \frac{4x_3(x_1 - y_1)p_z^2}{\mu^2}}{(x_1 - y_1)y_3} + \frac{x_3 \ln \frac{4x_1x_3p_z^2}{\mu^2}}{y_3(y_1 + y_3)}, & y_1 < x_1 < y_1 + y_3 \\ \frac{(x_1 + y_1)(x_3 + y_3) \ln \frac{y_3 - x_3}{-x_3}}{y_1y_3(y_3 - x_3)} - \frac{x_1(x_3 + 2y_1 + y_3) \ln \frac{x_1}{-x_3}}{y_1(y_3 - x_3)(y_1 + y_3)}, & x_1 > y_1 + y_3 \end{cases}$$

$$C_3(x_1, x_2, y_1, y_2) = \begin{cases} \frac{(x_1x_2 + y_1y_2) \ln \frac{x_2 - y_2}{x_2}}{y_1(x_2 - y_2)y_2} - \frac{x_1(x_2 + y_1) \ln \frac{-x_1}{x_2}}{y_1(x_2 - y_2)(y_1 + y_2)}, & x_1 < 0 \\ \frac{1}{x_1 - y_1} + \frac{2x_1 + x_2}{y_1(y_1 + y_2)} + \frac{[(x_1 + y_2)y_1 - x_1^2] \ln \frac{x_2 - y_2}{x_2}}{y_1(x_2 - y_2)y_2} + \frac{x_1 \ln \frac{4x_1(x_2 - y_2)p_z^2}{\mu^2}}{y_1(x_2 - y_2)} + \frac{x_1 \ln \frac{4x_1x_2p_z^2}{\mu^2}}{y_1(y_1 + y_2)}, & 0 < x_1 < y_1 \\ \frac{1}{x_2 - y_2} + \frac{x_1 + 2x_2}{y_2(y_1 + y_2)} + \frac{[(x_2 + y_1)y_2 - x_2^2] \ln \frac{x_1 - y_1}{x_1}}{(x_1 - y_1)y_1y_2} + \frac{x_2 \ln \frac{4x_2(x_1 - y_1)p_z^2}{\mu^2}}{(x_1 - y_1)y_2} + \frac{x_2 \ln \frac{4x_1x_2p_z^2}{\mu^2}}{y_2(y_1 + y_2)}, & y_1 < x_1 < y_1 + y_2 \\ \frac{(x_1x_2 + y_1y_2) \ln \frac{x_1 - y_1}{x_1}}{y_1(x_1 - y_1)y_2} - \frac{x_2(x_1 + y_2) \ln \frac{-x_2}{x_1}}{y_2(x_1 - y_1)(y_1 + y_2)}, & x_1 > y_1 + y_2. \end{cases}$$





3. Renormalization: RI/MOM



in the space-like $p^2 = -\rho(p^z)^2 < 0$ kinematics

$$(a_1 + a_2 \not{n}_t \not{n}_z + a_3 \not{n}_\perp \not{n}_z + a_4 \not{n}_\perp \not{n}_t) \tilde{S}$$

In the on-shell limit, the third and the last term a_3, a_4 disappear after integrating out the momentum q , and the product $\not{n}_t \not{n}_z$ goes to a unit matrix. Therefore, the summation $a_1 + a_2$ captures all terms that lead to UV divergences in the on-shell limit.

$$\tilde{\phi}(x_1, x_2, \mu)_{\text{RI/MOM}} = \frac{\tilde{\phi}(x_1, x_2, \mu)}{\tilde{\phi}(x_1, x_2, \mu)_{\text{OF}}}$$





3. Renormalization: RI/MOM



$$\begin{aligned} \tilde{\phi}(x_1, x_2, \mu)_{\text{OF}} &= \delta(x_1 - x_{1,0})\delta(x_2 - x_{2,0}) + \frac{\alpha_s C_F}{8\pi} \\ &\times \left\{ \left[g'_2 \delta(x_2 - x_{2,0}) + g'_3 \delta(x_3 - x_{3,0}) \right. \right. \\ &\left. \left. + \{x_2 \leftrightarrow x_1, x_{2,0} \leftrightarrow x_{1,0}\} \right] \right\}_{\oplus}, \end{aligned}$$





3. Renormalization: RI/MOM



$$\begin{aligned} C^{\mathcal{R}}(x_1, x_2, y_1, y_2, \mu) &= \delta(x_1 - y_1)\delta(x_2 - y_2) + \frac{\alpha_s C_F}{8\pi} \\ &\times \left[C'_2(x_1, x_2, y_1, y_2)\delta(x_2 - y_2) \right. \\ &+ C'_3(x_1, x_2, y_1, y_2)\delta(x_3 - y_3) \\ &\left. + \{x_1 \leftrightarrow x_2, y_1 \leftrightarrow y_2\} \right]_{\oplus}, \quad (46) \end{aligned}$$

where $C'_2 = C_2 - g'_2|_{x_{1,0} \rightarrow y_1, x_{2,0} \rightarrow y_2}$ and $C'_3 = C_3 - g'_3|_{x_{1,0} \rightarrow y_1, x_{2,0} \rightarrow y_2}$.





4. Expansion by regions



$$q^\mu = (q^+, q^-, q_\perp)$$

Expansion by regions:

- ✓ **Hard:** $q^\mu \sim (Q, Q, Q)$
- ✓ **Collinear:** $q^\mu \sim (Q, \Lambda^2/Q, \Lambda)$
- ✓ **Soft:** $q^\mu \sim (\Lambda, \Lambda, \Lambda)$

$$\tilde{\phi}_{(1/0)}^a = ig^2 \frac{C_F}{2} p^z \delta(x_2 - x_{2,0}) \int \frac{d^4 q}{(2\pi)^4} \frac{\delta(x_1 p^z - q^z - k_1^z)}{(q + k_1)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{q_\perp^2}{(k_s - q)^2 + i\epsilon}$$

$$\tilde{\phi}_{(1/0)}^d|_C = ig^2 C_F p^z \left[\delta(x_2 - x_{2,0}) \int \frac{d^4 q}{(2\pi)^4} \frac{\delta(x_1 p^z - q^z - k_1^z)}{(k_1 + q)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{2k_1^z + q^0 + q^z}{q^z} \right]_\oplus$$





4. Expansion by regions

$$q^\mu = (q^+, q^-, q_\perp)$$

Expansion by regions:

✓ **Hard:** $q^\mu \sim (Q, Q, Q)$



✓ **Collinear:** $q^\mu \sim (Q, \Lambda^2/Q, \Lambda)$

✓ **Soft:** $q^\mu \sim (\Lambda, \Lambda, \Lambda)$

$$\tilde{\phi}_{(1/0)}^a = ig^2 \frac{C_F}{2} p^z \delta(x_2 - x_{2,0}) \int \frac{d^4 q}{(2\pi)^4} \frac{\delta(x_1 p^z - q^z - k_1^z)}{(q+k_1)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{q_\perp^2}{(k_s - q)^2 + i\epsilon}$$

$$\tilde{\phi}_{(1/0)}^d|_C = ig^2 C_F p^z \left[\delta(x_2 - x_{2,0}) \int \frac{d^4 q}{(2\pi)^4} \frac{\delta(x_1 p^z - q^z - k_1^z)}{(k_1 + q)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{2k_1^z + q^0 + q^z}{q^z} \right]_\oplus$$

Leading Power!





4. Expansion by regions

✓ **Collinear:** $q^\mu \sim (Q, \Lambda^2/Q, \Lambda)$ **quasi-DA = LCDA**

$$\tilde{\phi}_{(1/0)}^a|_C = ig^2 \frac{C_F}{2} p^z \delta(x_2 - x_{2,0}) \int \frac{d^4q}{(2\pi)^4} \frac{\delta(x_1 p^z - q^z - k_1^z)}{(q+k_1)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{q_\perp^2}{(k_3 - q)^2 + i\epsilon} |_C$$

quasi

$$= ig^2 \frac{C_F}{2} \frac{p^+ - p^-}{\sqrt{2}} \delta(x_2 - x_{2,0}) \int \frac{d^4q}{(2\pi)^4} \frac{\sqrt{2} \delta((x_1 p^+ - q^+ - k_1^+) - (x_1 p^- - q^- - k_1^-))}{(q+k_1)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{q_\perp^2}{(k_3 - q)^2 + i\epsilon} |_C$$

$$= ig^2 \frac{C_F}{2} p^+ \delta(x_2 - x_{2,0}) \int \frac{d^4q}{(2\pi)^4} \frac{\delta(x_1 p^+ - q^+ - k_1^+)}{(q+k_1)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{q_\perp^2}{(k_3 - q)^2 + i\epsilon}$$

= $\phi_{(1/0)}^a$ · **Light-cone**

$$\tilde{\phi}_{(1/0)}^d|_C = ig^2 C_F p^z \left[\delta(x_2 - x_{2,0}) \int \frac{d^4q}{(2\pi)^4} \frac{\delta(x_1 p^z - q^z - k_1^z)}{(k_1 + q)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{2k_1^z + q^0 + q^z}{q^z} \right] |_C$$

quasi



$$= ig^2 C_F \frac{p^+}{\sqrt{2}} \left[\delta(x_2 - x_{2,0}) \int \frac{d^4q}{(2\pi)^4} \frac{\delta(\frac{x_1 p^+}{\sqrt{2}} - \frac{q^+ - q^-}{\sqrt{2}} - \frac{k_1^+}{\sqrt{2}})}{(k_1 + q)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{2\frac{k_1^+}{\sqrt{2}} + \frac{q^+ + q^-}{\sqrt{2}} + \frac{q^+ - q^-}{\sqrt{2}}}{\frac{q^+ - q^-}{\sqrt{2}}} \right] |_C$$

$$= ig^2 C_F \frac{p^+}{\sqrt{2}} \left[\delta(x_2 - x_{2,0}) \int \frac{d^4q}{(2\pi)^4} \frac{\sqrt{2} \delta(x_1 p^+ - q^+ - k_1^+)}{(k_1 + q)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{2k_1^+ + q^+ + q^+}{q^+} \right] |_C$$

$$= ig^2 C_F p^+ \left[\delta(x_2 - x_{2,0}) \int \frac{d^4q}{(2\pi)^4} \frac{\delta[(x_1 - x_{1,0}) p^+ - q^+]}{(k_1 + q)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{2(k_1^+ + q^+)}{q^+} \right] |_C$$

= $\phi_{(1/0)}^d$ · **Light-cone**

Leading Power!





4. Expansion by regions

$$q^\mu = (q^+, q^-, q_\perp)$$



Expansion by regions:

- ✓ Hard: $q^\mu \sim (Q, Q, Q)$
- ✓ Collinear: $q^\mu \sim (Q, \Lambda^2/Q, \Lambda)$
- ✓ Soft: $q^\mu \sim (\Lambda, \Lambda, \Lambda)$

$$\tilde{\phi}_{(1/0)}^a = ig^2 \frac{C_F}{2} p^z \delta(x_2 - x_{2,0}) \int \frac{d^4 q}{(2\pi)^4} \frac{\delta(x_1 p^z - q^z - k_1^z)}{(q + k_1)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{q_\perp^2}{(k_s - q)^2 + i\epsilon}$$

$$\tilde{\phi}_{(1/0)}^d|_C = ig^2 C_F p^z \left[\delta(x_2 - x_{2,0}) \int \frac{d^4 q}{(2\pi)^4} \frac{\delta(x_1 p^z - q^z - k_1^z)}{(k_1 + q)^2 + i\epsilon} \frac{1}{q^2 + i\epsilon} \frac{2k_1^z + q^0 + q^z}{q^z} \right]_\oplus$$

$\mathcal{O}(\Lambda_{\text{QCD}}/Q)$ **Power suppressed!**





4. Expansion by regions

- The one-loop LCDA and quasi-DA for baryon Λ does not contain the soft contributions.
- The one-loop quasi-DA contain the collinear and hard mode.
- QCD factorization shows that the hard and collinear modes in the quasi-DA can be factorized into a convolution of the hard matching coefficient and the LCDA which only contains collinear modes.

Hard Mode + Collinear Mode

Collinear Mode

$$\tilde{\Phi}(x_1, x_2, \mu) = \int dy_1 dy_2 \underbrace{\mathcal{C}(x_1, x_2, y_1, y_2, \mu)}_{\text{Hard Mode}} \underbrace{\Phi(y_1, y_2, \mu)}_{\text{Collinear Mode}} + \mathcal{O}\left(\frac{1}{x_1 p^z}, \frac{1}{x_2 p^z}, \frac{1}{(1-x_1-x_2)p^z}\right)$$





5. Summary and outlook



- LCDAs of a light baryon can be obtained through a simulation of a quasi-DA calculable on lattice QCD under the LaMET.
- We have calculated the one-loop perturbative contributions to LCDA and quasi-DA and explicitly have **demonstrated the factorization of quasi-DA at the one-loop level.**
- Our result **provides a first step to obtaining the baryon LCDA from first principle** lattice QCD calculations ***in the future.***

See Jun Hua's slides!

Thanks

