

# BOREXINO and the Sun




Marco Giammarchi


*Istituto Nazionale di Fisica Nucleare – Milano*  
<http://pcgiammarchi.mi.infn.it/giammarchi>

*On behalf of the BOREXINO Collaboration*

- 1 Sun and Solar Models
- 2 BOREXINO (in a nutshell)
- 3 BOREXINO Solar Physics results
- 4 Implications for other stars



# Sun and Solar Models



**William Thomson  
(Lord Kelvin)**

Energy from Gravity


Age of the Sun  
less than 20.000.000 years

A longstanding debate (19th c.)

**What is the energy source of the  
Sun and of the stars in general?**



**Unknown source of energy**




**Charles Darwin**


Geology and Biology

Earth is older than  
300.000.000 years

# Nuclear Physics for the Sun (1938)




Carl F. von Weizsäcker




Hans Bethe

Energy from nuclear reactions: **hydrogen burning** through **pp chain** and **CNO cycle**  
**... the Sun is 5.000.000.000 years old!**


# pp chain (99%)




# Solar Neutrinos



# CNO cycle (1%)



Dominant in the sun  
 $T_{core} \sim 15 \times 10^6 K$




- Dominant in heavier stars (>1.3 Sun mass)

Detection of neutrinos to confirm the working principles of the Sun (R. Davis experiment)

# Solar Neutrino Spectrum: experiment and predictions


- Davis experiment
- Solar Models (J. Bahcall)
- Discrepancy between the Davis experiment and the Model (Solar Neutrino Problem)
- Astrophysics/Neutrino physics possible solutions
- Neutrino Oscillations! (2002)



Predicted Neutrino Spectrum from the Sun, according to the Standard Solar Model

# BOREXINO


## Detection reaction: neutrino-electron elastic scattering



**Elastic Scattering**

**All types (NC)**


$$\nu_x + e^- \rightarrow \nu_x + e^-$$



**Electron neutrinos (NC, CC)**

$$\nu_e + e^- \rightarrow \nu_e + e^-$$

# The BOREXINO detector



# The BOREXINO saga


**1990:** idea of a sub-Mev solar neutrino detector  
A real time neutrino detection

**1995:** CTF testing the record radiopurity  
 $^{238}\text{U}, ^{232}\text{Th} < 10^{-16} \text{ g/g}$  &  $^{14}\text{C}/^{12}\text{C} < 10^{-18}$

**1996-1997:** Approval of the experiment



**Mid-2007:** Beginning of the data taking

Radiopure materials (structure and scintillator)  
Purification: distillation,  $\text{N}_2$  stripping, water extraction.






# BOREXINO at the Laboratori Nazionali del Gran Sasso




# Solar Physics Results

## Solar Neutrino Spectrum



## Reconstructed electron recoil spectrum and backgrounds in BOREXINO



- Reconstruction of  $\sim$ all components and backgrounds in the scintillator fiducial volume  
see e.g. M. Agostini et al. *Comprehensive measurement of pp-chain solar neutrinos* Nature 562 (2018) 505
- Identification and measurement of the **pp** main component (first observation made in real-time)
- Identification and measurement of the **CNO** neutrino contribution (first time)


## Summary of the BOREXINO Solar results (see also talk by M. Chen)

Neutrinos	References	Rate [cpd/100t]	Flux [cm <sup>-2</sup> s <sup>-1</sup> ]
pp	Nature 2014, Nature 2018, PRD 2019	$(134 \pm 10)_{-10}^{+6}$	$(6.1 \pm 0.5)_{-0.5}^{+0.3} \times 10^{10}$
<sup>7</sup> Be	PLB 2008, PRL 2011, Nature 2018, PRD 2019	$(48.3 \pm 1.1)_{0.7}^{+0.4}$	$(4.99 \pm 0.11)_{-0.08}^{+0.06} \times 10^9$
pep	PRL 2012, Nature 2018 PRD 2019	$(2.65 \pm 0.36)_{-0.24}^{+0.15}$ [HZ]	$(1.27 \pm 0.19)_{-0.12}^{+0.08} \times 10^8$ [HZ]
<sup>8</sup> B	PRD 2010, Nature 2018, PRD 2020	$0.223_{-0.022}^{+0.021}$	$5.68_{-0.41-0.03}^{+0.39+0.03} \times 10^6$
hep	Nature 2018, PRD 2020	<0.002 (90% CL)	<1.8x10 <sup>5</sup> (90% CL)
<b>CNO</b>	<b>Nature 2020</b>	<b><math>6.7_{-0.8}^{+2.0}</math></b>	<b><math>6.6_{-0.9}^{+2.0} \times 10^8</math></b>



# Implications for Main Sequence Stars

- Fuse Hydrogen to form Helium in their cores
- They are about 90% of the Stars
- Masses from  $\sim 0.2 M_{\odot}$  (nuclear fusion limit) to  $\sim 200 M_{\odot}$  (Eddington limit)
- Surface temperatures of
  - 50,000 K for  $100 M_{\odot}$
  - 30,000 K for  $18 M_{\odot}$
  - 5,800 K for  $1 M_{\odot}$
  - 3,600 K for  $0.5 M_{\odot}$
- Lifetime of
  - $90 \times 10^9$  y for  $0.5 M_{\odot}$
  - $10 \times 10^9$  y for  $1 M_{\odot}$
  - $20 \times 10^6$  y for  $10 M_{\odot}$
- Mass-Luminosity relations

Main Sequence: the most abundant stars in the Universe



All Main Sequence stars call for the same Physics!





BOREXINO measurements are relevant to all Main Sequence stars

(22,000 stars in the Hipparchos catalog)

BOREXINO measurements are relevant to all Main Sequence stars

BOREXINO error on the CNO measurement  $\sim 30\%$  (favoring high metallicity)

Error on C,N,O astrophysical reactions  $\sim 20\%$   
And decreasing because of new measurements planned (as in LUNA and other experiments)



Different role of heat transport mechanisms, depending on the T gradient (should not affect the conclusions!)

Fundamental step to the understanding and the experimental demonstration of the working principle of all stars in the Main Sequence

# Grazie!



**Premio  
Cocconi  
2021 - EPS**



**Premio Pontecorvo  
2015 G. Bellini**



**Premio Fermi  
2017 G. Bellini**



Ministerstwo  
Edukacji i Nauki

**Award of  
Polish Science  
Minister 2022**



**Bethe Prize  
2023 F. Calaprice**



Thank you for your attention



## Backup Slides

