

(Very/Ultra) High Energy Astrophysics II – Gamma Ray Astronomy


Mathieu de Naurois

LLR- In2p3/CNRS – Ecole Polytechnique – France

denauroi@in2p3.fr


- High Energy Gamma-Ray Astronomy from Space
- Atmospheric Showers
- Very High Energy Gamma-Ray Astronomy from ground

High Energy γ -ray Astronomy (From Space)



Pair Creation Telescopes

- ❑ Atmosphere is opaque to gamma rays
- ❑ Conversion of γ into pair $e^+ e^-$
 - ❑ Threshold $E_\gamma > 2 m_e c^2$ (1.022 MeV)
 - ❑ Angular resolution better as energy increase
 - $\theta \sim 1.5^\circ$ @ 100 MeV
 - $\theta \sim 0.1^\circ$ > 10 GeV
 - ❑ Almost no deviation for $E_\gamma \gg 2 m_e c^2$,
- ❑ $e^+ e^-$ reconstructed in a tracker \Rightarrow incident γ ray
- ❑ Anti-coincidence shield against charged cosmic rays



History of pair creation telescopes

1967-1968, OSO-3

621 γ ,
Galactic Plane

1972-1973, SAS-2,

~8,000 γ ,
3 sources

1975-1982, COS-B

~200,000 γ ,
25 sources (3C 273)


1991-2000, EGRET

$>1.4 \times 10^6 \gamma$,
271 sources


2007- AGILE

2008- FERMI


OSO-3





SAS-2



COS-B





EGRET




FERMI Large Area Telescope

- High precision tracker
 - 18 X/Y planes, Si strips (228 μm)
 - 900 000 channels
 - Triggers on 3 X/Y planes
- Hodoscopic Calorimeter
 - 1536 CsI(Tl) crystals(8 layers)
 - Shower imaging capabilities
- Anti-coïncidence shield
 - Segmented to avoid self-veto,
 - 89 folds



Fermi LAT > 100 MeV

□ 3×10^7 photons, $\langle E \rangle = 800$ MeV, $\Delta E/E = 100\%$




4FGL Catalogue

□ Release February 2019

<https://arxiv.org/abs/1902.10045>

- 5064 sources
- 232 pulsars (with pulsations)
- 17 PWNs
- 40 SNRs
- 11 binaries
- ~ 3130 blazars
- 42 radio galaxies
- 7 starburst galaxies
- 15 globular clusters
- 1336 unassociated



□ Updated June 22th, 2022
(4FGL - DR3)

<https://arxiv.org/abs/2201.11184>

□ 6658 sources

Pulsars


□ Compact object (neutron star) highly magnetized, rapidly rotating:

□ $R \sim 10 \text{ km}$

□ $M \sim 1.4 M_{\odot}$


□ Period $\text{ms} \Rightarrow \text{s}$

□ Several possible acceleration regions




Gamma-ray pulsars

- Gamma-ray pulsar: seen only in GeV.
- EGRET: A single gamma-ray pulsar (Geminga) out of 8 pulsars
- Fermi: discovery of a full population (16 in 6 month)
- Strong constraints on the geometry of emission zone (cone opening angle vs energy)



Pulsar catalog




- Population growing rapidly (> 240 in 2018)
- Millisecond pulsars: re-accelerated (in binary systems)
- Cutoff energy measurement, ...

Active Galactic Nuclei



- ❑ Supermassive black hole (10^6 - $10^9 M_{\odot}$) surrounded by accretion disk
- ❑ Giant ultra-relativistic jets of plasma (size \sim Mpc)

Artist's view (ESA/NASA)





Blazars

- Active Galactic Nuclei
 - Supermassive black hole surrounded by an accretion disk
 - Ultrarelativistic jets (Mpc)
 - Blazars: jets pointing towards the earth
 - Highly variable TeV emission: two model classes:
 - Leptonic
 - Hadronic (through π^0 decay)
 - Possible connection with UHECRS



Blazars

- Good quality MWL data
- Sequence of blazars emerging
- Population studies
- Tests of models (correlations $X/\gamma, \dots$)




Gamma-Ray bursts

- ❑ Most violent, and most distant explosions in Universe, lasting ~ few seconds to few minutes, observed ~ every day
- ❑ 2 envisaged mechanisms
 - ❑ Coalescence of neutron stars (Kilonova)
 - ❑ Collapse of very massive star (aka Collapsar)




NASA Astrophysics

Gamma-ray bursts




- ❑ Short burst of radiation (0.1 → 100 s)
- ❑ Binary Mergers
- ❑ “Hypernova”
- ❑ Several hundreds of GRB
 - ❑ A dozen seen in LAT
 - ❑ Time resolved spectra
 - ❑ Prompt emission > 10 GeV




13.6 GeV

Very High Energy γ -ray Astronomy




Atmospheric Showers

- ❑ Interaction of primary particle with a nucleus in the atmosphere (~ 10 km altitude)
- ❑ Creation of pairs (electron & positrons) & bremsstrahlung processes redistribute the energy to a large number of particles




Atmospheric Showers

- Primary interaction in the high atmosphere
 - Shower of secondary particles, detectable up to a few 100 m
 - Large effective areas ($>10^5 \text{ m}^2$, increasing with energy)
 - ⇒ High energies ($\geq 1000 \text{ TeV}$)
- Atmosphere used as an inhomogeneous calorimeter.
- Observables: charged particles, Cerenkov light, fluorescence light, radio emission
- Reconstruction: Energy, direction, impact, nature of primary particle (γ -hadron ; light (p, He) – heavy nuclei (Fe))




Atmospheric Showers

γ , 200 GeV




Atmospheric Showers




- Atmospheric showers initiated by protons and nuclei fluctuate more, and contain penetrating particles (muons)

Cherenkov Light

- ❑ As in any transparent medium (e.g. water), light is a bit **slower** in air than in vacuum
- ❑ Relativistic particles in the shower travel **faster** than light **in air**
- ❑ They emit Cherenkov light, analogue for light of sound wall



Sound Wall



- When a plane flies faster than speed of sound, sound emitted at each time accumulate on a cone → **Sonic Boom**

Observables and experimental techniques ($E > 10$ GeV)

TIBET



Primary (γ , hadron)

Scintillator
or water

Atmospheric
Shower

Hadron
detector

μ

Primary (γ , hadron)

Atmospheric
Shower

Cherenkov
Emission

Cherenkov
Telescope



MILAGRO

HESS, VERITAS, MAGIC, CANGAROO




STACEE, CELESTE, SOLAR II, GRAAL



2 Complementary Techniques


- Atmospheric Cherenkov Telescopes:
 - Small F.O.V.
 - Low duty cycle
 - High rejection
 - High resolution

Detailed study of a few sources




- Sampling experiments (Water Cherenkov, Particle Arrays,...)
 - Large F.O.V.
 - High duty cycle
 - Poor rejection
 - Poor resolution



Long term survey instruments




Atmospheric Cherenkov Telescopes



- Cherenkov light-pool ~ 120 m
- Image the shower on a fast camera ($\Delta T \sim 2$ ns)
- Large effective area (10^5 m²) even with modest reflector



- Key parameter : speed (< 10 ns)
- Image shape used in discrimination



VHE γ -ray world



High Energy Stereoscopic System



□ H.E.S.S. phase 1 (2003):

- 4 telescopes: $\text{Ø } 12 \text{ m}, 107 \text{ m}^2$
- 960 pixels/camera, FOV : 5°
- Observations : $\sim 1000 \text{ h/an}$
- Localisation accuracy : $\sim 10''$

□ H.E.S.S. phase 2 (2012):

- 5th telescope, $\text{Ø } 28 \text{ m}, 600 \text{ m}^2$
- 2048 pixels, FOV : 3.5°
- Pointing in $< 1 \text{ min}$ for 50 % sky
- Extended energy range

High Energy Stereoscopic System


©Sabine Gloaguen



Sabine Gloaguen
PHOTOGRAPHY

The Milky way in Gamma-rays

Major HESS project: 2700 h of data (2004 – 2010)








78 sources of different types (SNRs, PWNs, Binaries, ...)

VHE Universe much richer than expected

Galactic Bestiary in a nutshell

A complete zoo :

- ❑ Supernova Remnants
- ❑ Galactic Centre
- ❑ Pulsar Wind Nebular
- ❑ Interacting Binary Systems
- ❑ Young, Energetic Pulsars
- ❑ Diffuse Galactic Emission



Supernova Remnant


Artist's view (ESA/Nasa)

- ❑ Remnant of the explosion of a massive star
- ❑ Shock wave (> 1000 km/s) sweeping up interstellar medium
- ❑ Thermal emission from hot material inside the shell in X rays (10^6 K)
- ❑ Place of heavy element nucleosynthesis
→ at the origin of life!
- ❑ Enormous Released energy $\sim 10^{44}$ J
($10^{16} \times$ Hiroshima)
- ❑ Best candidates for galactic cosmic rays
 - ❑ Matching rate 1 SNR/Galaxy/century
 - ❑ Conversion efficiency $\sim 10\%$




State of the art

- ❑ SNRs are the second population of VHE sources in the Milky Way
- ❑ Morphologies in X/ γ -rays are often very similar, suggesting that the same particles give rise to the two emissions
- ❑ Maximum acceleration energy around 100 TeV




Supernova Remnant Evolution



Binary Systems

Artist's view (ESA/NASA)



- ❑ 70% of stars in the Galaxy live in couples of 2 or more!
- ❑ Binary systems are formed of :
 - ❑ A compact object (black hole or neutron star)
 - ❑ A companion star
 - ❑ An accretion disk
 - ❑ Sometimes jets of plasma
- ❑ Complex physics, variable systems
- ❑ 5 VHE gamma-ray systems known.



*Tatooine
(Star War)*



LS 5039 : light is opaque to light !

- ❑ Compact binary system with tight orbit (0.1 A.U) and massive (20 M_{\odot}), UV (17 000 K) star
- ❑ γ -rays are absorbed by pair creation on the stellar photons ($\gamma + h_{\nu} \rightarrow e^{+} + e^{-}$)
- ❑ Geometry introduces a modulation of VHE emission



Blazars


- Active Galactic Nuclei, seen face-on
- Jets outshine completely the galaxy & accretion disk
- Very variable emission, on short time scales (< min)
- Can be used to check that the speed of light is really a constant (not depending on light energy)



$$c' = c \left(1 \pm \xi \frac{E}{E_P} \pm \zeta^2 \frac{E^2}{E_P^2} \right)$$

TeV Gamma-ray bursts

- Two recent detections :
 - GRB 180721B, observed 10 hours after the initial burst, one of the most intense ever seen
 - GRB 190829A, seen in TeV during 2 days
 - In both case, “remnant phase” ↔ interaction of plasma jet with interstellar medium
- GRBs are very efficient particle accelerators!
- More recently a monster GRB (GRB 221009A) was seen up to ~ 18 TeV !







Results – Many Others

□ Wealth of results:


- Supernova remnants
- Plerions
- Galactic Center
- Binary Systems
- Interacting Stellar Winds
- Starburst galaxies
- Galactic Novae
- Huge flares from blazars,
- Tests of Lorentz Invariance
- Indirect dark matter searches,
-

Unveiling the
sources of cosmic
rays!



HAWC

- ❑ High altitude Water Cherenkov detector
- ❑ Array of water tanks
 - ❑ 900 tanks each 4.3 m high and 5 m diameter 1 PMT at bottom
 - ❑ 150 m x 150 m array
 - ❑ 75% ground coverage
- ❑ Sierra Negra, Mexico, alt: 4100 m lat: +19°



HAWC



Advantages:

- Wide field of view
- 100% duty cycle

Drawbacks


- Sensitivity much lower than IACTs
- Angular resolution $\sim 1^\circ$
- Bad energy reconstruction

Complementary to IACTs, starting to produce results



Conclusions

- Gamma-ray astronomy (from ground & space) underwent a revolution in the last ~ 15 years
 - New window in the Universe opened: violent phenomena, sources of cosmic rays
 - H.E.S.S. in Namibia leader of the field
 - Lots of unexpected results (binary systems, millisecond pulsars, ..;)
- Major project to come in the field: Cherenkov Telescope Array



Cherenkov Telescope Array




One night with HESS



Backup

Events



- ❑ These are the actual “photos” taken by the telescopes
- ❑ We then need to infer the particle type, its energy & direction to generate maps of the sky