

Lévy α -stable generalization of the ReBB model

based on [Universe 2023, 9\(8\), 361](#) & [Universe 2024, 10\(3\), 127](#)

T. Csörgő^{1,2}, S. Hegyi², [I. Szanyi](#)^{1,2,3}

¹MATE KRC, Gyöngyös, Hungary

²Wigner RCP, Budapest, Hungary

³Eötvös University, Budapest, Hungary

19th International Scientific Days
26 June 2024, Gyöngyös, Hungary

Outline

- **preliminaries**
- **the $p=(q,d)$ Bialas-Bzdak model and its extended version**
- **motivation for Lévy α -stable generalization**
- **Lévy α -stable generalization of the Bialas-Bzdak model**
- **an approximate simple Lévy α -stable model and fits to data**
- **relation between the parameters of the simple Lévy α -stable model and the full generalized model**

Preliminaries: ReBB model analysis of pp and p \bar{p} data

- the Real extended Bialas-Bzdak (ReBB) model describes elastic pp and p \bar{p} $d\sigma/dt$ data in a statistically acceptable way (CL \geq 0.1%) in the kinematic region:


$$546 \text{ GeV} \leq \sqrt{s} \leq 8 \text{ TeV}$$

$$0.38 \text{ GeV}^2 \leq -t \leq 1.2 \text{ GeV}^2$$

- significant model dependent odderon signal is observed
- main goal:** to improve the ReBB model to have a statistically acceptable (CL \geq 0.1%) description to elastic pp and p \bar{p} data in a wider kinematic range

T. Csörgő, I. Szanyi, *Eur. Phys. J. C* **81**, 611 (2021)

I. Szanyi, T. Csörgő, *Eur. Phys. J. C* **82**, 827 (2022)



ReBB model description to the 8 TeV pp data

Unitarity and the elastic scattering amplitude

- the unitarity of the S -matrix expresses the conservation of probability

$$SS^\dagger = I$$

- the unitarity relation in impact parameter (b) representation at high energies is

$$\tilde{\sigma}_{in}(s, b) = 2 \operatorname{Im} t_{el}(s, b) - |t_{el}(s, b)|^2 \quad (\sqrt{s} \text{ is the CM energy})$$

$$0 \leq \tilde{\sigma}_{in}(s, b) \leq 1 \quad \leftarrow \text{unitarity constraint}$$

- the elastic scattering amplitude $t_{el}(s, b)$ can be written as a solution of the unitarity equation in terms of $\tilde{\sigma}_{in}(s, b)$ (inelastic cross section / probability of inelastic scattering)
- $\tilde{\sigma}_{in}(s, b)$ can be calculated by using probability calculus and R. J. Glauber's multiple diffractive scattering theory


The Bialas-Bzdak (BB) $p=(q,d)$ model

A. Bialas, A. Bzdak, *Acta Phys.Polon. B 38, 159-168 (2007)*

- in the **Bialas-Bzdak (BB) $p=(q,d)$ model** the proton is a **bound state of a constituent quark and constituent a diquark**
- the inelastic scattering probability of two protons at a fixed impact parameter vector (\vec{b}) and at fixed constituent transverse position vectors ($\vec{s}_q, \vec{s}_d, \vec{s}'_q, \vec{s}'_d$) is given by a **Glauber expansion**:

$$\sigma(\vec{s}_q, \vec{s}_d; \vec{s}'_q, \vec{s}'_d; \vec{b}) = 1 - \prod_{a \in \{q,d\}} \prod_{b \in \{q,d\}} [1 - \sigma_{ab}(\vec{b} + \vec{s}'_b - \vec{s}_a)]$$

- $\sigma_{ab}(\vec{x}) \equiv \frac{d^2 \sigma_{ab}(\vec{x})}{dx^2}$ is the inelastic differential cross section (inelastic scattering probability) for the collision of two constituents at a fixed relative transverse position \vec{x} of the constituents
- the **Glauber expansion sums the probabilities of all possible single and multiple binary inelastic collisions of the constituents (back scattering is prohibited)**
- the collision of two protons is inelastic if at least one constituent-constituent collision is inelastic



Proton-proton collision in the quark-diquark model

- the **probability of inelastic scattering** of protons at a fixed impact parameter vector (\vec{b}) is given by averaging over the constituent positions inside the protons:

$$\tilde{\sigma}_{in}(\vec{b}) = \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} d^2 s_q d^2 s'_q d^2 s_d d^2 s'_d D(\vec{s}_q, \vec{s}_d) D(\vec{s}'_q, \vec{s}'_d) \sigma(\vec{s}_q, \vec{s}_d; \vec{s}'_q, \vec{s}'_d; \vec{b})$$

- $D(\vec{s}_q, \vec{s}_d)$ is the (transverse) distribution of constituents inside a proton
- the scattering amplitude in the original BB model is considered to be completely imaginary by neglecting its relatively small real part

$$\tilde{t}_{el}(s, b) = i \left(1 - \sqrt{1 - \tilde{\sigma}_{in}(s, b)} \right)$$

$$b = |\vec{b}|$$

$$T(s, t) = 2\pi \int_0^{\infty} \tilde{t}_{el}(s, b) J_0(qb) b db$$

$$q = \sqrt{-t}$$

- the s -dependence of the amplitude happens through the s -dependencies of the model parameters

The Bialas-Bzdak (BB) $p=(q,d)$ model

A. Bialas, A. Bzdak, *Acta Phys.Polon.*
B 38, 159-168 (2007)

- in the original BB model the distribution of constituents inside a proton is given in terms of products of Gaussians

$$D(\vec{s}_q, \vec{s}_d) = \frac{1 + \lambda^2}{R_{qd}^2 \pi} e^{-\frac{\vec{s}_q^2}{R_{qd}^2}} e^{-\frac{\vec{s}_d^2}{R_{qd}^2}} \delta^2(\vec{s}_d + \lambda \vec{s}_q)$$

$$\lambda = \frac{m_q}{m_d}$$

$$\vec{s}_d = -\lambda \vec{s}_q$$

$$\vec{s}'_d = -\lambda \vec{s}'_q$$

- the constituent-constituent inelastic differential cross sections have also Gaussian shapes

$$\sigma_{ab}(\vec{x}) = A_{ab} e^{-\frac{\vec{x}^2}{R_a^2 + R_b^2}}$$

$$a, b \in \{q, d\}$$

- the constituent-constituent inelastic integrated cross sections are

$$\sigma_{ab}^{int} = \iint \sigma_{ab}(\vec{x}) d^2x = \pi A_{ab} (R_a^2 + R_b^2)$$

- assuming that the diquark contains twice as many partons than the quark and the colliding constituents do not shadow each other, $\sigma_{qq}^{int} : \sigma_{qd}^{int} : \sigma_{dd}^{int} = 1 : 2 : 4$ and the number of free parameters reduces to five: A_{qq} , λ , R_q , R_d , and R_{qd}

Real extended Bialas-Bzdak (ReBB) model

- in the original BB model the differential cross section is zero around the position of the diffractive minimum
- as a solution, the elastic scattering amplitude was extended in a unitary manner leading to the Real extended Bialas-Bzdak (ReBB) model

F. Nemes, T. Csörgő, M. Csanád, Int. J. Mod. Phys. A Vol. 30, 1550076 (2015)

$$\begin{array}{l} \tilde{t}_{el}(s, b) = i[1 - e^{-\Omega(s, b)}] \\ \text{Re}\Omega(s, b) = -1/2 \ln[1 - \tilde{\sigma}_{in}(s, b)] \\ \text{Im}\Omega(s, b) = 0 \end{array} \quad \longrightarrow \quad \begin{array}{l} \text{new free parameter} \\ \text{Im}\Omega(s, \vec{b}) = -\alpha_R \tilde{\sigma}_{in}(s, \vec{b}) \end{array}$$

$$\tilde{t}_{el}(s, b) = i \left(1 - \sqrt{1 - \tilde{\sigma}_{in}(s, b)} \right) \quad \longrightarrow \quad \tilde{t}_{el}(s, b) = i \left(1 - e^{i \alpha_R \tilde{\sigma}_{in}(s, b)} \sqrt{1 - \tilde{\sigma}_{in}(s, b)} \right)$$

- the ReBB model gives a statistically acceptable description (CL \geq 0.1%) to elastic pp and p \bar{p} scattering in the kinematic region:

$$0.546 \text{ TeV} \leq \sqrt{s} \leq 8 \text{ TeV} \quad \& \quad 0.38 \text{ GeV}^2 \leq -t \leq 1.2 \text{ GeV}^2$$

T. Csörgő, I. Szanyi, Eur. Phys. J. C 81, 611 (2021)

I. Szanyi, T. Csörgő, Eur. Phys. J. C 82, 827 (2022)

- the main goal is to have a statistically acceptable description in a wider kinematic range
- the TOTEM measurement at LHC at $\sqrt{s} = 8$ TeV excluded a purely exponential pp differential cross-section in the range of four-momentum transfer squared $0.027 \text{ GeV}^2 \leq -t \leq 0.2 \text{ GeV}^2$ with a significance greater than 7σ TOTEM Collab., *Nucl. Phys. B*, 899, 527 (2015)
- a simple model with Gaussian impact parameter amplitude yields a purely exponential t-distribution while a simple model with Levy α -stable impact parameter amplitude and $\alpha_L < 2$ yields a non-exponential t-distribution

$$\tilde{T}_{el}(s, b) = \frac{i + \rho_0(s)}{2} \sigma_{tot}(s) \frac{1}{2\pi B_0(s)} e^{-\frac{1}{2} \frac{b^2}{B_0(s)}} \quad \rightarrow \quad \tilde{T}_{el}(s, b) = \frac{i + \rho_0(s)}{2} \sigma_{tot}(s) \frac{1}{4\pi^2} \int d^2 q e^{-i\vec{q} \cdot \vec{b}} e^{-\frac{1}{2} |q^2 B_L(s)|^{\alpha_L(s)/2}}$$

$$T_{el}(s, t) = \int d^2 b e^{i\vec{q} \cdot \vec{b}} \tilde{T}_{el}(s, b) \quad \frac{d\sigma_{el}}{dt}(s, t) = \frac{1}{4\pi} |T_{el}(s, t)|^2 \quad a(s) = \frac{1 + \rho_0^2(s)}{16\pi} \sigma_{tot}^2(s)$$

$$\frac{d\sigma_{el}}{dt}(s, -t) = a(s) e^{-t B_0(s)} \quad \rightarrow \quad \frac{d\sigma_{el}}{dt}(s, -t) = a(s) e^{-|t B_L(s)|^{\alpha_L(s)/2}}$$


Lévy α -stable distributions in HEP

- the application of Lévy α -stable distributions is not new in the field of high-energy physics
- the Cauchy-Lorentz or Breit-Wigner distribution ($\alpha_L = 1$ case) is used to model unstable particles
- the Lévy expansion technique was applied to describe elastic pp scattering also at 13 TeV

T. Csörgő, R. Pasechnik, A. Ster, *Eur. Phys. J. C* 79, 62 (2019)

- the application of stable distributions is widespread in heavy ion physics

M. Csanád, D. Kincses, *Universe* 10 (2024) 2, 54



Description to pp elastic differential cross section data at 13 TeV using the Lévy expansion technique

Gaussian vs Lévy α -stable distribution

- the bivariate Gaussian distribution centered at $\vec{0}$ is

$$G(\vec{x}|R_G) = \frac{1}{2\pi R_G^2} e^{-\frac{\vec{x}^2}{2R_G^2}}$$


- the bivariate symmetric Levy α -stable distribution centered at $\vec{0}$ is

$$L(\vec{x}|\alpha_L, R_L) = \frac{1}{(2\pi)^2} \int d^2 q e^{-i\vec{q}\cdot\vec{x}} e^{-|\vec{q}^2 R_L^2|^{\alpha_L/2}}$$

$$0 < \alpha_L \leq 2$$

- for $\alpha_L = 2$ the the Lévy α -stable distribution is the Gaussian distribution

$$L(\vec{x}|\alpha_L = 2, R_L = R_G/\sqrt{2}) \equiv G(\vec{x}|R_G)$$



The bivariate symmetric Levy α -stable distribution for $R_L = 1$ as a function of $x = |x|$

Lévy α -stable distributions with $\alpha_L < 2$ have tails behaving asymptotically as a power law (infinite variance)

- the inelastic differential cross section for the collision of two constituents can be written **in terms of a convolution of their parton distributions**
- in the original BB model the parton distributions of the constituents are Gaussian distributions


$$\sigma_{ab}(\vec{x}) = A_{ab} \pi S_{ab}^2 \int d^2 r_a G(\vec{r}_a | R_a / \sqrt{2}) G(\vec{x} - \vec{r}_a | R_b / \sqrt{2})$$

$$\equiv A_{ab} \pi S_{ab}^2 G(\vec{x} | S_{ab} / \sqrt{2})$$

$$\vec{x} = \vec{b} + \vec{s}'_b - \vec{s}_a$$

$$S_{ab}^2 = R_a^2 + R_b^2$$

$$a, b \in \{q, d\}$$



The picture of the proton in the quark-diquark model


The quark-diquark distribution

- in the original BB model $D(\vec{s}_q, \vec{s}_d)$, the distribution of constituents inside a proton is given in terms of products of Gaussians
- **considering the relative distance** between the quark and diquark ($\vec{s}_q - \vec{s}_d$) one can write $D(\vec{s}_q, \vec{s}_d)$ in terms of a single Gaussian distribution:

$$D(\vec{s}_q, \vec{s}_d) = (1 + \lambda)^2 G(\vec{s}_q - \vec{s}_d | R_{qd}/\sqrt{2}) \delta^2(\vec{s}_q + \lambda \vec{s}_d)$$

$$\lambda = m_q/m_d$$

- the Dirac δ fixes the center of the mass of the proton making the calculations easier
- $D(\vec{s}_q, \vec{s}_d)$ is normalized as $\int d^2 s_q d^2 s_d D(\vec{s}_q, \vec{s}_d) = 1$



The picture of the proton in the quark-diquark model

Lévy α -stable generalized Bialas-Bzdak (LBB) model

Universe 2023, 9(8), 361

- **the parton distributions of the constituent quark and diquark are now Levy α -stable distributions** and the inelastic differential cross section for the collision of two constituents is:

$$\sigma_{ab}(\vec{x}) = A_{ab}\pi S_{ab}^2 \int d^2r_a L(\vec{r}_a|\alpha_L, R_a/2)L(\vec{x} - \vec{r}_a|\alpha_L, R_b/2) \equiv A_{ab}\pi S_{ab}^2 L(\vec{x}|\alpha_L, S_{ab}/2)$$

$$S_{ab}^{\alpha_L} = R_a^{\alpha_L} + R_b^{\alpha_L}$$

- **the distribution of the constituents inside the proton is now given in terms of a Levy α -stable distribution:**

$$D(\vec{s}_q, \vec{s}_d) = (1 + \lambda)^2 L(\vec{s}_q - \vec{s}_d|\alpha_L, R_{qd}/2)\delta^2(\vec{s}_q + \lambda\vec{s}_d)$$

$$\int d^2s_q d^2s_d D(\vec{s}_q, \vec{s}_d) = 1$$

α_L is a new free parameter of the model and if $\alpha_L = 2$ the BB model with Gaussian distributions is recovered

Difficulties with LBB model

- $\tilde{\sigma}_{in}(\vec{b})$ can be written as sum of 11 different terms that are integrals of products of Lévy α -stable distributions

$$\tilde{\sigma}_{in}(\vec{b}) = \tilde{\sigma}_{in}^{qq}(\vec{b}) + 2\tilde{\sigma}_{in}^{qd}(\vec{b}) + \tilde{\sigma}_{in}^{dd}(\vec{b}) - [2\tilde{\sigma}_{in}^{qq,qd}(\vec{b}) + \tilde{\sigma}_{in}^{qd,dq}(\vec{b}) + \tilde{\sigma}_{in}^{qq,dd}(\vec{b}) + 2\tilde{\sigma}_{in}^{qd,dd}(\vec{b})] \\ + [\tilde{\sigma}_{in}^{qq,qd,dq}(\vec{b}) + 2\tilde{\sigma}_{in}^{qq,qd,dd}(\vec{b}) + \tilde{\sigma}_{in}^{dd,qd,dq}(\vec{b})] - \tilde{\sigma}_{in}^{qq,qd,dq,dd}(\vec{b})$$

- difficulties with the calculation of integrals of products of Lévy α -stable distributions
- the calculation is easy only if the integral can be written in a convolution form as in case of the leading order terms in $\tilde{\sigma}_{in}(s, \vec{b})$

Leading order terms in $\tilde{\sigma}_{in}$ in the LBB model

Universe 2023, 9(8), 361

$$\begin{aligned}\tilde{\sigma}_{in}^{qq}(\vec{b}) &= \pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} \times \\ &\times \int d^2 s_q d^2 s'_q L(\vec{s}_q | \alpha_L, R_{qd^*}/2) L(\vec{s}'_q | R_{qd^*}/2) L(\vec{b} + \vec{s}'_q - \vec{s}_q | (2R_q^{\alpha_L})^{1/\alpha_L}/2) \\ &= \pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} L(\vec{b} | \alpha_L, (2R_{qd^*}^{\alpha_L} + 2R_q^{\alpha_L})^{1/\alpha_L}/2),\end{aligned}$$

$$\begin{aligned}\tilde{\sigma}_{in}^{qd}(\vec{b}) &= 2\pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} \times \\ &\times \int d^2 s_q d^2 s'_q L(\vec{s}_q | R_{qd^*}/2) L(\vec{s}'_q | R_{qd^*}/2) L(\vec{b} - \lambda \vec{s}'_q - \vec{s}_q | \alpha_L, (R_q^{\alpha_L} + R_d^{\alpha_L})^{1/\alpha_L}/2) \\ &= 2\pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} L(\vec{b} | \alpha_L, ((1 + \lambda^{\alpha_L})R_{qd^*}^{\alpha_L} + R_q^{\alpha_L} + R_d^{\alpha_L})^{1/\alpha_L}/2),\end{aligned}$$

$$\begin{aligned}\tilde{\sigma}_{in}^{dd}(\vec{b}) &= 4\pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} \times \\ &\times \int d^2 s_q d^2 s'_q L(\vec{s}_q | R_{qd^*}/2) L(\vec{s}'_q | R_{qd^*}/2) L(\vec{b} + \lambda(\vec{s}_q - \vec{s}'_q) | \alpha_L, (2R_d^{\alpha_L})^{1/\alpha_L}/2) \\ &= 4\pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} L(\vec{b} | \alpha_L, (2\lambda^{\alpha_L} R_{qd^*}^{\alpha_L} + 2R_d^{\alpha_L})^{1/\alpha_L}/2).\end{aligned}$$

Difficulties with LBB model fits to the data

- since multivariate Lévy α -stable distributions can be given only in terms of special functions, it is hard to perform a numerical fitting procedure
- a relatively high computing capacity and improved analytic insight is needed to proceed with the full model
- **quick solution:** approximations that are valid at the low $-t$ domain
- at low $-t$ values, the original ReBB model had difficulties to describe the strongly non-exponential features of the experimental data on $d\sigma/dt$
- a simple model which is valid at the low $-t$ domain easily illustrates the power of the Lévy α -stable generalization

Simple Lévy α -stable model for low- $|t|$ pp $d\sigma/dt$

Universe 2023, 9(8), 361

- low- $|t|$ scattering corresponds to high- b scattering and at high b values $\tilde{\sigma}_{in}(s, b)$ is small
- leading order term in the Taylor expansion of the amplitude in $\tilde{\sigma}_{in}(s, b)$ dominates at low $-t$ values if α_R is small too

$$\tilde{t}_{el}(s, b) = i \left(1 - e^{i \alpha_R(s) \tilde{\sigma}_{in}(s, b)} \sqrt{1 - \tilde{\sigma}_{in}(s, b)} \right) \longrightarrow \tilde{t}_{el}(s, b) = \left(\alpha_R(s) + \frac{i}{2} \right) \tilde{\sigma}_{in}(s, b)$$

- motivated by the fact that the leading order terms in $\tilde{\sigma}_{in}(s, \vec{b})$ have Lévy α -stable shapes in the LBB model, $\tilde{\sigma}_{in}(s, \vec{b})$ is approximated with a single Lévy α -stable shape

$$\tilde{\sigma}_{in}(s, \vec{b}) = \tilde{c}(s) L(\vec{b} | \alpha_L(s), r(s))$$



- **a simple Lévy α -stable model model for low- $|t|$ pp $d\sigma/dt$ arises**

$$t_{el}(s, t) = \int d^2 b e^{i\vec{q}\cdot\vec{b}} \tilde{t}_{el}(s, \vec{b}), |\vec{\Delta}| = \sqrt{-t} \longrightarrow \frac{d\sigma}{dt}(s, t) = \frac{1}{4\pi} |t_{el}(s, t)|^2 = a(s) e^{-|tb(s)|^{\alpha_L(s)/2}}$$

- the model has three adjustable parameters, α_L , a , and b , to be determined at a given energy

Simple Lévy α -stable model and the data

Universe 2023, 9(8), 361



- the non-exponential Lévy α -stable model with $\alpha_L = 1.953 \pm 0.004$ represents the LHC TOTEM $\sqrt{s} = 8 \text{ TeV}$ low- $|t|$ differential cross section data with a confidence level of 55% (published)
- similarly good description is obtained to all the LHC data on low- $|t|$ pp (and $p\bar{p}$) $d\sigma/dt$

Fits with simple Lévy α -stable model

Universe 2024, 10(3), 127

- fits to the existing pp and p \bar{p} $d\sigma/dt$ data in the kinematic range:

$$546 \text{ GeV} \leq \sqrt{s} \leq 13 \text{ TeV}$$

$$0.02 \text{ GeV}^2 \leq -t \leq 0.15 \text{ GeV}^2$$



- the CL values of the fits range between 8.8% and 96%.
- statistical, systematic and normalization errors are taken into account using the χ^2 definition developed by PHENIX Collab.

\sqrt{s} , GeV	α_L	a , mb/GeV ²	b , GeV ⁻²	CL, %
546	1.93 ± 0.09	209 ± 15	15.8 ± 0.9	18.1
1800	2.0 ± 1.5	270 ± 24	16.2 ± 0.2	77.1
2760	1.600 ± 0.3	637 ± 252	28 ± 11	20.5
7000 (T)	1.95 ± 0.01	535 ± 30	20.5 ± 0.2	8.8
7000 (A)	1.97 ± 0.01	463 ± 13	19.8 ± 0.2	96.0
8000 (T1)	1.955 ± 0.005	566 ± 31	20.09 ± 0.08	43.86
8000 (T2)	1.90 ± 0.03	582 ± 33	20.9 ± 0.4	19.6
8000 (A)	1.97 ± 0.01	480 ± 11	19.9 ± 0.1	55.8
13000 (T1)	1.959 ± 0.006	677 ± 36	20.99 ± 0.08	76.5
13000 (T2)	1.958 ± 0.003	648 ± 10	21.06 ± 0.05	89.1
13000 (A)	1.968 ± 0.006	569 ± 17	20.84 ± 0.07	29.7

Values of the fitted parameters of the simple Lévy- α stable model at different energies

A. Adare *et al.* (PHENIX Collab.) *Phys. Rev. C* 77, 064907

$\alpha_L = 2$ versus $\alpha_L < 2$ results @ 13 TeV




Energy dependence of the α_L parameter

Universe 2024, 10(3), 127

- the value of the α_L parameter does not depend on energy
- its value is 1.959 ± 0.002 , i.e., slightly but in a statistical sense significantly different from 2

→ strong non-exponential behavior at low $-t$ in the differential cross section, power law tail at high- \vec{b} in $\tilde{\sigma}_{in}(s, \vec{b})$



Energy dependence of the α_L parameter of the simple Lévy- α stable model

Energy dependence of the optical point parameter


Universe 2024, 10(3), 127

- the energy dependence of the a parameter is quadratically logarithmic:

$$a(s) = p_0 + p_1 \ln \frac{s}{1 \text{ GeV}^2} + p_2 \ln^2 \frac{s}{1 \text{ GeV}^2}$$

- ATLAS and TOTEM data result slightly different energy dependencies
- reason: ATLAS and TOTEM use different methods to obtain the absolute normalization of the measurements

ATLAS Collab., *Eur. Phys. J. C* 83 (2023) 441



Energy dependence of the a parameter of the simple Lévy- α stable model

Energy dependence of the slope parameter


Universe 2024, 10(3), 127

- the energy dependence of the b parameter for TOTEM and ATLAS data together, and for $p\bar{p}$ data alone are linearly logarithmic:

$$b(s) = p_0 + p_1 \ln \frac{s}{1 \text{ GeV}^2}$$

- the LHC pp and the lower energy $p\bar{p}$ data do not lie on the same curve
- reason: the slope parameter data have a jump in the energy dependence around 3-4 GeV

TOTEM Collab., *Eur. Phys. J. C* (2019) 79:103



Energy dependence of the b parameter of the simple Lévy- α stable model

Simple Lévy α -stable & LBB model parameters

- parameters of the simple Levy α -stable model and the measurable quantities at $t \rightarrow 0$ can be approximately expressed in terms of the parameters of the LBB model *Universe 2023, 9(8), 361*
- only leading order terms in $\tilde{\sigma}_{in}(s, \vec{b})$ are considered; $A_{qq} = 1$ and $\lambda = 1/2$ are fixed

$$\frac{d\sigma}{dt}(s, t = 0) = a(s) = \frac{81}{16} \pi \left(2R_q^{\alpha_L(s)}(s) \right)^{4/\alpha_L} (1 + 4\alpha_R^2(s))$$

$$b(s) = \frac{1}{36} \left(\frac{4}{3} \right)^{2/\alpha_L(s)} \left((2 + 2^{\alpha_L(s)}) R_{qd}^{\alpha_L(s)}(s) + 3^{\alpha_L(s)} \left(2R_d^{\alpha_L(s)}(s) + R_q^{\alpha_L(s)}(s) \right) \right)^{2/\alpha_L(s)}$$

(obtained after a Taylor expansion in $t^{\alpha_L/2}$)

$$\sigma_{tot}(s) = 9\pi \left(2R_q^{\alpha_L(s)}(s) \right)^{2/\alpha_L(s)}$$

$$\rho_0(s) = \frac{Ret_{el}(s, t = 0)}{Imt_{el}(s, t = 0)} = 2\alpha_R$$

$$\sigma_{el}(s) = \frac{a(s)}{b(s)} \Gamma \left(\frac{2 + \alpha_L(s)}{\alpha_L(s)} \right)$$

- according to the analysis of elastic pp and $p\bar{p}$ data in the energy region $0.5 \text{ TeV} \leq \sqrt{s} \leq 8 \text{ TeV}$ only α_R is different for pp and $p\bar{p}$ scattering (T. Csörgő, I. Szanyi, *Eur. Phys. J. C* **81**, 611 (2021))
- in the low- $|t|$ approximation, difference between pp and $p\bar{p}$ scattering could be seen in the data on $d\sigma/dt$, ρ_0 , a (optical point), and σ_{el} , no difference in the data on σ_{tot} and b

Summary

- the formal Lévy α -stable generalization of the Bialas-Bzdak model is done, the $\alpha_L = 2$ limit corresponds to the original model
- solution of difficult and complex technical (mathematical and computational) problems is needed to fit the experimental data with the generalized model
- based on approximations a highly simplified Levy α -stable model of the pp (and $p\bar{p}$) differential cross section is deduced and successfully fitted to the data in the low- $|t|$ region
- the energy dependences of the parameters of the simple model are determined; the parameters of the simple model are related to the parameters of the Lévy α -stable generalized Real extended Bialas-Bzdak (LBB) model
- final conclusion: the successful fit results indicate promising prospect for the future utility of the LBB model in describing experimental data



Thank you for your attention!

Backup slides

ρ_0 & α_R : connection between $t = 0$ and $t \neq 0$ data

- there is a connection between the ρ_0 parameter and the α_R parameter of the ReBB model regulating the real part of the scattering amplitude and the minimum-maximum structure of the $d\sigma/dt$
- α_R is determined by the $d\sigma/dt$ data at the minimum-maximum region but at the same time specifies the value of the ρ_0 in the ReBB model

T. Csörgő, I. Szanyi, *Eur. Phys. J. C* **81**, 611 (2021)



Most general term in $\tilde{\sigma}_{in}$

$$\tilde{\sigma}_{in}^{qq,qd,dq,dd}(\vec{b}) = \int d^2 s_q d^2 s'_q L(\vec{s}_q | R_{qd^*}/2) L(\vec{s}'_q | R_{qd^*}/2) \times \sigma_{qq}(\vec{s}_q, \vec{s}'_q; \vec{b}) \sigma_{qd}(\vec{s}_q, -\lambda \vec{s}'_q; \vec{b}) \sigma_{dq}(\vec{s}'_q, -\lambda \vec{s}_q; \vec{b}) \sigma_{dd}(-\lambda \vec{s}_q, -\lambda \vec{s}'_q; \vec{b})$$

$$\sigma_{qq}(\vec{s}_q, \vec{s}'_q; \vec{b}) = \pi A_{qq} (2R_q^\alpha)^{2/\alpha} \times L(\vec{b} + \vec{s}'_q - \vec{s}_q | \alpha, (2R_q^\alpha)^{1/\alpha} / \sqrt{2})$$



$$\sigma_{qd}(\vec{s}_q, \vec{s}'_d; \vec{b}) = 2\pi A_{qq} (2R_q^\alpha)^{2/\alpha} \times L(\vec{b} + \vec{s}'_d - \vec{s}_q | \alpha, (R_q^\alpha + R_d^\alpha)^{1/\alpha} / \sqrt{2})$$

$$\sigma_{dd}(\vec{s}_d, \vec{s}'_d; \vec{b}) = 4\pi A_{qq} (2R_q^\alpha)^{2/\alpha} \times L(\vec{b} + \vec{s}'_d - \vec{s}_d | \alpha, (2R_d^\alpha)^{1/\alpha} / 2)$$

$$\sigma_{dq}(\vec{s}_d, \vec{s}'_q; \vec{b}) = 2\pi A_{qq} (2R_q^\alpha)^{2/\alpha} \times L(\vec{b} + \vec{s}'_q - \vec{s}_d | \alpha, (R_q^\alpha + R_d^\alpha)^{1/\alpha} / 2)$$

Energy dependences of the ReBB model parameters

T. Csörgő, I. Szanyi, *Eur. Phys. J. C* 81, 611 (2021)





The energy dependences of the scale parameters, $R_q(s)$, $R_d(s)$, and $R_{qd}(s)$ are **linear logarithmic** and the **same** for pp and $p\bar{p}$ processes!

The energy dependence of the α parameter, $\alpha(s)$ is **linear logarithmic** too, but **not** the same for pp and $p\bar{p}$ processes!

Energy dependences of the ReBB model parameters

T. Csörgő, I. Szanyi, *Eur. Phys. J. C* 81, 611 (2021)



The energy dependences of the scale parameters, $R_q(s)$, $R_d(s)$, and $R_{qd}(s)$ are linear logarithmic and the same for pp and $p\bar{p}$ processes!

The energy dependence of the α parameter, $\alpha(s)$ is linear logarithmic too, but not the same for pp and $p\bar{p}$ processes!

Fit method

- least squares fitting with the method developed by the PHENIX collaboration
- this method is **equivalent to the diagonalization of the covariance matrix** if the experimental errors are separated into three different types:
 - type A: point-to-point varying uncorrelated statistical and systematic errors
 - type B: point-to-point varying 100% correlated systematic errors
 - type C: point-independent, overall systematic uncertainties
- i.e least squares fitting with:

[A. Adare et al. \(PHENIX Collab.\)
Phys. Rev. C 77, 064907](#)

$$\chi^2 = \left(\sum_{j=1}^M \left(\sum_{i=1}^{n_j} \frac{(d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj} - th_{ij})^2}{\tilde{\sigma}_{ij}^2} \right) + \epsilon_{bj}^2 + \epsilon_{cj}^2 \right) + \left(\frac{d_{\sigma_{tot}} - th_{\sigma_{tot}}}{\delta\sigma_{tot}} \right)^2 + \left(\frac{d_{\rho_0} - th_{\rho_0}}{\delta\rho_0} \right)^2$$

$$\tilde{\sigma}_{ij}^2 = \tilde{\sigma}_{aij} \left(\frac{d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj}}{d_{ij}} \right)$$

$$\tilde{\sigma}_{kij} = \sqrt{\sigma_{kij}^2 + (d'_{ij}\delta_k t_{ij})^2}, \quad k \in \{a, b\}$$

- minimization with **CERN Root MINUIT**, parameter error estimation by **MINOS**.

Fit method

- the method takes into account (in M separately measured t ranges):
 - the t -dependent statistical (**type A**) and systematic (**type B**) errors (both vertical σ_k and horizontal $\delta_k t$) $\rightarrow \epsilon_b$ parameters;
 - the t -independent σ_c normalization uncertainties (**type C**) $\rightarrow \epsilon_c$ parameters;
 - the measured total cross-section $d_{\sigma_{tot}}$ and ratio d_{ρ_0} and their total uncertainties $\delta\sigma_{tot}$ and $\delta\rho_0$.

[A. Adare et al. \(PHENIX Collab.\)
Phys. Rev. C 77, 064907](#)

- i.e least squares fitting with:**

$$\chi^2 = \left(\sum_{j=1}^M \left(\sum_{i=1}^{n_j} \frac{(d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj} - th_{ij})^2}{\tilde{\sigma}_{ij}^2} \right) + \epsilon_{bj}^2 + \epsilon_{cj}^2 \right) + \left(\frac{d_{\sigma_{tot}} - th_{\sigma_{tot}}}{\delta\sigma_{tot}} \right)^2 + \left(\frac{d_{\rho_0} - th_{\rho_0}}{\delta\rho_0} \right)^2$$

$$\tilde{\sigma}_{ij}^2 = \tilde{\sigma}_{aij} \left(\frac{d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj}}{d_{ij}} \right)$$

$$\tilde{\sigma}_{kij} = \sqrt{\sigma_{kij}^2 + (d'_{ij}\delta_k t_{ij})^2}, \quad k \in \{a, b\}$$

- minimization with **CERN Root MINUIT**, parameter error estimation by **MINOS**.

Fit method

- the method takes into account (in M separately measured t ranges):
 - the ϵ_i -s must be considered as both measurements and fit parameters not effecting the NDF (since they have known central value of zero and known standard deviation of one)
 - the measured total cross-section $d_{\sigma_{tot}}$ and ratio d_{ρ_0} and their total uncertainties $\delta\sigma_{tot}$ and $\delta\rho_0$.

[A. Adare et al. \(PHENIX Collab.\)
Phys. Rev. C 77, 064907](#)

- i.e least squares fitting with:

$$\chi^2 = \left(\sum_{j=1}^M \left(\sum_{i=1}^{n_j} \frac{(d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj} - th_{ij})^2}{\tilde{\sigma}_{ij}^2} + \epsilon_{bj}^2 + \epsilon_{cj}^2 \right) + \left(\frac{d_{\sigma_{tot}} - th_{\sigma_{tot}}}{\delta\sigma_{tot}} \right)^2 + \left(\frac{d_{\rho_0} - th_{\rho_0}}{\delta\rho_0} \right)^2 \right)$$

$$\tilde{\sigma}_{ij}^2 = \tilde{\sigma}_{aij} \left(\frac{d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj}}{d_{ij}} \right)$$

$$\tilde{\sigma}_{kij} = \sqrt{\sigma_{kij}^2 + (d'_{ij}\delta_k t_{ij})^2}, \quad k \in \{a, b\}$$

- minimization with **CERN Root MINUIT**, parameter error estimation by **MINOS**.

Fit method

The PHENIX method is validated by evaluating the χ^2 from a full covariance matrix fit of the $\sqrt{s} = 13$ TeV TOTEM differential cross-section data using the Lévy expansion method from [T. Csörgő, R. Pasechnik, & A. Ster, Eur. Phys. J. C 79, 62 \(2019\)](#).

- the t -independent σ_c normalization uncertainties $\rightarrow \epsilon_c$ parameters;
- the measured total cross-section $d_{\sigma_{tot}}$ and ratio d_{ρ_0} and their total uncertainties $\delta\sigma_{tot}$ and $\delta\rho_0$.

[A. Adare et al. \(PHENIX Collab.\)
Phys. Rev. C 77, 064907](#)

- i.e least squares fitting with:

$$\chi^2 = \left(\sum_{j=1}^M \left(\sum_{i=1}^{n_j} \frac{(d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj} - th_{ij})^2}{\tilde{\sigma}_{ij}^2} \right) + \epsilon_{bj}^2 + \epsilon_{cj}^2 \right) + \left(\frac{d_{\sigma_{tot}} - th_{\sigma_{tot}}}{\delta\sigma_{tot}} \right)^2 + \left(\frac{d_{\rho_0} - th_{\rho_0}}{\delta\rho_0} \right)^2$$

$$\tilde{\sigma}_{ij}^2 = \tilde{\sigma}_{aij} \left(\frac{d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj}}{d_{ij}} \right)$$

$$\tilde{\sigma}_{kij} = \sqrt{\sigma_{kij}^2 + (d'_{ij}\delta_k t_{ij})^2}, \quad k \in \{a, b\}$$

- minimization with **CERN Root MINUIT**, parameter error estimation by **MINOS**.

Fit method

The PHENIX method is validated by evaluating the χ^2 from a full covariance matrix fit of the $\sqrt{s} = 13$ TeV TOTEM differential cross-section data using the Lévy expansion method from [T. Csörgő, R. Pasechnik, & A. Ster, Eur. Phys. J. C 79, 62 \(2019\)](#).

The PHENIX method and the fit with the full covariance matrix result in the same minimum within one standard deviation of the fit parameters.

[A. Adare et al. \(PHENIX Collab.\)
Phys. Rev. C 77, 064907](#)

- i.e least squares fitting with:

$$\chi^2 = \left(\sum_{j=1}^M \left(\sum_{i=1}^{n_j} \frac{(d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj} - th_{ij})^2}{\tilde{\sigma}_{ij}^2} \right) + \epsilon_{bj}^2 + \epsilon_{cj}^2 \right) + \left(\frac{d_{\sigma_{tot}} - th_{\sigma_{tot}}}{\delta\sigma_{tot}} \right)^2 + \left(\frac{d_{\rho_0} - th_{\rho_0}}{\delta\rho_0} \right)^2$$

$$\tilde{\sigma}_{ij}^2 = \tilde{\sigma}_{aij} \left(\frac{d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj}}{d_{ij}} \right)$$

$$\tilde{\sigma}_{kij} = \sqrt{\sigma_{kij}^2 + (d'_{ij}\delta_k t_{ij})^2}, \quad k \in \{a, b\}$$

- minimization with **CERN Root MINUIT**, parameter error estimation by **MINOS**.

Proportionality between $\rho_0(s)$ and $\alpha(s)$

$$t_{el}(s, b) = i \left(1 - e^{i \alpha \tilde{\sigma}_{in}(s, b)} \sqrt{1 - \tilde{\sigma}_{in}(s, b)} \right)$$

$$\alpha \tilde{\sigma}_{in} \ll 1$$


$$\text{Im } t_{el}(s, b) \simeq \lambda(s) \exp\left(-\frac{b^2}{2R^2(s)}\right)$$



$$\rho_0(s) = \alpha(s) \left(2 - \frac{3}{2} \lambda(s) + \frac{1}{3} \lambda^2(s) \right)$$

$$\lambda(s) = \text{Im } t_{el}(s, b = 0)$$

→ by rescaling one can get additional α parameter values at energies where ρ_0 is measured (and vice versa)



The dependence of ρ_0/α on $\lambda = \text{Im } t_{el}(s, b = 0)$ in the TeV energy range. The data points are generated numerically by using the trends of the ReBB model scale parameters and the experimentally measured ρ -parameter values.

Measurable quantities

- differential cross section:

$$\frac{d\sigma}{dt}(s, t) = \frac{1}{4\pi} |T(s, t)|^2$$

- total, elastic and inelastic cross sections:

$$\sigma_{tot}(s) = 2\text{Im}T(s, t=0)$$

$$\sigma_{el}(s) = \int_{-\infty}^0 \frac{d\sigma(s, t)}{dt} dt$$

$$\sigma_{in}(s) = \sigma_{tot}(s) - \sigma_{el}(s)$$

- ratio ρ_0 :

$$\rho_0(s) = \lim_{t \rightarrow 0} \rho(s, t) \equiv \frac{\text{Re}T(s, t \rightarrow 0)}{\text{Im}T(s, t \rightarrow 0)}$$

- slope of $d\sigma/dt$:

$$B(s, t) = \frac{d}{dt} \left(\ln \frac{d\sigma}{dt}(s, t) \right)$$

$$B_0(s) = \lim_{t \rightarrow 0} B(s, t)$$