## **Deep Inelastic Scattering 2025**



Contribution ID: 60 Type: not specified

## New measurement of $K^+ \to \pi^+ \nu \bar{\nu}$ branching ratio at the NA62 experiment

Wednesday 26 March 2025 09:00 (20 minutes)

The  $K^+ \to \pi^+ \nu \bar{\nu}$  decay is a golden mode for flavour physics. Its branching ratio is predicted with high precision by the Standard Model to be less than  $10^{-10}$ , and this decay mode is highly sensitive to indirect effects of new physics up to the highest mass scales. A new measurement of the  $K^+ \to \pi^+ \nu \bar{\nu}$  decay by the NA62 experiment at the CERN SPS is presented, using data collected in 2021 and 2022. This new dataset was collected after modifications to the beamline and detectors and at a higher instantaneous beam intensity with respect to the previous 2016–2018 data taking. Using the NA62 datasets from 2016–2022, a new measurement of  $\mathcal{B}(K^+ \to \pi^+ \nu \bar{\nu}) = \left(13.0^{+3.3}_{-2.9}\right) \times 10^{-11}$  is reported, and for the first time the  $K^+ \to \pi^+ \nu \bar{\nu}$  decay is observed with a significance exceeding  $5\sigma$ .

Author: FIORENZA, Renato (INFN Napoli)

Presenter: FIORENZA, Renato (INFN Napoli)

Session Classification: WG3: Electroweak Physics and Beyond the Standard Model

Track Classification: Electroweak Physics and Beyond the Standard Model