

Deep Inelastic Scattering 2025



Contribution ID: 49

Type: **not specified**

ATLAS ITk Pixel Detector Overview

Tuesday 25 March 2025 10:06 (22 minutes)

In the high-luminosity era of the Large Hadron Collider, the instantaneous luminosity is expected to reach unprecedented values, resulting in up to 200 proton-proton interactions in a typical bunch crossing. To cope with the resulting increase in occupancy, bandwidth and radiation damage, the ATLAS Inner Detector will be replaced by an all-silicon system, the Inner Tracker (ITk). The innermost part of the ITk will consist of a pixel detector, with an active area of about 13 m^2 . To deal with the changing requirements in terms of radiation hardness, power dissipation and production yield, several silicon sensor technologies equipped with novel ASICs connecting by bump-bonding technique will be employed in the five barrel and endcap layers. As a timeline, it is facing to pre-production of components, sensor, building modules, mechanical structures and services.

Authors: CRESCIOLI, Francesco (Univ. P. et Marie Curie (Paris VI) (FR)); CRESCIOLI, Francesco (Centre National de la Recherche Scientifique (FR))

Presenters: CRESCIOLI, Francesco (Univ. P. et Marie Curie (Paris VI) (FR)); CRESCIOLI, Francesco (Centre National de la Recherche Scientifique (FR))

Session Classification: WG6: Future Experiments

Track Classification: Future Experiments