
















# Looking for strange particles in ALICE




## Today's periodic system of the fundamental building blocks

	<i>Quarks</i>		<i>Leptons</i>	
<i>Generation 3</i>	 <b>t</b> Top	 <b>b</b> Bottom	 <b>τ</b> Tau	 <b>ν<sub>τ</sub></b> Tau-neutrino
<i>Generation 2</i>	 <b>c</b> Charm	 <b>s</b> Strange	 <b>μ</b> Muon	 <b>ν<sub>μ</sub></b> Muon-neutrino
<i>Generation 1</i>	 <b>u</b> Up	 <b>d</b> Down	 <b>e</b> Electron	 <b>ν<sub>e</sub></b> Electron-neutrino


# What are strange particles ?

meson



$\bar{d}s, ds$

baryon



uds

hadrons (baryons or mesons) containing at least one strange (s) quark

We will be looking for **neutral** strange particles, which travel **some distance (mm or cm) from the point of production (collision point)** before they decay into **two oppositely charged particles**

$$K_s^0 \rightarrow \pi^+ \pi^- \quad \tau = 0.89 \times 10^{-10} \text{ s}$$

$$c\tau = 3 \times 10^{10} \text{ cm s}^{-1} \times 8.9 \times 10^{-11} \text{ s}$$

2.67 cm from the point of interaction


$$\Lambda \rightarrow \pi^- p \quad \tau = 2.6 \times 10^{-10} \text{ s}$$

$$c\tau = 3 \times 10^{10} \text{ cm s}^{-1} \times 2.6 \times 10^{-10} \text{ s}$$



7.2 cm distance from the point of interaction

$$\bar{\Lambda} \rightarrow \pi^+ \bar{p}$$

Weak decays : strangeness is not conserved




## How do we find V0s ?










We look for two opposite tracks, having the same origin, which is not the interaction (collision) point

# How do we find V0s ?



We look for two opposite tracks, having the same origin, which is not the interaction (collision) point

# How do we identify each V0?




V0 decay :  
a neutral particle (no track) gives suddenly two tracks

- $P = Q \cdot B \cdot R$   
 P momentum  
 Q electric charge  
 B magnetic field  
 R radius of curvature

**Identify V0s from the decay topology**

# How do we identify each V0?



## Calculate the (invariant) mass

Energy conservation

$$E = E_1 + E_2$$

Momentum conservation

$$\mathbf{p} = \mathbf{p}_1 + \mathbf{p}_2$$

Total energy

$$E^2 = p^2 c^2 + m^2 c^4$$

$c=1$

$$E^2 = p^2 + m^2$$

$$E = E_1 + E_2 \quad E_1^2 = p_1^2 + m_1^2 \quad E_2^2 = p_2^2 + m_2^2$$

$$E^2 = p^2 + m^2 \quad m^2 = E^2 - p^2 = (E_1 + E_2)^2 - (p_1 + p_2)^2 = m_1^2 + m_2^2 + 2E_1 E_2 - 2\mathbf{p}_1 \cdot \mathbf{p}_2$$

## Calculate the mass of the initial particle from the values of the mass and the momentum of the final particles

Particle Identification (done by a number of PID detectors)  $\Rightarrow m_1 m_2$

Radius of curvature of the particle tracks due to magnetic field  $\Rightarrow p_1 p_2$


$P=Q \cdot B \cdot R$  (P momentum, Q electric charge, R radius of curvature, B magnetic field)



## Strangeness enhancement in lead-lead collisions

- Analysis of large event samples from lead collisions
- Find number of  $K_s$ ,  $\Lambda$ , anti- $\Lambda$
- Calculate particle yields
- Calculate strangeness enhancement taking into account particle yields in proton collisions

# Strangeness enhancement : one of the first signals of QGP



Enhancement increases with number of strange quarks in the hadron ( $\Omega$  has 3,  $\Xi$  has 2,  $\Lambda$  has 1)





ALI-DER-80680

Particle yield for Pb-Pb collisions/ the number of participants

---

Particle yield for proton-proton collisions / 2

# Fit functions describing the invariant mass distributions




2<sup>nd</sup> degree polynomial for the background  $f(x) = ax^2 + bx + c$   
Gaussian for the peak


$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

Find the number of  $K_s$ ,  $\Lambda$ , anti- $\Lambda$  after subtraction of the background

# Geometry of a Pb-Pb collision



- Peripheral collision
  - Large **distance** between the centres of the nuclei
  - Small number of **participants**
  - Few charged particles produced (low multiplicity)





- Central collision
  - Small **distance** between the centres of the nuclei
  - Large number of **participants**
  - Many charged particles produced (high multiplicity)

# Centrality of Pb-Pb collisions


Distribution of the signal amplitude of V0 (plastic scintillators)  
 red line : described by model (Glauber)



Centrality	$dN_{ch}/d\eta$	$\langle N_{part} \rangle$	$(dN_{ch}/d\eta)/(\langle N_{part} \rangle/2)$
0%–5%	$1601 \pm 60$	$382.8 \pm 3.1$	$8.4 \pm 0.3$
5%–10%	$1294 \pm 49$	$329.7 \pm 4.6$	$7.9 \pm 0.3$
10%–20%	$966 \pm 37$	$260.5 \pm 4.4$	$7.4 \pm 0.3$
20%–30%	$649 \pm 23$	$186.4 \pm 3.9$	$7.0 \pm 0.3$
30%–40%	$426 \pm 15$	$128.9 \pm 3.3$	$6.6 \pm 0.3$
40%–50%	$261 \pm 9$	$85.0 \pm 2.6$	$6.1 \pm 0.3$
50%–60%	$149 \pm 6$	$52.8 \pm 2.0$	$5.7 \pm 0.3$
60%–70%	$76 \pm 4$	$30.0 \pm 1.3$	$5.1 \pm 0.3$
70%–80%	$35 \pm 2$	$15.8 \pm 0.6$	$4.4 \pm 0.4$



peripheral collisions



central collisions

# Strangeness enhancement calculation

**Yield** : number of particles produced per interaction =  $N_{\text{particles(produced)}}/N_{\text{events}}$

**Efficiency** =  $N_{\text{particles(measured)}}/N_{\text{particles(produced)}}^*$

**Yield** =  $N_{\text{particles(measured)}}/(\text{efficiency} \times N_{\text{events}})$

$K_s$ -Yield (pp) = 0.25 /interaction ;  $\Lambda$ -Yield(pp) = 0.0617 /interaction ;  $\langle N_{\text{part}} \rangle = 2$  for pp

**Strangeness enhancement**: the particle yield normalised by the number of participating nucleons in the collision, and divided by the yield in proton-proton collisions\*\*

\*assumption on efficiency values : to match yields in Analysis Note  
Measurement of  $K_s$  and  $\Lambda$  spectra and yields in Pb–Pb collisions at  $\sqrt{s_{NN}}=2.76$  TeV with the ALICE experiment

\*pp yields at 2.76 TeV from interpolation between 900 GeV and 7 TeV  
Analysis Note “ $K_s$ ,  $\Lambda$  and anti $\Lambda$  production in pp collisions at 7 TeV”