

# Recent results from Belle and Belle II for exotic hadrons



Youngjoon Kwon (Yonsei U.)

*on behalf of Belle & Belle II*

Dec. 10, 2024 @ BCVSPIN 2024



# Overview

- Quick intro. to Belle & Belle II


- Charmed Pentaquark searches

✓  $P_c^+ \rightarrow pJ/\psi$  in  $\Upsilon(1S)$  and  $\Upsilon(2S)$

✓  $P_{cs}(4459)^0 \rightarrow \Lambda J/\psi$  in  $\Upsilon(1S)$  and  $\Upsilon(2S)$



- Updates on  $\Upsilon(10753)$

- Closing




from Yuan & Olsen, Nature Rev. Phys. 1 (2019) no.8, 480-494

# $e^+e^-$ collision near $\Upsilon$ resonances



# Luminosities of Belle II and Belle



## Belle (1999-2010) Luminosity

$$\int \mathcal{L}_{total} dt = 1039 \text{ fb}^{-1}$$

$$\int \mathcal{L}_{\Upsilon(4S)} dt = 711 \text{ fb}^{-1}$$


$$\int \mathcal{L}_{\Upsilon(1S)} dt = 5.8 \text{ fb}^{-1}$$

$$\int \mathcal{L}_{\Upsilon(2S)} dt = 24.5 \text{ fb}^{-1}$$

$$\int \mathcal{L}_{\Upsilon(5S)} dt = 121 \text{ fb}^{-1}$$



# *Search for $P_c^+$ states in $pJ/\psi$*

# motivation



- Charmed pentaquark ( $P_c$ ) states have been discovered by LHCb.
  - $P_c(4312)^+$ ,  $P_c(4440)^+$ , and  $P_c(4457)^+$  in  $\Lambda_b \rightarrow K + pJ/\psi$
- not possible to confirm with  $e^+e^-$  B factory,
  - not enough energy to produce  $\Lambda_b$  pair
  - OTOH, deuterons are observed in  $\Upsilon(nS)$  by ARGUS, CLEO and BaBar.
- Why not then look for  $P_c$  in  $\Upsilon(nS)$ ?
- Belle has world-largest sample of  $\Upsilon(1S)$  and  $\Upsilon(2S)$ .
- We search for  $P_c^+ \rightarrow pJ/\psi$  from  $\Upsilon(1S)$  and  $\Upsilon(2S)$  at Belle.

# Analysis procedure



## Event selection


- 3 well-measured charged tracks
- Identification of  $e^\pm$ ,  $\mu^\pm$  and  $p$
- $\Lambda$  veto for  $p$  candidates
- sideband for non- $J/\psi$  bkg.




## Cut on $M^2_{\text{recoil}}(pJ/\psi)$

- to suppress non- $J/\psi$  bkg. with  $M^2_{\text{recoil}}(pJ/\psi) > 10 \text{ GeV}^2$



## Study $M(pJ/\psi)$ distributions (next page)

- in  $\Upsilon(1S)$ ,
- in  $\Upsilon(2S)$ ,
- in continuum ( $\sqrt{s} = 10.52 \text{ GeV}$ )



 for  $P_c(4312)^+$ 
 for  $P_c(4440)^+$ 
 for  $P_c(4457)^+$





- no significant  $P_c^+$  signals in any place
- major sources of systematic uncertainties
  - particle ID (2.1 %)
  - MC modeling (2.2 %, 2.8 %)
  - $N_{1S}, N_{2S}$  ( $\sim 2.2$  %)

● We set upper limits on  $P_c^+$  productions from  $\Upsilon(1,2S)$



—	$\Upsilon(1S)$ decays			$\Upsilon(2S)$ decays		
	$P_c(4312)^+$	$P_c(4440)^+$	$P_c(4457)^+$	$P_c(4312)^+$	$P_c(4440)^+$	$P_c(4457)^+$
$N_{fit}^A$	$10 \pm 8$	$14 \pm 12$	$-3 \pm 9$	$30 \pm 16$	$33 \pm 15$	$0 \pm 3$
$N_{fit}^{A,UL}$	26	37	14	52	60	6
$N_{fit}^B$	$10 \pm 8$	$12 \pm 11$	$3 \pm 9$	$29 \pm 12$	$31 \pm 15$	$0 \pm 3$
$N_{fit}^{B,UL}$	26	33	17	50	57	7
$N_{sig}^{UL}$	31	47	34	56	77	26
$\mathcal{B}^{UL} (\times 10^{-6})$	4.5	6.8	4.9	5.3	7.2	2.4

*Evidence for  $P_{cs}(4459)^0$   
in  $\Lambda J/\psi$*


# motivation and procedure

- Similar motivation as the previous paper (arXiv:2403.04340)
  - for neutral charmed pentaquark  $P_{cs}(4459)^0$  in  $\Upsilon(1S)$  and  $\Upsilon(2S)$
- We search for  $P_{cs}(4459)^0 \rightarrow \Lambda J/\psi$  from  $\Upsilon(1S)$  and  $\Upsilon(2S)$  at Belle.
  - $J/\psi \rightarrow \ell^+ \ell^-$ ,  $\Lambda \rightarrow p\pi$
  - 2D sideband for  $M_{p\pi}$  vs.  $M_{\ell^+\ell^-}$

(a)  $\Upsilon(1S)$  sample  
(b)  $\Upsilon(2S)$  sample




# $\Lambda J/\psi$ Yield



## ● Assess signal yield in $M(\Lambda J/\psi)$

- use  $M_{\Lambda J/\psi} = M_{\ell^+\ell^-p\pi} - M_{\ell^+\ell^-} - M_{p\pi} + m_{J/\psi} + m_{\Lambda}$   
to improve mass resolution  $\sigma_M$  (11.6  $\rightarrow$  2.8 MeV)
- excess seen near 4.46 GeV in both  $\Upsilon(1S)$  and  $\Upsilon(2S)$  data

# Results



## ● Signal yield of $M(\Lambda J/\psi)$

- determined by a binned max. likelihood fit, with

$$f_{\text{PDF}} = f_{\text{R}} + f_{\text{no}P_{cs}} + f_{\text{SB}}$$

- fit with fixed mass, width (from LHCb value) gives

$$N_{P_{cs}} = 19 \pm 5$$

$$\Delta(-2 \ln \mathcal{L}) = 13.01 \text{ (3.4}\sigma \text{ evidence by pseudo-experiment technique)}$$

## ● Fit result with free mass, width

$$M_{\text{R}} = 4469.5 \pm 4.1 \pm 4.1 \text{ MeV}$$

$$\Gamma_{\text{R}} = 14.3 \pm 9.2 \pm 6.3 \text{ MeV}$$

## ● Systematic uncertainty

- $\Lambda$  selection (determined by  $B^{\pm} \rightarrow K^{\pm} \Lambda \bar{\Lambda}$ )  $\sim O(5\%)$ ; BF of  $\Upsilon(2S) \rightarrow \Upsilon(1S) \sim O(6\%)$
- for  $M_{\text{R}}$ ,  $\Gamma_{\text{R}}$  parameters: fit range (2.5, 3.5 MeV), N(bins) (3.2, 5.2 MeV)

# *Updates regarding* $\Upsilon(10753)$



# Energy scan for $\Upsilon(10753)$

## ● $\Upsilon(10753)$ — a reminder


- first observed by Belle, [JHEP 10 (2019) 220] with  $5.2\sigma$
- in the energy dependence of  $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
- $\exists$  several competing interpretations
- Belle also had exotic candidates  $Z_b(10610)^\pm$ ,  $Z_b(10650)^\pm$  [PRL 108, 122001 (2012)]

## ● Belle II added scan points

- JHEP 07 (2024) 116



# Confirmation of $\Upsilon(10753)$ signal



- Left-column figures for all events
- Right-column figures for  $p(\pi\pi\mu\mu) < 0.1$  GeV to suppress events from ISR
- Red dash (---) corresponding to  $\Upsilon(nS)$



# Energy scan for $\Upsilon(10753)$

## ● $\Upsilon(10753)$ — a reminder


- first observed by Belle, [JHEP 10 (2019) 220] with  $5.2\sigma$
- in the energy dependence of  $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
- $\exists$  several competing interpretations
- Belle also had exotic candidates  $Z_b(10610)^\pm$ ,  $Z_b(10650)^\pm$  [PRL 108, 122001 (2012)]

## ● Belle II added scan points

- JHEP 07 (2024) 116
- $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$  with  $\Upsilon(nS) \rightarrow \mu^+\mu^-$
- confirms Belle results of  $\Upsilon(10753)$

	$\mathcal{R}_{\sigma(1S/2S)}^{\Upsilon(10753)}$	$\mathcal{R}_{\sigma(3S/2S)}^{\Upsilon(10753)}$
Ratio	$0.46^{+0.15}_{-0.12}$	$0.10^{+0.05}_{-0.04}$ <b>small</b>

- no signals for  $Z_b(10610)^\pm$ ,  $Z_b(10650)^\pm$



# Cross-section ratios, etc.



	$\mathcal{R}_{\sigma(1S/2S)}^{\Upsilon(10753)}$	$\mathcal{R}_{\sigma(3S/2S)}^{\Upsilon(10753)}$	$\mathcal{R}_{\sigma(1S/2S)}^{\Upsilon(5S)}$	$\mathcal{R}_{\sigma(3S/2S)}^{\Upsilon(5S)}$	$\mathcal{R}_{\sigma(1S/2S)}^{\Upsilon(6S)}$	$\mathcal{R}_{\sigma(3S/2S)}^{\Upsilon(6S)}$
Ratio	$0.46^{+0.15}_{-0.12}$	$0.10^{+0.05}_{-0.04}$	$0.45^{+0.04}_{-0.04}$	$0.32^{+0.04}_{-0.03}$	$0.64^{+0.23}_{-0.13}$	$0.41^{+0.16}_{-0.12}$

**Table 2.** Cross-section ratios at resonance peaks above the  $\Upsilon(4S)$ . Uncertainty in this table combines statistical and systematic uncertainties.

Mode	$N_{Z_{b1}}$	$N_{Z_{b1}}^{\text{UL}}$	$\sigma_{Z_{b1}}$ (pb)	$\sigma_{Z_{b1}}^{\text{UL}}$ (pb)	$N_{Z_{b2}}^{\text{UL}}$	$N_{Z_{b2}}$	$\sigma_{Z_{b2}}$ (pb)	$\sigma_{Z_{b2}}^{\text{UL}}$ (pb)
10.746 GeV								
$\pi\Upsilon(1S)$	$0.0^{+1.6}_{-0.0}$	< 4.9	$0.00^{+0.04}_{-0.00}$	< 0.13	—	—	—	—
$\pi\Upsilon(2S)$	$5.8^{+5.9}_{-4.6}$	< 13.8	$0.06^{+0.06}_{-0.05}$	< 0.14	—	—	—	—
10.805 GeV								
$\pi\Upsilon(1S)$	$2.5^{+2.4}_{-1.6}$	< 5.2	$0.21^{+0.20}_{-0.13}$	< 0.43	$0.0^{+0.7}_{-0.0}$	< 5.8	$0.00^{+0.03}_{-0.00}$	< 0.28
$\pi\Upsilon(2S)$	$5.2^{+3.8}_{-3.0}$	< 12.3	$0.15^{+0.11}_{-0.09}$	< 0.35	$0.0^{+0.8}_{-0.0}$	< 6.0	$0.00^{+0.04}_{-0.00}$	< 0.30




**Table 3.** Signal yields and upper limits at 90% credibility for  $e^+e^- \rightarrow \pi Z_b(10610, 10650)$ ,  $Z_b(10610, 10650) \rightarrow \pi\Upsilon(1S, 2S)$  processes and corresponding Born cross-section measurement limits. Uncertainties for the numbers of signal events are statistical only. Here we use  $Z_{b1}$  and  $Z_{b2}$  as shorthand for  $Z_b(10610)$  and  $Z_b(10650)$ , respectively.

# Di-pion mass distribution for $\Upsilon(10753)$



- similar to both phase-space model and  $\Upsilon(2S) \rightarrow \pi^+\pi^-\Upsilon(1S)$  for  $\pi^+\pi^-\Upsilon(1S)$
- but similar to  $\Upsilon(2S) \rightarrow \pi^+\pi^-\Upsilon(1S)$  only for  $\pi^+\pi^-\Upsilon(2S)$

# $\Upsilon(10753) \rightarrow \chi_{bJ}\omega$





- cross section shows a peak at  $\Upsilon(10753)$ , hence a confirmation and a new decay channel
- the ratio  $\chi_{b1}\omega/\pi\pi\Upsilon(nS) \sim$  one order of magnitude higher at  $\Upsilon(10753)$  than at  $\Upsilon(5S)$

# $\Upsilon(10753) \rightarrow \chi_{b0}\omega$ and $\eta_b\omega$

- Tetraquark interpretation of this state predicts enhancement of  $\Upsilon(10753) \rightarrow \eta_b(1S)\omega$

$$\frac{\Gamma(\omega\eta_b)}{\Gamma(\Upsilon\pi^+\pi^-)} \sim 30$$

- we measure  $\eta_b$  indirectly by using recoil mass  $M_{\text{recoil}}(\omega) = \sqrt{(E_{\text{cm}} - E_\omega)^2 - p_\omega^2}$
- no signals observed in either modes  $\rightarrow$  set upper limits



$$\sigma_B(e^+e^- \rightarrow \eta_b(1S)\omega) < 2.5 \text{ pb},$$

$$\sigma_B(e^+e^- \rightarrow \chi_{b0}(1P)\omega) < 8.7 \text{ pb}.$$

# Summary

- As a B-factory, Belle II continues being a strong player in the study of exotic hadrons as well as spectroscopy of conventional ones.
- In this talk, we present the searches of charmed pentaquark states by Belle
  - Search for  $P_c^+ \rightarrow pJ/\psi$  in  $\Upsilon(1S)$  and  $\Upsilon(2S)$
  - Evidence of  $P_{cs}(4459)^0 \rightarrow \Lambda J/\psi$  in  $\Upsilon(1S)$  and  $\Upsilon(2S)$
- We also show Belle II results regarding  $\Upsilon(10753)$ , a new  $b\bar{b}$ -like state first observed by Belle in 2019.
  - Confirmation of  $\Upsilon(10753)$
  - New decays channel  $\Upsilon(10753) \rightarrow \chi_{bJ}\omega$
- Run 2 is underway with goal of collecting a several  $\text{ab}^{-1}$  data in the next few years. Please stay tuned!

***Thank you!***