

# Searches for rare/BSM top decays (Tevatron)



Daniel Wicke  
Bergische Universität Wuppertal  
*for the CDF and DØ collaborations*




# Introduction and Outline

Top quark decay by modified interactions or new particles:

- Elm interaction → Charge,  $t\bar{t}\gamma$
- Weak interaction →  $Wtb$ ,  $Ztq$  and Higgs
- BSM particles → Charged Higgs
- Strong interaction → Colour flow

go from rare to BSM to strong force



# Top Quark Width

Significant BSM contributions should alter the top quark width (SM: 1.34 GeV)


## Direct measurements

CDF

- Reconstructed top quark mass distribution  $4.3 \text{ fb}^{-1} \rightarrow \Gamma_t < 7.5 \text{ GeV}$

## Combination of Branching Fraction with $t$ -channel Single Top

DØ



Top pairs: amount of 0,1 and 2 identified  $b$ -Jets

$$R = \frac{B(t \rightarrow Wb)}{B(t \rightarrow Wq)} = 0.97 \pm 0.09$$

Single Top:

$\Gamma_t = \Gamma_t(t \rightarrow Wb)/R$  with  $\Gamma_t(t \rightarrow Wb) \propto \sigma_{1t}$ ,  $t$ -channel

$$1 - 2.3 \text{ fb}^{-1}: \quad \Gamma_t = 1.99^{+0.69}_{-0.55} \text{ GeV}$$

*Gross picture of top quark decay as expected*

# Rare Processes

# Photon Radiation in Top Pair Events CDF 6.0 $\text{fb}^{-1}$



## Semileptonic selection

- lepton,  $\cancel{E}_T$ , at least 3 jets one  $b$ -tag
- $H_T > 200 \text{ GeV}$
- + isolated energetic photon

## Background estimates

- photon fake rate from jets events  
(as function of  $\cancel{E}_T$ )
- rate of fake  $b$ -tags from  $b\bar{b}$  configurations


applied to events without photon/ $b$ -tag requirement, respectively



# Photon Radiation in Top Pair Events (2)

CDF 6.0 fb<sup>-1</sup>

A total of 30 candidate events observed




Run 193396 Event 1050006

$$\sigma_{t\bar{t}\gamma} = 0.18 \pm 0.08 \text{ pb} \quad \text{SM: } 0.17 \text{ pb (NLO from Madgraph+K-faktor)}$$

p-value: 0.0015 (3.0 $\sigma$ )

# Associated Higgs Production

- Check of the top quark's Yukawa coupling
- Enhancements possible
  - in 2HDM (e.g. MSSM) for low  $\tan \beta$
  - anomalous contributions to top-Yukawa coupling
  - in presence of a quark singlett  $T$ .




## Signature

In semileptonic top decay  $\ell, \not{E}_T, 4 b$ - and 2 light-jets


Signal to background discrimination

- DØ:  $H_T$  and  $n_{\text{jets}}$  most sensitive




$\ell, E_T$ , at least 4 jets

- sub-channels 4,  $\geq 5$  jets  
1, 2,  $\geq 3$   $b$ -tags



- Limits on  $\sigma_{ttH}\mathcal{B}(H \rightarrow bb)$
- Compared to BSM models
  - Here:  $G'$  with  $t'$



# Associated Higgs

$\ell, E_T$ , at least 4 jets

Signal to background discrimination

- Ensemble of neural networks
  - 1000 networks for each  $M_H$  value
  - each network with random set of 10 of 22 observables
  - including  $H_T$  and  $n_{\text{jets}}$


- Combination of network outputs  
Supra-Bayesian method

Major systematics:

- Process cross-section & JES

At  $M_H = 115 \text{ GeV}$  exclude  $> 22.9 \cdot \sigma_{\text{SM}}$

# Lepton+jets CDF, $7.5 \text{ fb}^{-1}$



# Associated Higgs

$\cancel{E}_T$ , at least 5 jets (but no charged lepton)

- $p_T > 50(40)$  GeV for (2nd) leading jet
- $H_T > 300$  GeV
- Additional QCD suppression with NNs


Signal to background discrimination

- NN with 13 input variables


At  $M_H = 115$  GeV exclude  $> 28.1 \cdot \sigma_{\text{SM}}$

# No lepton CDF, $5.7 \text{ fb}^{-1}$

All jets signal region (3-tag)



Limits for  $t\bar{t}H$  in missing  $E_T + \text{Jets}$  and All Jets



# Flavour Changing Neutral Currents

- SM expects  $B(t \rightarrow Zq) \sim 10^{-14}$
- BSM physics may yield measurable contribution

## Dilepton Analysis, CDF



$1.9 \text{ fb}^{-1}$

Selection:  $e^+e^-$  und  $\mu^+\mu^- + 4\text{jets}$ ,  $M_{\ell\ell} \approx M_Z$

Observable:  $\chi^2$  from mass constraints  
(Reconstructed  $Z$ ,  $W$  and top masses)

Fit of signal and SM simulation  
to  $\chi^2$ -distribution from data

$$B(t \rightarrow Zq) < 3.7\% \text{ (95\% C.L.)}$$



# Trilepton Analysis



DØ 4.1 fb<sup>-1</sup>

Selection:  $3\ell, \cancel{E}_T$  and possibly jets.

- 4 channels:  $eee, ee\mu, \mu\mu e, \mu\mu\mu$
- At least one  $\ell^+\ell^-$  pair with  $M_{\ell\ell} \approx M_Z$  (channel dependent distance cut)
- assignment of leptons to  $Z$  and  $W$   
 $\sim 100\%$  correct for  $ee\mu$  and  $\mu\mu e$   
 $\sim 90\%$  correct for  $eee$  and  $\mu\mu\mu$ .

Background estimates

- $WZ, ZZ$  and  $t\bar{t}$  from simulation
- $Z+jets$  and  $Z\gamma$  contribution from data



# Trilepton Analysis (2)

## Signal to background separation

- $H_T$
- $m_t^{\text{rec}} = M(\ell^+ \ell^- j)$   
use jet that yields  
 $m_t^{\text{rec}}$  closest to 172.5 GeV

## Result


$$B(t \rightarrow Zq) < 3.2\%$$

Expected  $B(t \rightarrow Zq) < 3.8\%$


## Couplings

$$v_{tqZ} < 0.19 \quad (q = u, c)$$

for only one non-vanishing coupling



# Summary of FCNC in Top Quark Decays



# BSM Particles



# Decay to Charged Higgs Boson

New particles in the decay alter deduced  $\sigma_{t\bar{t}}$   
depending on decay channel

$$\sigma_{t\bar{t}}^C = \sigma_{t\bar{t}} \cdot \frac{B^{\text{BSM}}(t\bar{t} \rightarrow C)}{B^{\text{SM}}(t\bar{t} \rightarrow C)}$$

$C = \ell + \text{jets}, \text{Dilepton, tauonic}$

- Check cross-section ratios
- Consider decay  $t \rightarrow bH^\pm$ 
  - with  $H^\pm \rightarrow cs$ , leptophobic  $H^\pm$
  - with  $H^\pm \rightarrow \tau\nu$ , tauonic  $H^\pm$
- Within MSSM relevant
  - leptophobic: at low  $\tan\beta$
  - tauonic: at high  $\tan\beta$



# Leptophobic Charged Higgs

DØ  $1 \text{ fb}^{-1}$


Sensitivity from semileptonic to dilepton cross-section

- $B(H^\pm \rightarrow cs) = 100\%$ 
  - semileptonic decay enhanced
  - dileptonic decay reduced
- fix  $\sigma_{t\bar{t}}$  at nominal value (7.48 pb)
- maximise likelihood wrt  $B(t \rightarrow bH^\pm)$

Dominating systematics:

- uncertainty of assumed  $\sigma_{t\bar{t}}$ , luminosity and  $b$ -tagging

Limits obtained w/ Feldman-Cousins:



# Leptophobic Charged Higgs

DØ  $1 \text{ fb}^{-1}$

Sensitivity from semileptonic to dilepton cross-section


- $B(H^\pm \rightarrow cs) = 100\%$ 
  - semileptonic decay enhanced
  - dileptonic decay reduced
- fix  $\sigma_{t\bar{t}}$  at nominal value (7.48 pb)
- maximise likelihood wrt  $B(t \rightarrow bH^\pm)$

Dominating systematics:

- uncertainty of assumed  $\sigma_{t\bar{t}}$ , luminosity and  $b$ -tagging

Limits obtained w/ Feldman-Cousins

... transferred to exclusions for specific models (e.g. CPX)



# Leptophobic Charged Higgs

CDF 2.2 fb<sup>-1</sup>

Bump search in dijet mass in  $\ell + \text{jets}$

- Reconstruct (fit) decay kinematics w/ constraints
  - top quark mass
  - leptonic  $W$ -mass
- Fifth jet in  $p_T$  joined with closest other jet if  $\Delta R < 1.0$



$\Rightarrow$  Obtain  $M(\text{dijet})$  with assignment of best  $\chi^2$

- Binned likelihood to extract  $B(t \rightarrow bH^\pm)$

Dominating uncertainties

- Simulation
- JES (for  $M_{H^\pm} \simeq M_W$ )

Limits also applicable to  $H^\pm \rightarrow ud$




Differences between dilepton and lepton+ $\tau$  yields sensitivity to total cross-section

- maximise likelihood simultaneously  
wrt  $B(t \rightarrow bH^\pm)$  and  $\sigma_{t\bar{t}}$
- reduces assumptions

Systematics:

- luminosity reduced
- multijet description dominates



Limits w/ Feldman-Cousins for a range of  $M_{H^\pm}$

In next-to-minimal SSM additional CP-odd Higgs bosons occur


$$t \rightarrow H^\pm b \rightarrow W^\pm a_1^0 b$$

- for  $m_{a_1^0} < 2m_b$  likely decay is  $a_1^0 \rightarrow \tau^+ \tau^-$
- decay products have low momenta  
⇒ undetected in other  $t\bar{t}$  analyses

## Analysis

- $t\bar{t}$  semileptonic selection
- require additional soft tracks (from taus)
- cross-checks for soft tracks in control regions

Limits on  $B(t \rightarrow bH^\pm)$  for various  $m_a$



# Colour Flow in Top Pair Decays

Verify colour singlet structure of hadronically decaying  $W$  in top decays

- distinguish from colour octet
- applicable to separate e.g.  $Z(H \rightarrow b\bar{b})$  from  $Z(g \rightarrow b\bar{b})$

## Observables


### Jet Pull:

Sum energy and distance weighted cell directions (in  $\eta$  and  $\varphi$ )

$$\vec{t} = \frac{1}{E_T^{\text{jet}}} \sum_{\text{Cal. cells}} E_{T,i} |r_i| \vec{r}_i$$

### Relative Jet Pull:

Angle between  $\vec{t}$  and connection to additional jet:  $\theta_{\text{rel},i}^{\text{pull}}$ .



# Color Flow Analysis

## Selection

- Consider  $\ell + \text{jets}$  sample with 2 (loose)  $b$ -tags
- 90% pure top pairs

Signal expectation:

- $W$ -jets point to each other
- $b$ -jets point away from each other


Control sample rather flat.

*Rel. jet pulls shows qualitatively expected behaviour*


Signal sample: no  $b$ -tags

$w$ -pairs

$$|m_{jj} - M_W| < 30 \text{ GeV}$$




$b$ -pairs




Control sample: no  $b$ -tags

$\theta_{\text{rel}}^{\text{pull}}$  of leading jet



second leading jet



# Color Flow: Result

D0 5.3 fb<sup>-1</sup>

Selection of  $\theta_{\text{rel}}^{\text{pull}}$ :

- $\Delta R > 2$ :  $\theta_{\text{rel}}^{\text{pull}}$  of leading  $p_T$  jet
- $\Delta R < 2$  both  $|\eta_d| < 1$ : use smaller  $\theta_{\text{rel}}^{\text{pull}}$ .
- $\Delta R < 2$  one  $|\eta_d| < 1$ :  $\theta_{\text{rel}}^{\text{pull}}$  of jet with  $|\eta_d| < 1$ .



## Dominating Systematics

- Signal shapes (1 vs. 8):  $\delta f_{\text{Singlet}} = \pm 0.118$
- Jet pull reconstruction:  $\delta f_{\text{Singlet}} = {}^{+0.100}_{-0.093}$

Simultaneously determine singlet vs octet fract.  
and top quark pair cross-section

$$f_{\text{Singlet}} = 0.56 \pm 0.36_{\text{stat}} \pm 0.22_{\text{syst}}$$

consistent with SM  $f_{\text{Singlet}} = 1$



# Summary and Conclusions

Searches for rare and BSM top quark decays

- Top Quark Width (35% uncertainty)
- Rare decays
  - Photon radiation in top pairs ( $3\sigma$  evidence)
  - $t\bar{t}H$  ( $\sigma_{\text{Limit}} \lesssim 23\sigma_{\text{SM}}$  at  $M_H = 115 \text{ GeV}$ )
  - FCNC in top decays ( $B < 3.2\%$ )
- Decays with BSM particles
  - Charged Higgs ( $B \lesssim 10...20\%$ )
- Colour structure

Top quark decay is verified to the few percent level

