



UNICA

UNIVERSITÀ
DEGLI STUDI
DI CAGLIARI



Asymmetric collisions in MadGraph5_aMC@NLO

Carlo Flore

Università di Cagliari & INFN - Sezione di Cagliari

Quarkonia as Tools 2025
Centre Paul Langevin, Aussois
Jan 9th, 2025

Work done in collaboration with
L. Manna, A. Safronov, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer

Outline

- Introduction
- NLO computations in MG5
- Implementation of asymmetric hadronic collisions at NLO in MG5
- Implementation of ep and eA collisions at NLO in MG5
- Conclusions and outlook

Introduction

- Asymmetric collisions offer the opportunity to study many aspects of the hadronic structure not accessible in symmetric pp reactions

Introduction

- Asymmetric collisions offer the opportunity to study many aspects of the hadronic structure not accessible in symmetric pp reactions
- they give access to the **nPDFs**, or allows for the study of **meson structure**

Introduction

- Asymmetric collisions offer the opportunity to study many aspects of the hadronic structure not accessible in symmetric pp reactions
- they give access to the **nPDFs**, or allows for the study of **meson structure**
- hadron structure analyses require **precise perturbative computations** for a large variety of processes

Introduction

- Asymmetric collisions offer the opportunity to study many aspects of the hadronic structure not accessible in symmetric pp reactions
- they give access to the **nPDFs**, or allows for the study of **meson structure**
- hadron structure analyses require **precise perturbative computations** for a large variety of processes
- **NLO pQCD codes** (e.g. MadGraph5_aMC@NLO, POWHEG, MCFM, HELAC-Onia, HERWIG, PYTHIA and SHERPA) have **limited capabilities** to perform calculations for asymmetric reactions

Introduction

- Asymmetric collisions offer the opportunity to study many aspects of the hadronic structure not accessible in symmetric pp reactions
- they give access to the **nPDFs**, or allows for the study of **meson structure**
- hadron structure analyses require **precise perturbative computations** for a large variety of processes
- **NLO pQCD codes** (e.g. MadGraph5_aMC@NLO, POWHEG, MCFM, HELAC-Onia, HERWIG, PYTHIA and SHERPA) have **limited capabilities** to perform calculations for asymmetric reactions
- **implementation** of asymmetric collisions is **often limited to private modified codes**, not publicly available

Introduction

- Asymmetric collisions offer the opportunity to study many aspects of the hadronic structure not accessible in symmetric pp reactions
- they give access to the **nPDFs**, or allows for the study of **meson structure**
- hadron structure analyses require **precise perturbative computations** for a large variety of processes
- **NLO pQCD codes** (e.g. MadGraph5_aMC@NLO, POWHEG, MCFM, HELAC-Onia, HERWIG, PYTHIA and SHERPA) have **limited capabilities** to perform calculations for asymmetric reactions
- **implementation** of asymmetric collisions is **often limited to private modified codes**, not publicly available
- we would like to **provide the community a tool performing reliable and precise automated perturbative computations at NLO in pQCD for any asymmetric reaction** (pA , AB , πA , ep , eA ...)

Introduction

- Asymmetric collisions offer the opportunity to study many aspects of the hadronic structure not accessible in symmetric pp reactions
- they give access to the **nPDFs**, or allows for the study of **meson structure**
- hadron structure analyses require **precise perturbative computations** for a large variety of processes
- **NLO pQCD codes** (e.g. MadGraph5_aMC@NLO, POWHEG, MCFM, HELAC-Onia, HERWIG, PYTHIA and SHERPA) have **limited capabilities** to perform calculations for asymmetric reactions
- **implementation** of asymmetric collisions is **often limited to private modified codes**, not publicly available
- we would like to **provide the community a tool performing reliable and precise automated perturbative computations at NLO in pQCD for any asymmetric reaction** (pA , AB , πA , ep , eA ...)

we have done it extending MG5

Symmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$d\sigma_{hh \rightarrow X} = \sum_{a,b} \int dx_a dx_b f_{a/h}(x_a, \mu_F; \text{LHAID_h}) f_{b/h}(x_b, \mu_F; \text{LHAID_h}) \\ \times d\hat{\sigma}_{ab \rightarrow X}(x_a, x_b, \mu_F, \alpha_S(\mu_R; \text{LHAID_h}))$$

Symmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$d\sigma_{hh \rightarrow X} = \sum_{a,b} \int dx_a dx_b f_{a/h}(x_a, \mu_F; \text{LHAID_h}) f_{b/h}(x_b, \mu_F; \text{LHAID_h}) \\ \times d\hat{\sigma}_{ab \rightarrow X}(x_a, x_b, \mu_F, \alpha_S(\mu_R; \text{LHAID_h}))$$

$$d\sigma^{\text{NLO}} = \sum_{\alpha} d\sigma^{(\text{NLO}, \alpha)}, \quad \alpha = \{E, S, C, SC\}$$

E: Event, fully resolved configuration

S, C, SC: soft, collinear, soft-collinear limits of **E (Counterevents)**

Symmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$d\sigma_{hh \rightarrow X} = \sum_{a,b} \int dx_a dx_b f_{a/h}(x_a, \mu_F; \text{LHAID_h}) f_{b/h}(x_b, \mu_F; \text{LHAID_h}) \\ \times d\hat{\sigma}_{ab \rightarrow X}(x_a, x_b, \mu_F, \alpha_S(\mu_R; \text{LHAID_h}))$$

$$d\sigma^{\text{NLO}} = \sum_{\alpha} d\sigma^{(\text{NLO}, \alpha)}, \quad \alpha = \{E, S, C, SC\}$$

E: Event, fully resolved configuration

S, C, SC: soft, collinear, soft-collinear limits of **E** (**Counterevents**)

$$d\sigma^{(\text{NLO}, \alpha)} = f_{a/h}(x_a^{(\alpha)}, \mu_F^{(\alpha)}; \text{LHAID_h}) f_{b/h}(x_b^{(\alpha)}, \mu_F^{(\alpha)}; \text{LHAID_h}) \\ \times W^{(\alpha)}(x_a^{(\alpha)}, x_b^{(\alpha)}, \mu_F^{(\alpha)}, \alpha_S(\mu_R^{(\alpha)}; \text{LHAID_h})) d\chi_{12} d\chi_{n+1}$$

Symmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$W^{(\alpha)} = g_S^{2b+2}(\mu_R^{(\alpha)}; \text{LHAID_h}) \left[\widehat{W}_0^{(\alpha)} + \widehat{W}_F^{(\alpha)} \log \left(\frac{\mu_F^{(\alpha)}}{Q} \right)^2 + \widehat{W}_R^{(\alpha)} \log \left(\frac{\mu_R^{(\alpha)}}{Q} \right)^2 \right] \\ + g_S^{2b}(\mu_R^{(\alpha)}; \text{LHAID_h}) \widehat{W}_B \delta_{\alpha S}$$

Symmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$W^{(\alpha)} = g_S^{2b+2}(\mu_R^{(\alpha)}; \text{LHAID_h}) \left[\widehat{W}_0^{(\alpha)} + \widehat{W}_F^{(\alpha)} \log \left(\frac{\mu_F^{(\alpha)}}{Q} \right)^2 + \widehat{W}_R^{(\alpha)} \log \left(\frac{\mu_R^{(\alpha)}}{Q} \right)^2 \right] \\ + g_S^{2b}(\mu_R^{(\alpha)}; \text{LHAID_h}) \widehat{W}_B \delta_{\alpha S}$$

- $\widehat{W}_0, \widehat{W}_R, \widehat{W}_F, \widehat{W}_B$ scale and PDF independent

Symmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$W^{(\alpha)} = g_S^{2b+2}(\mu_R^{(\alpha)}; \text{LHAID_h}) \left[\widehat{W}_0^{(\alpha)} + \widehat{W}_F^{(\alpha)} \log \left(\frac{\mu_F^{(\alpha)}}{Q} \right)^2 + \widehat{W}_R^{(\alpha)} \log \left(\frac{\mu_R^{(\alpha)}}{Q} \right)^2 \right] \\ + g_S^{2b}(\mu_R^{(\alpha)}; \text{LHAID_h}) \widehat{W}_B \delta_{\alpha S}$$

- $\widehat{W}_0, \widehat{W}_R, \widehat{W}_F, \widehat{W}_B$ scale and PDF independent
- \widehat{W}_B shares same kinematics of Soft Counterparts

Symmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$W^{(\alpha)} = g_S^{2b+2}(\mu_R^{(\alpha)}; \text{LHAID_h}) \left[\widehat{W}_0^{(\alpha)} + \widehat{W}_F^{(\alpha)} \log \left(\frac{\mu_F^{(\alpha)}}{Q} \right)^2 + \widehat{W}_R^{(\alpha)} \log \left(\frac{\mu_R^{(\alpha)}}{Q} \right)^2 \right] \\ + g_S^{2b}(\mu_R^{(\alpha)}; \text{LHAID_h}) \widehat{W}_B \delta_{\alpha S}$$

- $\widehat{W}_0, \widehat{W}_R, \widehat{W}_F, \widehat{W}_B$ scale and PDF independent
- \widehat{W}_B shares same kinematics of Soft Counterparts
- $\mu_F^{(\alpha)}$ may be different for Events and Counterparts

Symmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$W^{(\alpha)} = g_S^{2b+2}(\mu_R^{(\alpha)}; \text{LHAID_h}) \left[\widehat{W}_0^{(\alpha)} + \widehat{W}_F^{(\alpha)} \log \left(\frac{\mu_F^{(\alpha)}}{Q} \right)^2 + \widehat{W}_R^{(\alpha)} \log \left(\frac{\mu_R^{(\alpha)}}{Q} \right)^2 \right] + g_S^{2b}(\mu_R^{(\alpha)}; \text{LHAID_h}) \widehat{W}_B \delta_{\alpha S}$$

- $\widehat{W}_0, \widehat{W}_R, \widehat{W}_F, \widehat{W}_B$ scale and PDF independent
- \widehat{W}_B shares same kinematics of Soft Counterparts
- $\mu_F^{(\alpha)}$ may be different for Events and Counterparts
- output of a fixed-order NLO computation \Rightarrow **HwU files**

$$N_{\text{entr}} = \underbrace{4}_{\substack{2 \text{ bin edges} \\ 1 \text{ cross section} \\ 1 \text{ MC error}}} + \underbrace{(3 + N_\mu)}_{\substack{1 \text{ central value} \\ 2 \text{ min/max values} \\ N_\mu \text{ scale combinations}}} + \sum_{i=1}^{N_{\text{LHAID}}} \underbrace{(3 + N_{\text{mem}}^i)}_{\substack{1 \text{ central value} \\ 2 \text{ min/max values} \\ N_{\text{mem}} \text{ PDF members}}}$$

Asymmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$d\sigma_{h_1 h_2 \rightarrow X} = \sum_{a,b} \int dx_a dx_b f_{a/h_1}(x_a, \mu_F; \text{LHAID}_{h_1}) f_{b/h_2}(x_b, \mu_F; \text{LHAID}_{h_2}) \\ \times d\hat{\sigma}_{ab \rightarrow X}(x_a, x_b, \mu_F, \alpha_S(\mu_R; \text{LHAID}_{h_2}))$$

Asymmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$d\sigma_{h_1 h_2 \rightarrow X} = \sum_{a,b} \int dx_a dx_b f_{a/h_1}(x_a, \mu_F; \text{LHAID}_{h_1}) f_{b/h_2}(x_b, \mu_F; \text{LHAID}_{h_2}) \\ \times d\hat{\sigma}_{ab \rightarrow X}(x_a, x_b, \mu_F, \alpha_S(\mu_R; \text{LHAID}_{h_2}))$$

- strong coupling constant from the LHAPDF set of hadron h_2
- for pA collisions, full consistency if corresponding proton baseline for nPDF is used

Asymmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$d\sigma_{h_1 h_2 \rightarrow X} = \sum_{a,b} \int dx_a dx_b f_{a/h_1}(x_a, \mu_F; \text{LHAID}_{h_1}) f_{b/h_2}(x_b, \mu_F; \text{LHAID}_{h_2}) \\ \times d\hat{\sigma}_{ab \rightarrow X}(x_a, x_b, \mu_F, \alpha_S(\mu_R; \text{LHAID}_{h_2}))$$

- strong coupling constant from the LHAPDF set of hadron h_2
- for pA collisions, full consistency if corresponding proton baseline for nPDF is used
- output in the HwU files:
 1. **symmetric** cross sections **for each LHAID**

Asymmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$d\sigma_{h_1 h_2 \rightarrow X} = \sum_{a,b} \int dx_a dx_b f_{a/h_1}(x_a, \mu_F; \text{LHAID}_{h_1}) f_{b/h_2}(x_b, \mu_F; \text{LHAID}_{h_2}) \\ \times d\hat{\sigma}_{ab \rightarrow X}(x_a, x_b, \mu_F, \alpha_S(\mu_R; \text{LHAID}_{h_2}))$$

- strong coupling constant from the LHAPDF set of hadron h_2
- for pA collisions, full consistency if corresponding proton baseline for nPDF is used
- output in the HwU files:
 1. **symmetric** cross sections **for each LHAID**
 2. **asymmetric** cross section **for $h_1 h_2$ and $h_2 h_1$** ; h_1 PDF taken as the **first PDF set in the list** of LHAIDs, h_2 runs over the list of LHAIDs starting **from the second**

Asymmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$d\sigma_{h_1 h_2 \rightarrow X} = \sum_{a,b} \int dx_a dx_b f_{a/h_1}(x_a, \mu_F; \text{LHAID}_{h_1}) f_{b/h_2}(x_b, \mu_F; \text{LHAID}_{h_2}) \\ \times d\hat{\sigma}_{ab \rightarrow X}(x_a, x_b, \mu_F, \alpha_S(\mu_R; \text{LHAID}_{h_2}))$$

- strong coupling constant from the LHAPDF set of hadron h_2
- for pA collisions, full consistency if corresponding proton baseline for nPDF is used
- output in the HwU files:
 1. **symmetric** cross sections **for each LHAID**
 2. **asymmetric** cross section **for $h_1 h_2$ and $h_2 h_1$** ; h_1 PDF taken as the **first PDF set in the list** of LHAIDs, h_2 runs over the list of LHAIDs starting **from the second**
 3. **scale uncertainty** computed for the (central) PDF **for all the LHAIDs**

Asymmetric hadronic collisions in MG5 at NLO

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$d\sigma_{h_1 h_2 \rightarrow X} = \sum_{a,b} \int dx_a dx_b f_{a/h_1}(x_a, \mu_F; \text{LHAID}_{h_1}) f_{b/h_2}(x_b, \mu_F; \text{LHAID}_{h_2}) \\ \times d\hat{\sigma}_{ab \rightarrow X}(x_a, x_b, \mu_F, \alpha_S(\mu_R; \text{LHAID}_{h_2}))$$

- strong coupling constant from the LHAPDF set of hadron h_2
- for pA collisions, full consistency if corresponding proton baseline for nPDF is used
- output in the HwU files:
 1. **symmetric** cross sections **for each LHAID**
 2. **asymmetric** cross section **for $h_1 h_2$ and $h_2 h_1$** ; h_1 PDF taken as the **first PDF set in the list** of LHAIDs, h_2 runs over the list of LHAIDs starting **from the second**
 3. **scale uncertainty** computed for the (central) PDF **for all the LHAIDs**
 4. **Lab frame now default** one (by default, hidden flag `cms_frame = False`)

Asymmetric hadronic collisions in MG5 at NLO



CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation

$$d\sigma_{h_1 h_2 \rightarrow X} = \sum_{a,b} \int dx_a dx_b f_{a/h_1}(x_a, \mu_F; \text{LHAID}_{h_1}) f_{b/h_2}(x_b, \mu_F; \text{LHAID}_{h_2}) \\ \times d\hat{\sigma}_{ab \rightarrow X}(x_a, x_b, \mu_F, \alpha_S(\mu_R; \text{LHAID}_{h_2}))$$

- strong coupling constant from the LHAPDF set of hadron h_2
- for pA collisions, full consistency if corresponding proton baseline for nPDF is used
- output in the HwU files:
 1. **symmetric** cross sections **for each LHAID**
 2. **asymmetric** cross section **for $h_1 h_2$ and $h_2 h_1$** ; h_1 PDF taken as the **first PDF set in the list** of LHAIDs, h_2 **runs over the list** of LHAIDs starting **from the second**
 3. **scale uncertainty** computed for the (central) PDF **for all the LHAIDs**
 4. **Lab frame now default** one (by default, hidden flag `cms_frame = False`)
 5. in general $N_{\text{entr}}^{\text{asym}} > N_{\text{entr}}^{\text{symm}}$, may also be $N_{\text{entr}}^{\text{asym}} \gg N_{\text{entr}}^{\text{symm}}$

Validation - pA


CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation



- only nPDF uncertainty shown
- excellent agreement with MCFM

Validation - πA



CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation



- MG5 and JAM MC statistical uncertainties tend to grow in the most forward region
- MG5 and JAM results essentially identical and compatible with experimental data



Selection of results (I)

CF, D. Kikola, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation





Selection of results (I)

CF, D. Kikola, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation



Selection of results (I)


CF, D. Kikola, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation



- nPDF uncertainty dominant at forward (pseudo)rapidity
- scale and NPDF uncertainties comparable in the backward region
- such predictions can be now obtained in a single run by indicating three LHAIIDs in the run_card

Selection of results (II)

CF, D. Kikola, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Sfronov, in preparation



- nuclear modification factor: $R_{p\text{Pb}} = \frac{1}{A_{\text{Pb}}} \frac{d\sigma_{p\text{Pb}}}{d\sigma_{pp}}$
- approximation for uncertainty computation:

$$\delta(R_{AB}) = \frac{1}{A_A A_B} \frac{\delta(d\sigma_{AB})}{d\sigma_{pp}^{\text{central}}}$$

- $R_{p\text{Pb}}$ automatically computed in a single run

ep and *eA* collisions

L. Manna et al., in preparation; L. Manna, A. Safronov et al., PoS
EPS-HEP2023 (2024) 274; L. Manna, A. Safronov et al. PoS DIS2024 (2024)

We now focus on photoproduction:

ep and eA collisions

L. Manna et al., in preparation; L. Manna, A. Safronov et al., PoS
EPS-HEP2023 (2024) 274; L. Manna, A. Safronov et al. PoS DIS2024 (2024)

We now focus on photoproduction:

- direct contribution:

$$\sigma_{eh(A) \rightarrow C+X} = \sum_b \int dx_\gamma dx_b f_\gamma^e(x_\gamma, Q_{\max}^2) \\ \times f_b^{h(A)}(x_b, \mu_F; \text{LHAID_h}) \hat{\sigma}_{\gamma b \rightarrow C+X}(x_\gamma, x_b, \mu_F, \mu_R)$$

ep and eA collisions

L. Manna et al., in preparation; L. Manna, A. Safronov et al., PoS
EPS-HEP2023 (2024) 274; L. Manna, A. Safronov et al. PoS DIS2024 (2024)

We now focus on photoproduction:

- direct contribution:

$$\sigma_{eh(A)\rightarrow C+X} = \sum_b \int dx_\gamma dx_b f_\gamma^e(x_\gamma, Q_{\max}^2) \\ \times f_b^{h(A)}(x_b, \mu_F; \text{LHAID_h}) \hat{\sigma}_{\gamma b \rightarrow C+X}(x_\gamma, x_b, \mu_F, \mu_R)$$

- resolved contribution:

$$\sigma_{eh(A)\rightarrow C+X} = \sum_{a,b} \int dx_\gamma dx_a dx_b f_\gamma^e(x_\gamma, Q_{\max}^2) \\ \times f_a^\gamma(x_a, \mu_F; \text{LHAID_}\gamma) f_b^{h(A)}(x_b, \mu_F; \text{LHAID_h}) \hat{\sigma}_{ab \rightarrow C+X}$$

ep and eA collisions

L. Manna et al, in preparation; L. Manna, A. Safronov et al., PoS
EPS-HEP2023 (2024) 274; L. Manna, A. Safronov et al. PoS DIS2024 (2024)

We now focus on photoproduction:

- direct contribution:

$$\sigma_{eh(A)\rightarrow C+X} = \sum_b \int dx_\gamma dx_b f_\gamma^e(x_\gamma, Q_{\max}^2) \\ \times f_b^{h(A)}(x_b, \mu_F; \text{LHAID_h}) \hat{\sigma}_{\gamma b \rightarrow C+X}(x_\gamma, x_b, \mu_F, \mu_R)$$

- resolved contribution:

$$\sigma_{eh(A)\rightarrow C+X} = \sum_{a,b} \int dx_\gamma dx_a dx_b f_\gamma^e(x_\gamma, Q_{\max}^2) \\ \times f_a^\gamma(x_a, \mu_F; \text{LHAID_}\gamma) f_b^{h(A)}(x_b, \mu_F; \text{LHAID_h}) \hat{\sigma}_{ab \rightarrow C+X}$$

- fixed some bugs related to the WW distribution and modified the boost routines

ep and eA collisions

L. Manna et al, in preparation; L. Manna, A. Safronov et al., PoS
EPS-HEP2023 (2024) 274; L. Manna, A. Safronov et al. PoS DIS2024 (2024)

We now focus on photoproduction:

- direct contribution:

$$\sigma_{eh(A)\rightarrow C+X} = \sum_b \int dx_\gamma dx_b f_\gamma^e(x_\gamma, Q_{\max}^2) \\ \times f_b^{h(A)}(x_b, \mu_F; \text{LHAID_h}) \hat{\sigma}_{\gamma b \rightarrow C+X}(x_\gamma, x_b, \mu_F, \mu_R)$$



- resolved contribution:

$$\sigma_{eh(A)\rightarrow C+X} = \sum_{a,b} \int dx_\gamma dx_a dx_b f_\gamma^e(x_\gamma, Q_{\max}^2) \\ \times f_a^\gamma(x_a, \mu_F; \text{LHAID_}\gamma) f_b^{h(A)}(x_b, \mu_F; \text{LHAID_h}) \hat{\sigma}_{ab \rightarrow C+X}$$

- fixed some bugs related to the WW distribution and modified the boost routines
- lab frame now default

Inclusive photoproduction - validation at LO


L. Manna et al., in preparation; L. Manna, A. Safronov et al., PoS EPS-HEP2023 (2024) 274; L. Manna, A. Safronov et al. PoS DIS2024 (2024)



- validation for $c\bar{c}$ (left) and $b\bar{b}$ photoproduction (HERA kinematics) at LO
- agreement $\mathcal{O}(1\%)$ with HELAC-Onia and $\sim 4\%$ with FMNR

Inclusive photoproduction - validation at NLO



L. Manna et al., in preparation; L. Manna, A. Safronov et al., PoS EPS-HEP2023 (2024) 274; L. Manna, A. Safronov et al. PoS DIS2024 (2024)



- validation for $c\bar{c}$ (left) and $b\bar{b}$ photoproduction (HERA kinematics) at NLO
- very satisfactory agreement ($\mathcal{O}(1\%)$) with FMNR

Inclusive $b\bar{b}$ photoproduction - predictions


L. Manna et al., in preparation



- NLO predictions for future ep colliders
- limited P_T range at EicC
- P_T spectrum extend to $P_T \sim 60$ GeV at LHeC and FCC-eH

Resolved contribution - validation at NLO



L. Manna et al. in preparation; L. Manna, A. Safronov et al., PoS
EPS-HEP2023 (2024) 274; L. Manna, A. Safronov et al. PoS DIS2024 (2024)



- ratio MG5 to FMNR with combined Monte Carlo uncertainties
- $\sim \mathcal{O}(2\%)$ agreement with FMNR

$b\bar{b}$ photoproduction at the EIC





L. Manna et al. in preparation; L. Manna, A. Safronov et al., PoS EPS-HEP2023 (2024) 274; L. Manna, A. Safronov et al. PoS DIS2024 (2024)



- predictions for the EIC at $\sqrt{s_{ep}} = 45$ GeV (left) and $\sqrt{s_{ep}} = 140$ GeV (right)
- resolved contribution comparable with direct one around $P_T \sim 9 - 10$ GeV at $\sqrt{s_{ep}} = 45$ GeV
- direct contribution dominant but resolved one not negligible at $\sqrt{s_{ep}} = 140$ GeV





R_{eAu} @ EIC - $\sqrt{s_{ep}} = 45$ GeV

EIC Onium review, arXiv:2409.03691 [hep-ph]



R_{eAu} @ EIC - $\sqrt{s_{ep}} = 90$ GeV

EIC Onium review, arXiv:2409.03691 [hep-ph]



Conclusions and outlook

- **MG5** now available online in its **full NLO version on NLOAccess**
- validated and developed MG5 for **asymmetric collisions**
 - **photoproduction** in ℓh collisions
 - **nuclear modification factors** in pA , πA and AB collisions

Conclusions and outlook

- **MG5** now available online in its **full NLO version on NLOAccess**
- validated and developed MG5 for **asymmetric collisions**
 - **photoproduction** in ℓh collisions
 - **nuclear modification factors** in pA , πA and AB collisions
- next:
 - MG5 extension to (N)LO+PS for asymmetric collisions
 - DIS implementation at (N)LO+PS
 - automation of onium production computations at (N)LO in MG5

Alice's talk

- automation of spin and transverse momentum effects for quarkonium production

Conclusions and outlook

- **MG5** now available online in its **full NLO version on NLOAccess**
- validated and developed MG5 for **asymmetric collisions**
 - **photoproduction** in ℓh collisions
 - **nuclear modification factors** in pA , πA and AB collisions
- next:
 - MG5 extension to (N)LO+PS for asymmetric collisions
 - DIS implementation at (N)LO+PS
 - automation of onium production computations at (N)LO in MG5
- automation of spin and transverse momentum effects for quarkonium production

Alice's talk

Thank you



Backup

MG5 HwU output for asymmetric collisions

reweight_scale	reweight_PDF	No. of entries (symmetric): $N_{\text{entr}}^{\text{sym}}$	No. of entries (asymmetric): $N_{\text{entr}}^{\text{asym}}$
True	True	$4 + N_{\text{LHAID}} (3 + N_{\mu}) + \sum_{i=1}^{N_{\text{LHAID}}} (3 + N_{\text{mem}}^i)$	$N_{\text{entr}}^{\text{sym}} + 2(N_{\text{LHAID}} - 1) (3 + N_{\mu}) + 2 \sum_{i=2}^{N_{\text{LHAID}}} (3 + N_{\text{mem}}^i)$
True	False	$4 + N_{\text{LHAID}} (3 + N_{\mu}) + N_{\text{LHAID}} (3 + 1)$	$N_{\text{entr}}^{\text{sym}} + 2(N_{\text{LHAID}} - 1) (3 + N_{\mu}) + 2(N_{\text{LHAID}} - 1) (3 + 1)$
False	True	$4 + \sum_{i=1}^{N_{\text{LHAID}}} (3 + N_{\text{mem}}^i)$	$N_{\text{entr}}^{\text{sym}} + 2 \sum_{i=2}^{N_{\text{LHAID}}} (3 + N_{\text{mem}}^i)$
False	False	$4 + N_{\text{LHAID}} (3 + 1)$	$N_{\text{entr}}^{\text{sym}} + 2(N_{\text{LHAID}} - 1) (3 + 1)$

Selection of results



CF, D. Kikota, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation



- nPDF uncertainty dominant at forward (pseudo)rapidity
- scale and NPDF uncertainties comparable in the backward region
- such predictions can be now obtained in a single run by indicating three LHAIIDs in the run_card

Selection of results





CF, D. Kikota, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation



- nPDF uncertainty dominant at forward (pseudo)rapidity
- scale and NPDF uncertainties comparable in the backward region
- such predictions can be now obtained in a single run by indicating three LHAIDs in the run_card



Selection of results

CF, D. Kikoła, A. Kusina, J.-P. Lansberg, O. Mattelaer, A. Safronov, in preparation



Selection of results

L. Manna et al., in preparation



- NLO predictions for future ep colliders
- limited P_T range at EICc
- P_T spectrum extend to $P_T \sim 60 \text{ GeV}$ at LHeC and FCC-eh