





**FPF**  
*orward hysics acility*


The logo for the Forward Physics Facility (FPF) features a red double-headed arrow passing through a central point. From this point, two blue fan-shaped regions extend outwards, representing particle detectors. The text 'FPF' is written in large, bold, black letters, with 'orward hysics acility' written in a smaller, italicized font below it.



**FASER** designed to search for new, light, long-lived particles (**LLPs**), and study **neutrinos**.


These are produced in the **decay of light hadrons** which are produced in the LHC collisions.

Light hadron production is very peaked in the **forward direction**, extremely collimated with the beam collision axis line of sight (LOS), hence even small detectors covering the angular region less than a milliradian around the LOS have good physics sensitivity.



# Physics Motivation


- The **Forward Physics Facility** has a rich and **broad** physics programme
- Three main physics motivations
  - Beyond Standard Model (BSM) “dark sector” searches
  - Neutrino physics
  - QCD physics



- The **Forward Physics Facility** has a rich and **broad** physics programme
- Three main physics motivations
  - Beyond Standard Model (BSM) “dark sector” searches
  - Neutrino physics
  - QCD physics
- In order to fully benefit from the increase in luminosity from the HL-LHC, the FPF will allow:
  - **Longer detectors** to increase target/decay volume
  - **Wider detectors** to increase sensitivity to heavy flavour produced particles
  - Space for **new detectors** with complementary physics capabilities

# FORWARD PHYSICS FACILITY

- A comprehensive site selection study by the CERN Civil Engineering group has identified an ideal location ~600 m west of ATLAS.




- The site is on CERN land in France
- The cavern is 65 m-long, 9 m-wide/high
- Shielded from ATLAS by 200m of rock
- Disconnected from LHC tunnel
- Cost: ~CHF 40M
- Vibration, safety studies: can construct FPF without disrupting LHC operations
- Radiation studies: can work in FPF while LHC is running (HL-LHC starts 2029)

# FASER2 detector overview

FASER2 scaled up version of **FASER** experiment.

Designed to search for decaying **BSM LLPs** (**100x larger** transverse size to FASER, and 7x longer decay volume) and to measure charge/momentum of muons from **neutrino interactions** in FLArE and FASERv2)



Larger transverse size => need to use different technologies than in FASER:

Investigated different options for tracking detector technology and magnets.

Baseline tracker – **SciFi tracker** (like LHCb modules)

Two options for magnet:


- Custom made **super-conducting dipole** (2Tm bending power 1.7x1.7m<sup>2</sup> square air-core aperture):
  - Preliminary quote from Toshiba: 4.1MCHF 3-4y lead-time
- 4 off-the-shelf **crystal-puller magnet** units (2Tm (central) bending power 1.6m diameter air-core aperture):
  - Preliminary quote from Toshiba: 2.3MCHF 1-2y lead-time



# Summary



- The FPF greatly **expands the LHC physics programme**
- The **beam** is there already
- Broad and **interdisciplinary** physics programme
  - Neutrinos, QCD, Cosmic Ray, BSM, ...
- The cost is **modest**
- Fully aligned with existing **European strategy**
- Great **test-bed** for new ideas
- Could run in **LHC Run 4**





# Oxford position

- We have been involved in physics studies, design studies, mechanical engineering, magnet procurement in UK industry...
- The FPF has very many desirable features (capital cost, operational cost, environmental, timescale, training...)
- Medium-cost, medium term projects are needed
- There are scenarios in which it would proceed
- We are well positioned to exploit any such opportunity

# EXTRAS

**CERNCOURIER** | Reporting on international high-energy physics

Physics ▾ Technology ▾ Community ▾ In focus Magazine

SEARCHES FOR NEW PHYSICS | MEETING REPORT

## Looking forward at the LHC

1 September 2023




**New territory** 3D rendering of the proposed Forward Physics Facility and its experiments.  
Credit: A Navascues Corago

The Forward Physics Facility (FPF) is a proposed new facility to operate concurrently with the High-Luminosity LHC, housing several new experiments on the ATLAS collision axis. The FPF offers a broad, far-reaching physics programme ranging from neutrino, QCD and hadron-structure studies to beyond-the-Standard Model (BSM) searches. The project, which is being studied within the Physics Beyond Colliders initiative, would exploit the pre-existing HL-LHC beams and thus have minimal energy-consumption requirements.

On 8 and 9 June, the 6th workshop on the Forward Physics Facility was held at CERN and online. Attracting about 160 participants, the workshop was organised in sessions focusing on the facility design, the proposed experiments and physics studies, leaving plenty of time for discussion about the next steps.

- The FPF experiments have **strong sensitivity** in all of the dark sector PBC benchmark models
- In some cases the sensitivity is not as strong as that of SHiP
- There are classes of models where the FPF is **more powerful than SHiP**

$$\begin{array}{l}
 m_1 \text{ --- } \curvearrowright \\
 m_0 \text{ --- } \curvearrowleft \\
 \Delta \equiv \frac{\Delta m}{m_0} \equiv \frac{m_1 - m_0}{m_0} \\
 \mathcal{O}_m = \frac{1}{\Lambda_m} \bar{\chi}_1 \sigma^{\mu\nu} \chi_0 F_{\mu\nu}
 \end{array}$$




- LLPs can result from weak couplings
- They can also arise generically from compressed spectra
- In this case soft photons are difficult to detect
- At the LHC the parents have Lorentz boosts of order 1000.
- FASER2 can detect GeV particles with even  $\sim$ MeV mass splittings
- Difficult at SHiP

Further examples in the backup slides

# Neutrinos at the FPF

LHC provides a **strongly collimated** beam of **TeV energy neutrinos** of **all three flavours** in the far forward direction.





Proposed FPF experiment have potential to detect O(1M) neutrinos:  
 O(**10<sup>5</sup>**) electron neutrinos, O(**10<sup>6</sup>**) muon neutrinos, O(**10<sup>4</sup>**) tau neutrinos

# Application: Astroparticle Physics

Forward **charm** production at the LHC



Constraints on **prompt atmospheric neutrino flux** at IceCube




cosmic ray muon puzzle:  
observed excess of muons compared to hadronic interaction models

forward pion/kaons fluxes will provide crucial input

See also:  
Talk Subir Sarkar


# Application: QCD

## TeV Energy Neutrino Interaction




# Neutrino DIS at the FPF


Collider Neutrino Experiments are a **Neutrino-Ion Collider** at **EIC** center of mass energies



neutrino DIS data will **improve PDFs**  
 [FPF, P5 Input] [Cruz-Martinez et al. 2309.09581]






reduced PDF uncertainties for many LHC processes



**breaks PDF/BSM degeneracy** for main LHC experiments

[Rojo, FPF7 Talk]


# FPF CE Site Investigation Works



- 20cm diameter core taken to 100m depth at proposed FPF location
- Detailed geological study of core carried out
  - No showstoppers identified
  - Area looks **good for excavation**



- There are **4 current experiments** being designed for the FPF
  - Diverse technologies optimized for particular SM and BSM topics
  - FPF covers  $\eta > 5.5$ , experiments on LOS cover  $\eta \gtrsim 7$
- Experimental layout being optimised
  - Many **opportunities** for new groups



- At present there are **4 experiments** being designed for the FPF
  - Diverse technologies optimized for particular SM and BSM topics
  - FPF covers  $\eta > 5.5$ , experiments on LOS cover  $\eta \gtrsim 7$
- Experiment still in early design phase
  - Many **opportunities** for new groups
- All but one of the proposed FPF experiments have pathfinder versions **already running** at the LHC
  - **FASER, FASERv, milliQan**
  - Physics results from these experiments give confidence that the FPF physics potential can be realized

[Phys. Rev. D \*\*102\*\*, 032002](#)  
[Phys. Rev. Lett. \*\*131\*\*, 031801](#)  
[Phys. Lett B, \*\*848\*\*, 138378](#)  
[arxiv: 2403.12520](#)  
[CERN-FASER-CONF-2024-001](#)

PHYSICAL REVIEW LETTERS  
 Highlights Recent Accepted Collections Authors Referees Search Press About

Featured in Physics Editors' Suggestion Open Access

First Direct Observation of Collider Neutrinos with FASER at the LHC  
 et al. (FASER Collaboration)  
 Phys. Rev. Lett. **131**, 031801 – Published 19 July 2023

Viewpoint: The Dawn of Collider Neutrino Physics

References No Citing Articles PDF HTML Export Citation

ABSTRACT  
 We report the first direct observation of neutrino interactions at a particle collider experiment. Neutrino candidate events are identified in a 13.6 TeV center-of-mass energy pp collision dataset of 35.4 fb<sup>-1</sup> using the active electronic components of the FASER detector at the Large Hadron Collider. The candidates are required to have a track propagating through the entire length of the FASER detector and be consistent with a muon neutrino charged-current interaction. We infer 153<sup>+12</sup> neutrino interactions with a significance of 16 standard deviations above the background-only hypothesis. These events are consistent with the characteristics expected from neutrino interactions in terms of secondary particle production and spatial distribution, and they imply the observation of both neutrinos and anti-neutrinos with an incident neutrino energy of significantly above 200 GeV.

Physics Letters B  
 Volume 848, January 2024, 138378

Letter

Search for dark photons with the FASER detector at the LHC  
 FASER Collaboration\*

Show more

+ Add to Mendeley Share 55 Cite

https://doi.org/10.1016/j.physletb.2023.138378

Under a Creative Commons license

Abstract  
 The FASER experiment at the LHC is designed to search for light, weakly-interacting particles produced in proton-proton collisions at the ATLAS interaction point that travel in the far-forward direction. The first results from a search for dark photons decaying to an electron-positron pair, using a dataset corresponding to an integrated luminosity of 27.0 fb<sup>-1</sup> collected at centre-of-mass energy  $\sqrt{s} = 13.6$  TeV in 2022 in LHC Run 3, are presented. No events are seen in an almost background-free analysis, yielding world-leading constraints on dark photons with couplings  $\epsilon \sim 2 \times 10^{-5} - 1 \times 10^{-4}$  and masses  $\sim 17$  MeV – 70 MeV. The analysis is also used to probe the parameter space of a massive gauge boson from a U(1)<sub>B-L</sub> model, with couplings  $g_{B-L} \sim 5 \times 10^{-6} - 2 \times 10^{-5}$  and masses  $\sim 15$  MeV – 40 MeV excluded for the first time.

PHYSICAL REVIEW D  
 covering particles, fields, gravitation, and cosmology

Highlights Recent Accepted Collections Authors Referees Search Press About

Open Access

Search for millicharged particles in proton-proton collisions at  $\sqrt{s} = 13$  TeV  
 A. Ball et al.  
 Phys. Rev. D **102**, 032002 – Published 6 August 2020

Article References Citing Articles (23) PDF HTML Export Citation

ABSTRACT  
 We report on a search for elementary particles with charges much smaller than the electron charge using a data sample of proton-proton collisions provided by the CERN Large Hadron Collider in 2018, corresponding to an integrated luminosity of 37.5 fb<sup>-1</sup> at a center-of-mass energy of 13 TeV. A prototype scintillator-based detector is deployed to conduct the first search at a hadron collider sensitive to particles with charges  $\leq 0.1e$ . The existence of new particles with masses between 20 and 4700 MeV is excluded at 95% confidence level for charges between 0.006e and 0.3e, depending on their mass. New sensitivity is achieved for masses larger than 700 MeV.

PHYSICAL REVIEW D  
 covering particles, fields, gravitation, and cosmology

Highlights Recent Accepted Collections Authors Referees Search Press About

Open Access

Search for millicharged particles in proton-proton collisions at  $\sqrt{s} = 13$  TeV  
 A. Ball et al.  
 Phys. Rev. D **102**, 032002 – Published 6 August 2020

Article References Citing Articles (23) PDF HTML Export Citation

ABSTRACT  
 We report on a search for elementary particles with charges much smaller than the electron charge using a data sample of proton-proton collisions provided by the CERN Large Hadron Collider in 2018, corresponding to an integrated luminosity of 37.5 fb<sup>-1</sup> at a center-of-mass energy of 13 TeV. A prototype scintillator-based detector is deployed to conduct the first search at a hadron collider sensitive to particles with charges  $\leq 0.1e$ . The existence of new particles with masses between 20 and 4700 MeV is excluded at 95% confidence level for charges between 0.006e and 0.3e, depending on their mass. New sensitivity is achieved for masses larger than 700 MeV.

CERN-FASER-CONF-2024-001  
 April 6, 2024

Search for Axion-Like Particles in Photonic Final States with the FASER Detector at the LHC  
 FASER Collaboration

The first FASER search for a light, long-lived particle decaying into a pair of photons is reported. The search uses the collected 2022 and 2023 LHC proton-proton collision data at  $\sqrt{s} = 13.6$  TeV, corresponding to an integrated luminosity of 57.7 fb<sup>-1</sup>. A model with axion-like particles (ALPs) dominantly coupling to weak gauge bosons is targeted, probing a mass range between 20 and 500 MeV and couplings to the Standard Model particles,  $g_{\gamma\gamma}$ , between  $10^{-10}$  and  $10^{-9}$  GeV<sup>-1</sup>. Signal events are characterized by high energy deposits in the electromagnetic calorimeter and no signal in the muon scintillators. One event is observed, compared to a background expectation of  $0.62 \pm 0.88$  events, which is entirely dominated by muonino interactions. World-leading constraints on ALPs are obtained for masses up to 300 MeV and couplings around  $10^{-10}$  GeV<sup>-1</sup>, testing a previously unexplored region of parameter space.

CONTENTS

- I. Introduction 2
- II. Axion-Like Particles in FASER 2
- III. The FASER Detector 4
- IV. Data and Simulation Samples 6
- V. Event Selection and Reconstruction 7
- VI. Background Estimation 8
- A. Neutrino Background 9
- B. Other Background Contributions 12
- VII. Systematic Uncertainties 14
- VIII. Results 15
- IX. Conclusions 17

Appendix  
 1. Scintillator Efficiencies in 2022 and 2023 Data 18  
 2. FASER ALP Limit in Relation to Previous Experiments 18  
 3. Event Display 18  
 4. Neutrino Compositions by Flavour 18

References 22

# FASERv2 detector overview

**Emulsion** tungsten neutrino detector, scaled up version of **FASERv**.

20tonne target mass:

- 80x20cm<sup>2</sup> in transverse plane (64x25cm<sup>2</sup> also under consideration)
- 8m long
- 3300x 2mm thick tungsten plates interleaved with emulsion film

Electronic detectors to allow to **timestamp** events and connect muons from neutrino interactions with those reconstructed in the FASER2 spectrometer (for charge/momentum measurement).

Emulsion needs to be **replaced** with track occupancy of  $O(10^5)$  tracks/cm<sup>2</sup> (about 2months of HL-LHC running)




To save cost would like to improve this to have one set of emulsion per year:

- reduce track multiplicity by factor of 3 with **sweeper magnet**,
- or **improve tools** to allow analysis with higher occupancy

Under study...

Several testbeam studies and R&D ongoing related to:

- Detector assembly
- Emulsion film optimization (long term operation, higher track occupancy)
- Analysis tools (re-reconstruct FASERv data skipping every-other-film to mimic 2mm thick tungsten)




# FORMOSA detector overview


FORMOSA is a **scintillator** based **millicharged particle** detector (similar to the running **miliQan** experiment).

Idea is to see low scintillation signal in multiple bars pointing at IP.

Baseline design: 20x20 array of scintillator bars (with surrounding scintillators to veto backgrounds)



At start of 2024 a small **demonstrator detector** was installed in TI12 (FASER location) to study the performance with a large rate of through going muons from the IP. Using this demonstrator the DAQ concept for FORMOSA has been validated.



Team studying updated design with central submodule of CeBr3 scintillator **~35 higher light yield**, allows significant increase in sensitivity for low charges.

# FLArE detector overview

## Liquid Argon TPC

Builds on investment in liquid noble gas detectors

Fiducial (active) mass **10 (30) tons**

Single-wall 8.8 x 2.0 x 2.4 m foam-insulated cryostat



7x3 modules, for space charge, scintillator light (trigger)

Anode charge readout pixelated ~5mm size

Alternative readout 3D optical TPC (ARIADNE)

TimePix detects wavelength-shifted secondary scintillation light

THick Gaseous Electron Multiplier in gas phase of a dual-phase LArTPC



Horizontal “filing cabinet” installation

Cryogenic plant downstream of all detectors  
(noise/vibration)

Hadron calorimeter/muon spectrometer downstream  
for low mmtm

Magnetized iron plates interleaved with scintillator  
modules

FASER2 provides the spectrometer for high mmtm

# FPF Documentation

**FPF workshop series:**  
[FPF1](#), [FPF2](#), [FPF3](#), [FPF4](#), [FPF5](#),  
[FPF6](#), [FPF7](#), [FPF Theory Day](#)

**FPF Paper:**  
[2109.10905](#)


~75 pages, ~80 authors

**Snowmass Whitepaper:**  
[2203.05090](#)

~450 pages, ~250 authors

**Recent Summary:**  
[FPF Update](#)

**Technical Documents:**  
[Facility Technical Study](#)  
[Muon Flux Study](#)  
[Vibration Study](#)  
[Geotechnical Report](#)



- FPF is a proposed **new facility** to house several experiments on the collision axis line-of-sight in the HL-LHC era
  - **Maximise the physics from the HL-LHC**
- The FPF experiments would :
  - Have **world leading sensitivity** in several **BSM models**
  - Study O(1M) **TeV neutrinos** (covering all flavours) with important implications for:
    - **Neutrino physics**
    - **QCD**
    - **Astroparticle physics**
- Following many studies within the PBC we have come-up with a baseline design for the facility that can house the proposed experiments
  - No showstoppers identified for implementing the facility or for operating the experiments
  - Facility cost of **only ~40MCHF**
- Conceptual designs of the experiments are ongoing with **baseline solutions available** for the different detector components
  - More studies will allow better optimizations and improved physics reach
- Plan to submit a proposal for the FPF to the ESPP in Q1 2025
  - If viewed favourably the facility can be implemented to allow **first physics during Run 4**

# Documentation on facility studies

CERN-PBC-NOTE 2023-002

<https://cds.cern.ch/record/2851822/>

CERN-PBC-Notes-2024-004

<https://cds.cern.ch/record/2904086>

CERN-PBC-NOTE 2024-003

<https://cds.cern.ch/record/2901520/>



**CERN-PBC-NOTE 2023-002**  
7 March 2023  
Jamie.Boyd@cern.ch

### Update on the FPF Facility technical studies

**FPF PBC Working Group:**  
M. Andreini, G. Arduini, K. Balazs, J. Boyd, R. Bozzi, F. Cerutti, F. Corsanego, J-P. Corso, L. Elie, A. Infantino, A. Navasques Cornago, J. Osborne, G. Peon, M. Sabaté Gilarte  
CERN, CH-1211 Geneva, Switzerland

Keywords: FPF


---

**Summary**

The Forward Physics Facility (FPF) is a proposed new facility to house several new experiments at the CERN High Luminosity LHC (HL-LHC). The FPF is located such that the experiments can be aligned with the collision axis line of sight (LOS), a location which allows many interesting physics measurements and searches for new physics to be carried out. The status of technical studies related to the FPF, as well as the physics potential were documented in Ref. [1] which was released in March 2022. This note documents updates to the FPF technical studies completed since that time.

---

1



**CERN-PBC-NOTE 2024-XXX**  
18 June 2024  
Jamie.Boyd@cern.ch

### Update of Facility Technical Studies for the FPF

**FPF PBC Working Group**  
K. Balazs, J. Boyd, T. Bud, J-P. Corso, D. Gamba, A. Magazini, A. Navasques Cornago, J. Osborne (CERN, CH-1211 Geneva, Switzerland)

**Contributors from the FPF Experiments**  
FLArE: L. Bartoszek (Bartoszek Engineering), Y. Li (BNL), S. Linden (BNL), C. Miraval (BNL), S. Trabocchi (BNL)  
FASER2: S. Bosco (Bern)  
FASER2: N. Sumi (KEK), J. Carroll (Liverpool), A. Lowe (Oxford)  
FORMOSA: R. Loos (CERN)

Keywords: FPF

---

**Executive Summary**


The Forward Physics Facility (FPF) has been proposed to house a set of experiments to detect collider neutrinos and search for new particles in the High-Luminosity LHC era. This report provides an update to the space and infrastructure requirements of the Facility, a result of integration studies carried out by CERN technical teams in conjunction with the FPF experimental community.

Previous radio protection studies showed that access to the FPF cavern during LHC beam operation was expected to be possible. This update includes vibration studies, which indicate that no major disruptions to HL-LHC and SPS performance are expected during FPF excavation works. FPF construction, then, is not expected to interfere with the LHC and can proceed largely independent of the LHC schedule.

Since the last study, a site investigation, where a core was drilled to the depth of the FPF cavern, yielded broadly positive results, confirming the reliability of the Facility design. More detailed considerations of services have been incorporated, leading to a slight increase in size and an accompanying modest increase in cost for the civil engineering compared to previous estimates. The updated timeline for the works shows that the FPF could be constructed within a few years of approval, with no special R&D needed for the Facility design.

---

1



**CERN-PBC-NOTE 2024-003**  
4 July 2024  
davide.gamba@cern.ch

### Impact of Vibration to HL-LHC Performance During the FPF Facility Construction

**D. Gamba, H. Bartosik, M. Guinchard, J. A. Osborne, K. Pál, C. Vendeuvre, J. Wenninger, K. Widuch**  
CERN, CH-1211 Geneva, Switzerland

Keywords: excavation, forward physics facility, ground motion, tunnel deformation, vibration, FPF, LHC, HL-LHC, SPS

---

**Summary**

The Forward Physics Facility (FPF) is a proposed experimental site intended to be positioned approximately 630 meters from the ATLAS interaction point. It aims to capture long-lived particles and neutrinos that travel along the beam collision axis and fall outside the ATLAS detector's acceptance. The construction of this facility, particularly the excavation of the necessary shaft and cavern, could occur concurrently with beam operations at the CERN accelerator complex. Therefore, it is crucial to ensure that the ground motion resulting from these construction activities does not disrupt the normal functioning of the SPS and LHC. This study details how sensitive the SPS and LHC rings are to vibrations and misalignments close to the FPF construction site. It also examines the expected effects on beam operations, incorporating lessons learned from the HL-LHC infrastructure development near the ATLAS experiment, previous civil engineering projects, and established knowledge of slow ground movements in the vicinity.

---

1



**Steering Committee:** Jamie Boyd, Albert De Roeck, Milind Diwan, Jonathan Feng, Felix Kling

**Coordination Panel:** Aki Ariga, Alan Barr, Brian Batell, Jianming Bian, Jamie Boyd, Giovanni De Lellis, Albert De Roeck, Milind Diwan, Jonathan Feng, Chris Hill, Felix Kling, Juan Rojo, Dennis Soldin, Anna Stasto

**WG0 Facility:** Jamie Boyd

Physics WGs

**WG1 Neutrino Interactions:** Juan Rojo

**WG2 Charm Production:** Anna Stasto

**WG3 Light Hadron Prod:** Luis Anchordoqui, Dennis Soldin

**WG4 BSM:** Brian Batell, Sebastian Trojanowski

Detector WGs

**WG5 FASER2:** Alan Barr, Josh McFayden, Hide Otono

**WG6 FASERnu2:** Aki Ariga, Tomoko Ariga

**WG7 FLArE:** Jianming Bian, Milind Diwan

**WG9 FORMOSA:** Matthew Citron, Chris Hill

## Technical group:

Bringing together experts from CERN, BNL, and an external engineering contractor Bartoszek engineering:

BNL: M. Diwan, S. Linden, Y. Li, C. Miraval + L. Bartoszek

CERN: J. Boyd, J.P Corso (integration), A. Magazinik (integration), F. Resnati (neutrino platform)

Meeting bi-weekly on Fridays.

# Strongly Reviewed in Snowmass.

## Executive Summary (10 pages)

LHC

**The Energy Frontier (Science Drivers 1 – 3 & 5):** The Energy Frontier currently has a top-notch program with the Large Hadron Collider (LHC) and its planned High Luminosity upgrade (HL-LHC) at CERN, which sets the basis for the Energy Frontier vision. The fundamental lessons learned from the LHC thus far are that a Higgs-like particle exists at 125 GeV and there is no obvious and unambiguous signal of BSM physics. This implies that new physics either occurs at scales higher than we have probed, must be weakly coupled to the SM, or is hidden in backgrounds at the LHC. The immediate goal for the Energy Frontier is to continue to take and analyze the data from LHC Run 3, which will go on for about three more years, and carry out the 2014 P5 recommendations to complete the HL-LHC Upgrade and execute its physics program. The HL-LHC will measure the properties of the Higgs Boson more precisely, probe the boundaries of the SM further, and possibly observe new physics or point us in a particular direction for discovery.

A new aspect of the proposed LHC program is the emergence of a variety of auxiliary experiments that can use the interactions already occurring in the existing collision regions during the normal LHC and HL-LHC running of the ATLAS, CMS, LHCb, and ALICE experiments to explore regions of discovery space that are not currently accessible. These typically involve observing particles in the far forward direction or long-lived particles produced at larger angles but decaying far outside the existing detectors. These are mid-scale detectors in their own right and provide room for additional innovation and leadership opportunities for younger physicists at the LHC. The EF supports continued strong U.S. participation in the success of the LHC, and the HL-LHC construction, operations, and physics programs, including auxiliary experiments.


future collider

New colliders are the ultimate tools to extend the EF program into the next two decades thanks to the broad and complementary set of measurements and searches they enable. With a combined strategy of precision measurements and high-energy exploration, future lepton colliders starting at energies as low as the Z-pole up to a few TeV can shed substantial light on some of these key questions. It will be crucial to find a way to carry out experiments at higher energy scales, directly probing new physics at the 10 TeV energy scale and beyond. The EF supports a fast start for the construction of an  $e^+e^-$  Higgs Factory (linear or circular), and a significant R&D program for multi-TeV colliders (hadron and muon). The realization of a Higgs Factory will require an immediate, vigorous, and targeted accelerator and detector R&D program, while the study towards multi-TeV colliders will need significant and long-term investments in a broad spectrum of R&D programs for accelerators and detectors.

Finally, the U.S. EF community has expressed renewed interest and ambition to develop options for an energy-frontier collider that could be sited in the U.S., while maintaining its international collaborative partnerships and obligations with, for example, CERN.

*A new aspect of the proposed LHC program is the emergence of a variety of auxiliary experiments that can use the interactions already occurring in the existing collision ... to explore regions of discovery space that are not currently accessible. These typically involve observing particles in the far forward direction or long-lived particles ... decaying far outside the existing detectors. These are mid-scale detectors in their own right and provide room for additional innovation and leadership opportunities for younger physicists at the LHC. The EF supports continued strong U.S. participation ... including auxiliary experiments.*

# FASER2 Magnet





Two possible magnet solutions investigated for the FASER2 magnet:

- Custom made SC dipole based on (descoped) SAMURAI experiment magnet
  - 2Tm bending power, 1.7m x 1.7m aperture
  - SAMURAI magnet built by Toshiba, JP, but similar magnet could be built by TESLA, UK
  - Cost (Toshiba) 730 MJPY (4.1 MCHF today), 3-4y lead-time
- Off the shelf 'crystal puller' magnets:
  - 4 units: 0.4T central field 1.25m depth => 2Tm bending power (central region)
  - 1.6m diameter
  - Available from both Toshiba and TESLA
  - Cost (Toshiba) with modified mechanics to operate on side: 400 MJPY (2.3MCHF), 1-2y lead-time

- The FPF experiments have strong sensitivity in all of the dark sector PBC benchmark models
- In some of these the sensitivity is not as strong as that of SHiP
- There are several classes of models where the FPF is **more powerful than SHiP**

## MILLI-CHARGED PARTICLES


- The FPF accommodates a suite of experiments that can be optimized for various physics cases. This diversity is essential in probing a broad range of BSM physics possibilities.
- For example: FORMOSA, targeting milli-charged particles.
- Motivated by dark sectors with massless dark photons, but also new particles with magnetic or electric dipole moments, ...
- World-leading sensitivity for masses from  $\sim 100$  MeV to 100 GeV.
- Will not be probed by SHiP (and no fixed target experiment can produce particles with mass  $> 10$ -20 GeV).




- The FPF experiments have strong sensitivity in all of the dark sector PBC benchmark models
- In some of these the sensitivity is not as strong as that of SHiP
- There are several classes of models where the FPF is **more powerful than SHiP**

## STRONGLY-INTERACTING DARK SECTORS



- U(1) dark force  $\rightarrow$  dark photons, milli-charged particles.
- Any other dark force  $\rightarrow$  strongly-interacting dark sector. Dark particles (“quirks”) can be pair-produced at the LHC, but then oscillate down the beampipe, bound together by the dark color force.



- FASER2 can discover quirks with masses up to  $\sim$ TeV, as motivated by the gauge hierarchy problem (neutral naturalness).
- Requires LHC energies to produce new TeV particles, impossible to see at fixed target experiments.



Li, Pei, Ran, Zhang (2021); Li et al. (in progress)



Muon background flux estimate wd to be  $0.6 \text{ Hz/cm}^2$  at  $L=5e34 \text{ cm}^{-2} \text{ s}^{-1}$  in region 20cm from LOS.

Muon hotspots horizontally  $\pm 2 \text{ m}$  from the LOS.

Reducing the muon flux would be beneficial for some experiments, investigating the possibility of a sweeper magnet to reduce the flux.

Detailed FLUKA simulations used to:

- Estimate muon background
- Estimate radiation levels (RP)
- Estimate radiation to electronics / detector

All look encouraging for implemtning the FPF experiments

FLUKA simulation of Run 3 LHC setup validated at 20% level by FASER/SND@LHC data

(Note for HL-LHC much of the LSS changes => signifciantly reduces estimated muon flux per  $\text{fb}^{-1}$ )

Please find enclosed the first draft of the updated cost based on the results of the site investigations and the cost review done by ARUP. The estimated cost of the facility based on the existing design is 30.1MCHF, the estimate being a Class 4 (accuracy ranges being -15% to -30% on the low side, and +20% to +50% on the high side).

Ref.	Work Package	Cost [CHF]
1.	Underground Works	10,000,000.00
1.1	Preliminary activities	1,600,000.00
1.2	Access shaft	3,900,000.00
1.3	Experimental Cavern	4,500,000.00
2.	Surface Works	6,120,000.00
2.1	General items	640,000.00
2.2	Topsoil and earthworks	660,000.00
2.3	Roads and network	730,000.00
2.4	Buildings	4,090,000.00
2.4.1	Access building	2,000,000.00
2.4.2	Cooling and ventilation building	1,400,000.00
2.4.3	Electrical Building	490,000.00
2.4.5	External platforms	200,000.00
3.	General items	10,000,000.00
4.	Miscellaneous	4,000,000.00
	<b>TOTAL CE WORKS</b>	<b>30,120,000.00</b>

### Assumptions

1. Services not included
2. Technical galleries not included
3. Cranes not included
4. Access building as a conventional steel portal frame structure with cladding, only one floor
5. CV Building as a reinforced concrete building, only one floor
6. Finished floor level at 450m ASL
7. Sectional doors not included

Very preliminary estimate of cost increase to make cavern larger (to accommodate additional infrastructure etc.):


	Additional cost
5m longer cavern	700,000 CHF - 1,000,000.00 CHF
10m longer cavern	1,400,000 CHF - 2,000,000.00 CHF
Increase of the radius of the cavern by 1m	700,000 CHF - 1,000,000.00 CHF

Very preliminary costing of technical infrastructure for cavern at 10MCHF level.

**Total cost of facility (no experiments): ~40MCHF**

# Neutrinos at the FPF



LHC provides a **strongly collimated** beam of **TeV energy neutrinos** of **all three flavours** in the far forward direction.



Proposed FPF experiment have potential to detect O(1M) neutrinos:  
 O(**10<sup>6</sup>**) muon neutrinos, O(**10<sup>5</sup>**) electron neutrinos, O(**10<sup>4</sup>**) tau neutrinos



# Muon Background: Sweeper Magnet



Placing a sweeper magnet on the LOS can deflect these muons and reduce the background.

Best place for such a magnet would be between where LOS leaves LHC magnets and where it leaves the LHC tunnel (200m lever-arm for deflected muons).


Based on quick integration study required (will likely require some small local modifications to cryogenic infrastructure in the tunnel).



# Motivation



ANUBIS proposal:  
Extend fiducial volume  
→ harvest sensitivity!



'For free':  $\Delta\beta \approx 0.5\%$  (TOF)  
→ unique sensitivity to anomalous slow charged particles ( $dE/dx$ )


ANUBIS Detector requirements:

Parameter	Specification
Time resolution	$\delta t \lesssim 0.5$ ns
Angular resolution	$\delta\alpha \lesssim 0.01$ rad
Spatial resolution	$\delta x, \delta z \lesssim 0.5$ cm
Per-layer hit efficiency	$\varepsilon \gtrsim 98\%$

→ Adapt and exploit  
ATLAS Phase II RPC technology!

Ongoing RPC work at U Cambridge:

- RPC construction & characterisation
- Eco-gases for RPCs
- Quality assurance (Q/A) of ATLAS Phase II RPC
  - Gas leak studies
  - Mechanical stability
- Within DRD1 context



Adapted from ATL-PHYS-PUB-2022-007

Happy to share effort & expertise!

# Our involvement in ATLAS RPCs

- Our **current** involvement in **ATLAS RPCs**:
  - Quality Assurance (Q/A) & prospective Quality Checks (Q/C):
    - Study leak formation, electric faults & ageing in RPCs:
      - Apply mechanical stress through thermal cycling
        - Capitalise on our expertise in ITk Q/A & Q/C
        - USP: three (!) climate rooms at UCam
      - Our proposal endorsed by ATLAS RPC community
- Our **potential future** involvement in **ATLAS RPCs**:
  1. Q/C: commissioning tests on significant RPC fraction
  2. Contribute to installation & commissioning of Phase II RPCs
    - Need technicians (installation) & PDRAs (commissioning)
    - Project very short of experienced PRDAs
      - we could easily share expertise
      - build up expertise base in the UK
  3. Produce BOR/BOM type RPC chambers (80/120)
    - Establish RPC production line in UK!
  4. *Preserve UK leadership in ANUBIS also on detector front*



# proANUBIS location: similar to ANUBIS

