


# Compressed Baryonic Matter experiment at FAIR

Daniel Wielanek

Hanna Zbroszczyk

Warsaw University of Technology

# Road map



What are we pursuing and why?



Physics motivation






# QCD phase diagram

Low  $\mu_B$ , high  $T$ :

- Cross-over transition from hadronic to quark matter - comprehensive studies of QGP properties
- No critical point anticipated for  $\mu_B/T < 3$




High  $\mu_B$ , low  $T$ :

- Unknown phase structure (first-order phase transition, critical point possible, mixed phases, new phases, ...)
- Properties of matter to determine
- Characteristics of hadrons
- Equation of State (EoS) to establish
- Neutron Star (NS)

Bazavovet al. [HotQCD], PLB 795 (2019) 15-21  
 Dinget al., [HotQCD], PRL 123 (2019) 6, 062002  
 Borsanyi et al., PRL 125 (2020) 5, 052001  
 Isserstedt et al. PRD 100 (2019) 074011  
 Gao, Pawłowski, PLB 820 (2021) 136584

# NS puzzle




$$M_{\text{NS}} \approx 1 \div 2 M_{\odot}$$

$$R \approx 10\text{-}12 \text{ km}$$

$$\rho \approx 3 \div 5 \rho_0$$

$$\rho_0 \approx 2.8 \times 10^{14} \text{ g/cm}^3$$





- Observation of **NS** indicates their **mass**  $\sim 2M_{\odot}$   
(Shapiro-delay: Post-Keplerian parameters of orbits)
- **Hyperons:** Expected in core of NS, the conversion of N into Y is energetically favorable
- **Appearance of Hyperons:** The presence of Y alleviates Fermi pressure, resulting in a EoS and a reduction in NS mass (inconsistent with observations)

*Can they still be considered as components of NS?*


- **Proposed Solution:** A mechanism that provides additional pressure to ensure a stiffer EoS

One emergent mechanism involves many-body interactions, such as YN, YY, NNY, NYY

(Other: hypersonic three-body forces, Quark Matter Core - a transition to deconfined phase below hyperon threshold in density)



# NSM and HIC



Top row: simulation of NS mergers (NSM)

2 NSs of  $1.35 M_{\odot}$  each,

merging into a single object ( $2R \sim 10 \text{ km}$ ,  $n \sim 5n_0$ ,  $T \leq 20 \text{ MeV}$ ).


Overlap region:  $t \sim 20 \text{ ms}$ ,  $n \sim 2n_0$ ,  $T \sim 75 \text{ MeV}$

- max. temperature
- max. density


Bottom row: non-central Au+Au collision at  $\sqrt{s_{NN}} = 2.42 \text{ GeV}$

$n \simeq 3n_0$ ,  $T \simeq 80 \text{ MeV}$

HADES, *Nature Phys.* 15, 1040–1045 (2019)




Artist's depiction of a neutron star collision after inspiral, NASA/Swift/Dana Berry



Space and time scales vastly contrasting (km-NS / fm-HIC - 18 orders of magnitude; duration - 20 orders of magnitude)

Similar densities and temperatures achieved

# Road map



What are we pursuing and why?


Where are we now?




Physics motivation



Current status



# Critical point predictions




- LQCD frowns upon the location of the critical point at  $\mu_B/T < 3$
- Effective QCD and lattice-based theories estimate its location at  $T \sim 90 - 120$  MeV and  $\mu_B \sim 500 - 650$  MeV
- This corresponds to heavy-ion collisions at  $\sqrt{s_{NN}} \sim 3 - 5$  GeV
- The circumstance in which the critical point does not exist is also conceivable

DSE: Bernhardt, Fischer and Isserstedt, PLB 841 (2023)<sub>2</sub>  
 FRG: Fu, Pawłowski, Rennecke, PRD 101, 053032 (2020)<sub>3</sub>  
 BHE: Hippert et al., arXiv:2309.00579  
 lQCD-Pade: Basar, arXiv:2312.06952  
 lQCD-Pade: Clarke et al., PoS LATTICE2023 (2024),  
 Bazavov et al. [HotQCD], PLB 795 (2019) 15-21  
 Borsanyi et al. [Wuppertal-Budapest], PRL 125 (2020)  
 Cuteri, Philipsen, Sciarra, JHEP 11 (2021) 141  
 Vovchenko et al., PRD 97, 114030 (2018)



# Critical point searches



STAR CPOD 2024  
STAR, PRL 128 (2022) 20, 202303  
HADES, PRC 102 (2020) 2, 024914

$$\frac{\kappa_n(N_B - N_{\bar{B}})}{VT^3} = \frac{1}{VT^3} \frac{\partial^3 \ln Z(V, T, \mu_B)}{\partial (\mu_B/T)^n}$$

$\kappa_n$  experimentally measured


$$\kappa_n(N_B - N_{\bar{B}}) = \langle N_B \rangle + (-1)^n \langle N_{\bar{B}} \rangle = k_n(\text{Skellam})$$

- Non-monotonic trend in  $\kappa_4/\kappa_2$  of net-proton multiplicity distributions suggested as a **signature of the critical point**
- STAR collider program conducted comprehensive studies at  $\sqrt{s_{NN}} > 7.7$  GeV
- STAR fixed-target data investigation ongoing
- Sensitivity to the features of the QCD phase diagram increases with the order of the moment
- Higher-order moments requires prominent **statistics**

**Detailed systematics studies indispensable**




# E-M probes access the whole collision



Inscribes matter properties enabling estimation:

- degrees of freedom of the medium
- fireball's lifetime, temperature, acceleration, polarization
- transport properties
- restoration of chiral symmetry

late emission ← → early emission



Thermal dileptons in **LMR**:

- $T$  close to  $T_{ch}$  and  $T_{pc}$
- dominantly emitted around phase transition

Thermal dileptons in **IMR**:

- $T$  is higher than  $T_{pc}$
- Emitted fom QGP phase

$$\text{Effective size-signal: } S_{eff} \sim R \frac{S}{B}$$

R - interaction rate

S - signal

B- combinatorial background


**Prominent interaction rate mandatory**

EPJC (2009) 59 607-623

Nature Physics 15, 1040-1045 (2019)


JPS Conf.Proc. 21 (2020) 010079

# EoS to probe NS properties




EoS investigations include vast number of measurements:

- Chemistry (strangeness, charm, hyper nuclei, ...)
- Collectivity
- Vorticity
- Fluctuations and correlations
- Interactions in the final states (NN, NY, YY, many-body, hyper-nuclei, ...)




PRC 102 (2020) 34909 (STAR)

# EoS to probe NS properties




EoS investigations include vast number of measurements:

- Chemistry (strangeness, charm, hyper nuclei, ...)
- Collectivity
- Vorticity
- Fluctuations and correlations
- Interactions in the final states (NN, NY, YY, many-body, hyper-nuclei, ...)




PRC 102 (2020) 34909 (STAR)




PRL 113 (2014) 52302

# EoS to probe NS properties




EoS investigations include vast number of measurements:



- Chemistry (strangeness, charm, hyper nuclei, ...)
- Collectivity
- Vorticity
- Fluctuations and correlations
- Interactions in the final states (NN, NY, YY, many-body, hyper-nuclei, ...)



PRC 102 (2020) 34909 (STAR)




PRL 113 (2014) 52302




PRC 93 (2016) 14907 (STAR)

# EoS to probe NS properties




EoS investigations include vast number of measurements:


- Chemistry (strangeness, charm, hyper nuclei, ...)
- Collectivity
- Vorticity
- Fluctuations and correlations
- Interactions in the final states (NN, NY, YY, many-body, hyper-nuclei, ...)




PRC 102 (2020) 34909 (STAR)




PRL 113 (2014) 52302



STAR, PRL 112, 162301 (2014)




# EoS to probe NS properties





EoS investigations include vast number of measurements:



- Chemistry (strangeness, charm, hyper nuclei, ...)
  - Collectivity
  - Vorticity
  - Fluctuations and correlations
  - Interactions in the final states (NN, NY, YY, many-body, hyper-nuclei, ...)
- PLB 835, 137506 (2022)



PRC 102 (2020) 34909 (STAR)




PRL 113 (2014) 52302








# EoS to probe NS properties




EoS investigations include vast number of measurements:

- Chemistry (strangeness, charm, hyper nuclei, ...)
- Collectivity
- Vorticity
- Fluctuations and correlations
- Interactions in the final states (NN, NY, YY, many-body, hyper-nuclei, ...)





# EoS to probe NS properties




EoS investigations include vast number of measurements:

- Chemistry (strangeness, charm, hyper nuclei, ...)
- Collectivity
- Vorticity
- Fluctuations and correlations
- Interactions in the final states (NN, NY, YY, many-body, hyper-nuclei, ...)


PRC 96 (2017) no.2, 024911




STAR, PRL 112, 162301 (2014)



PRC 102 (2020) 34909 (STAR)




PRL 113 (2014) 52302



Invariant Mass ( $\text{GeV}/c^2$ )


# EoS to probe NS properties






EoS investigations include vast number of measurements:

- Chemistry (strangeness, charm, hyper nuclei, ...)
- Collectivity
- Vorticity
- Fluctuations and correlations
- Interactions in the final states (NN, NY, YY, many-body, hyper-nuclei, ...)



PRC 96 (2017) no.2, 024911



(a) QM2024, Nucl. Phys. A 259, (2024)



# Road map





# High $\mu_B$ facilities




CBM / HADES@ SIS100 (>2028)

MPD, BM@N@NICA


NA60@SPS(>2030)

NA61/SHINE@SPS

HADES@SIS18




STAR@RHIC




CBM / HADES:  
operations at  
 $\sqrt{s_{NN}} \sim 2 - 5 \text{ GeV}$

J-PARC-HI




CEE@HIAF (>2027)




T. Galatyuk, NPA 982 (2019), update 2024 [https://github.com/tgalatyuk/interaction\\_rate\\_facilities](https://github.com/tgalatyuk/interaction_rate_facilities)  
CBM, EPJA 53 3 (2017) 60

# Road map





# GSI, existing (upgraded to integrate with FAIR)




- Intensity
- Precision
- Antiproton beams

**FAIR,**  
under  
construction




# Facility for Antiproton and Ion Research





# Facility for Antiproton and Ion Research




SIS100 magnets,  
April, 2024



CBM cave,  
February, 2024



# Compressed Baryonic Matter experiment



Fixed-target experiment → highest rates achievable

Versatile subsystems → tailored for the physics program

Silicon-based tracking → fast and precise


Free-streaming front-end-electronics (FEE) → minimal dead-time while data acquisition

Online event selection → advanced data taking focused on customized needs


First beams in 2028/2029

**Years 1-3:** first energy scan, improved statistical uncertainties of factor 10 with respect to STAR

**Years 4-8:** high-statistics measurements: di-lepton IMR, ultra-rare probes




315 full members from  
10 countries  
47 full member  
institutions  
10 associated member  
institutions



**Campaign 2024:**  
 high-rate studies  
 online reconstruction  
 and selection  
 $\Lambda$  baryons in Ni+Ni at  
 1.0 - 1.93 AGeV


Free-streaming CBM data transport  
 Pre-series productions of all CBM detector systems  
 High-rate studies up to 10 MHz coll. rate in A+A collisions

## Rare signal reconstructed: $\Lambda \rightarrow p \pi^-$



Ni+Ni 1.93 AGeV  
 run 2391 (May '22):  
 10<sup>9</sup> collisions, 1:57h  
 400 kHz av. coll. rate


all detector systems involved  
 secondary vertex  
 velocity windows for p and  $\pi^-$   
 candidate




Free-streaming  
CBM data transport

Pre-series productions  
of all CBM detector  
systems

High-rate studies  
up to 10 MHz coll. rate  
in nucleus-nucleus  
collisions




## Rare signal reconstructed: $\Lambda \rightarrow p \pi^-$



Ni+Ni 1.93 AGeV  
run 2391 (May '22):  
 $10^9$  collisions, 1:57h  
400 kHz av. coll. rate


all detector systems involved  
secondary vertex  
velocity windows for p and  $\pi^-$   
candidate




Campaign 2024:  
high-rate studies  
online reconstruction  
and selection  
 $\Lambda$  baryons in Ni+Ni at  
1.0 - 1.93 AGeV



# Road map




# Key observables




## Systematic measurements:

- **Fluctuations:** System alteration through first-order phase transition, critical point
- **Dileptons :** Emissivity: system's lifetime, temperature, density, in-medium characteristics
- **Hadrons (Strangeness, Charm, Hyper-nuclei, Bound states):** EOS: vorticity, collectivity, correlations: NN, YN, YY, multi-body interactions




T. Galatyuk, NPA 982 (2019), update 2023  
[https://github.com/tgalatyuk/interaction\\_rate\\_facilities](https://github.com/tgalatyuk/interaction_rate_facilities)





HADES; *Nature Physics* 15 (2019) 10, 1040-1045

# Fluctuations



Corrections for volume fluctuations and conservation laws


- Event-by-event changes of efficiency
- Proper selection of  $y - p_T$  interval
- (Net-)baryons vs. protons, neutrons, nuclei



Expectations after  $\sim 3$  years of running

- Full coverage of  $\kappa_4(E)$  for protons
- First results of  $\kappa_6$
- Possible addition of strangeness:  $\kappa_4(\Lambda)$

# Dileptons




Electron thermal radiation, corrected for acceptance and efficiency,

Dominated by  $\rho$  contribution at LMR,




Can be reconstructed with 1.5-4.5% of precision,

Gives access to the fireball lifetime and electrical conductivity (transport properties)

$T$  vs. baryon density effects from partonic to hadronic fireballs






# Flow, polarization, correlations




Excellent acceptance coverage  
 Reconstruction efficiency  $\sim 30\%$

- Precise measurements of flow for  $S=1, 2, 3$
- Polarization measurement with precision of  $\sim 5\%$
- Thorough multi body  $N$  and  $Y$  correlations of  $S=1, 2, 3$  achievable









# Interactions: hyper-nuclei, bound-states




The most abundant production of hyper-nuclei anticipated at  $\sqrt{s_{NN}} \sim 2 - 5$  GeV

Prominent interaction rates and excellent particle identification will facilitate to search for multi-strange hyper-nuclei



Andronic et al., PLB 697 (2011)  
 Steinheimer et al., PLB 714 (2012)





To answer fundamental questions about the structure of the QCD phase diagram at high  $\mu_B$  and to explore neutron stars

## What are we pursuing and why?

## Where are we now?

Already operating at high  $\mu_B$  experiments are complete and exploration of new physics needs higher interaction rates

## Who is involved?

Many world-wide existing and planned facilities complement each other programs

## How to achieve the goal?

Compressed Baryonic Matter experiment with high interaction rates will explore the region of the energies of the highest importance

## What is the plan?

To start these exploration in 2028 and to answer fundamental questions in the first year of CBM running

# Postdoctoral Research Associate positions in Experimental Heavy-Ion Physics

Warsaw U. of Tech. (main) • Europe

hep-ex nucl-ex PostDoc • Experiments: [GSI-FAIR-CBM](#), [BNL-RHIC-STAR](#)

🕒 **Deadline on Dec 31, 2024**

<https://inspirehep.net/jobs/2811921>

## Job description:

The Heavy-Ion Reaction Group (HIRG) at the Faculty of Physics at Warsaw University of Technology participates in the experiments STAR at BNL, CBM at FAIR, HADES at GSI, ALICE, and NA61/SHINE at CERN.

The STAR, HADES, and CBM groups specialize in two-particle femtoscopic correlation analysis measurements. We closely cooperate with the ALICE group at WUT.

The successful candidates will work with Professor Hanna Zbroszczyk on the STAR experiment at RHIC, the HADES experiment at SIS18, or the [CBM experiment at SIS100](#), focusing on studies of two-particle correlations. One position can relate up to 2 experiments. Responsibilities include data analysis and publication of results, collaboration service work, mentoring students, and supporting the Heavy Ion Reaction Group's research activities. Occasional travel to BNL and/or GSI will be required. The successful candidates are expected to lead studies of femtoscopic correlations in the search for STAR, HADES, or CBM experiments. The contract can be extended up to 24 months, provided a satisfactory evaluation outcome after the first 12 months.


## Duties and Responsibilities:

- Taking part in the analysis of heavy-ion collision RHIC data recorded by the STAR detector to study two-particle correlations or
- Taking part in the analysis of heavy-ion collision SIS-18 data recorded by the HADES detector to study two-particle correlations or
- Taking part in the Monte Carlo data analysis for the CBM experiment
- Joining the STAR/HADES or CBM collaborations and the relevant Physics Working Group, participating in weekly deliberations and active participation in its meetings (at least every few weeks);
- Taking part in data-taking (experimental shifts);
- Presentation of results at meetings and conferences, as well as writing scientific articles and publishing papers in peer-reviewed journals;
- Maintaining close cooperation with colleagues in the PWG group will be essential for the progress of all stages of the project

- Thank you



# Current prospects and timeline



← until 2028  
**FAIR Phase-0**

← after 2028

Staged implementation recommended by the Heuer/Tribble Commission's report (2022) with the First Science stage endorsed by the FAIR Council as "the most appropriate starting scenario to achieve world-leading science."

37

