

Exploring the shape of QGP droplet using high- p_{\perp} tomography

Bithika Karmakar

Incubator of Scientific Excellence - Centre for Simulations of Superdense Fluids
University of Wrocław, Poland

Zimányi School 2024

Collaborators: Dusan Zigic, Jussi Auvinen, Pasi Huovinen, Marko Djordjevic and Magdalena
Djordjevic

Contents

- Introduction
- High- p_{\perp} energy loss: DREENA
- Probing the initial profiles
- Study of the early evolution
- Summary

Introduction

- Quark Gluon Plasma (QGP) is created in ultrarelativistic heavy-ion collisions.
- Consists of interacting quarks, antiquarks and gluons.
- Low- p_{\perp} ($p_{\perp} \leq 5\text{GeV}$) observables are generally used to study the medium properties.
- High- p_{\perp} partons propagate through the medium. Jet loses energy by interacting with the medium.
- The rare high- p_{\perp} particles can also become excellent probe of the QCD matter.

Initial condition model


- Initial condition models (e.g., Glauber, MC-KLN, IP Glasma etc) aim to determine the initial energy or entropy deposition.
- Bayesian analysis has been used to constrain the initial conditions using T_{RENTo} model. Nature Phys. 15, no. 11, 1113-1117 (2019)
- For a pair of projectiles A and B colliding along z axis, the participant thickness is $T_{A,B}(x, y) = \int dz \rho_{A,B}^{part}(x, y, z)$.
- The initial entropy density is $s(x, y) \propto T_R(p; T_A, T_B) = \left(\frac{T_A^p + T_B^p}{2} \right)^{1/p}$.
- $p = 0$ is found to be preferred in Bayesian analysis when initial free streaming is considered.

Probing the shape of QGP droplet

- Initial free streaming not favored by high- p_{\perp} data.
S. Stojku, J. Auvinen, M. Djordjevic, P. Huovinen and M. Djordjevic, Phys. Rev. C **105** (2022) 2, L021901
- Can lead to different results without free streaming.
- DREENA can be used to constrain the initial profiles with no free streaming.
- We use $p \in \{1/3, 0, -1/3, -2/3, -1\}$ to see which value the high- p_{\perp} data prefers.
- Other parameters of the bulk evolution are tuned to agree with the low- p_{\perp} data.
- We use constant η/s in each case.



Averaged initial temperature ($\tau = 1\text{fm}$)

BK, D. Zigic, P. Huovinen, M. Djordjevic, M. Djordjevic and J. Auvinen, Phys. Rev. C **110** (2024) 4, 044906



Initial state anisotropy

BK, D. Zigic, P. Huovinen, M. Djordjevic, M. Djordjevic and J. Auvinen, Phys. Rev. C **110** (2024) 4, 044906



Generalized DREENA-A (Adaptive)

- **Dynamical Radiative and Elastic ENergy loss Approach**

- Based on finite temperature field theory and generalized HTL approach

M. Djordjevic, PRC **74**, 064907, (2006) ; PRC **80**, 064909 (2009), M. Djordjevic and U. Heinz, PRL **101**, 022302

- Takes into account both radiative and collisional energy losses

- No fitting parameter

- $$\frac{E_f d^3 \sigma_q(H_Q)}{dp_f^3} = \frac{E_i d^3 \sigma(Q)}{dp_i^3} \otimes P(E_i \rightarrow E_f) \otimes D(Q \rightarrow H_Q),$$

- $$\frac{E_f d^3 \sigma_u(H_Q)}{dp_f^3} = \frac{E_i d^3 \sigma(Q)}{dp_i^3} \otimes D(Q \rightarrow H_Q).$$

- Includes event-by-event fluctuating arbitrary temperature profiles.

- Agrees with R_{AA} , high- p_{\perp} v_2 , v_3 . Predictions vastly underestimates high- p_{\perp} v_4 (High- p_{\perp} v_4 puzzle), allows to extract bulk medium properties. D. Zigic, J. Auvinen, I. Salom, M. Djordjevic and P.


Huovinen, Phys. Rev. C **106** (2022) 4, 044909

Dusan Zigic, Wed 15:00

Bojana Illic, Fri 11:50

Charged hadron R_{AA} and high- p_{\perp} v_n in Pb+Pb $\sqrt{s} = 5.02$ TeV

BK, D. Zigic, P. Huovinen, M. Djordjevic, M. Djordjevic and J. Auvinen, Phys. Rev. C **110** (2024) 4, 044906




Study of early evolution of heavy-ion collision

- Although high- p_{\perp} data prefer no initial free streaming and delayed onset of hydrodynamics, it may appear unrealistic.
- We try to accommodate earlier onset of hydrodynamic evolution by considering the highly anisotropic initial profiles ($T_{\text{RENTo}} p < 0$).
- We consider $\tau_0 = 1 \text{ fm}, 0.6 \text{ fm}$ and 0.2 fm .
- Readjusted η/s and T_{RENTo} normalization to reproduce the low- p_{\perp} data.

Charged hadron R_{AA} and high- p_{\perp} v_n in Pb+Pb $\sqrt{s} = 5.02$ TeV

BK, D. Zigic, P. Huovinen, M. Djordjevic, M. Djordjevic and J. Auvinen, Phys. Rev. C **110** (2024) 4, 044906




Summary

- The shape of the initial state has been modulated by varying the parameter p of T_RENTo model.
- It is found that $p \approx 0$ gives the best overall fit which is consistent with the Bayesian analysis of low- p_{\perp} data.
- We tested if larger anisotropy of the initial profiles ($p \ll 0$) would allow an earlier onset of fluid dynamical evolution.
- Lower values of p enhance R_{AA} and high- p_{\perp} v_2 , the enhancement is insufficient for facilitating an earlier onset of transverse expansion.

Extreme QCD 2025

- **Dates:** July 2 – 4, 2025
- **Venue:** University of Wroclaw, Wroclaw, Poland
- **Student support:** fee, travel, accommodation
- **Web:** INDICO
INSPIRE



XQCD 2025



Uniwersytet
Wrocławski

Thank you for your attention



European Research Council
Established by the European Commission

$$\epsilon_{m,n} = \frac{\int dx dy r^m \cos [n(\phi - \Psi_{m,n})] e(x, y)}{\int dx dy r^m e(x, y)}$$

$$\Psi_{m,n} = \frac{1}{n} \arctan \frac{\int dx dy r^m \sin(n\phi) e(x, y)}{\int dx dy r^m \cos(n\phi) e(x, y)} + \pi/n$$

Early evolution using DREENA-A

S. Stojku, J. Auvinen, M. Djordjevic, P. Huovinen and M. Djordjevic, Phys. Rev. C **105** (2022) 2, L021901

1. **Cyan** $\rightarrow \tau_q = \tau_0 = 0.2fm$

2. **Orange** $\rightarrow \tau_0 = 0.2fm; \tau_q = 1fm$

3. **Red** \rightarrow FS; $\tau_0 = \tau_q = 1fm$

4. **Black** $\rightarrow \tau_0 = \tau_q = 1fm$


- Fits low- p_{\perp} data well

- Divergent is disfavored by R_{AA} data

- Delaying τ_q hardly changes v_2

- Early FS does not fit data as well

- v_2 predictions approach data when $\tau_0 = \tau_q = 1fm$ (No early free steaming)



Pb + Pb $\sqrt{s} = 5.02$ TeV