

COMETA workshop on EFT in HH and VBS

Monday 17 March 2025 - Wednesday 19 March 2025

Paris

Book of Abstracts

Contents

Workshop introduction	1
VBS and HH theory review	1
VBS experimental review	1
HH experimental review	1
Discussion	1
Review of VBS EFT MC modelling in ATLAS and CMS	1
Review of HH EFT MC modelling in ATLAS and CMS	1
ML for reweighting	1
Pheno and theory overview : HH and VBS interplay, available tools	2
Discussion	2
Publishing experimental results	2
Discussion	2
VBS measurement in semileptonic final states with ATLAS Run-2 data	2
HEFT Multi-Higgs Production in VBF	3
Double Higgs Production in Vector Boson Fusion at NLO QCD in HEFT	3
The effect of single Higgs backgrounds and Higgs branching fractions in the SMEFT interpretation	4
Double Higgs production via VBF NLO in SMEFT	4
A novel implementation of the Matrix Element Method at next-to-leading-order (NLO) for the measurement of the Higgs tri-linear coupling in di-Higgs production at the LHC	4
VBS aQGC Reweighting study	5
The Art of Counting: where to cut-off SMEFT and HEFT in Higgs Pair Production	6
RGE effects in SMEFT predictions for HH production	6
HHH and HH production as probes of the Higgs self-couplings	6

Triple and double VBF-Higgs production as probes of SMEFT/HEFT interactions	6
SMEFT effects and CP violation in diboson production via gluon-fusion	6
VBS constraints on Higgs interactions in HEFT [TBC]	7
TBC	7

1

Workshop introduction

EFT Review : theory and experimental landscape / 2

VBS and HH theory review

Corresponding Author: alejonahuel.rossia@unipd.it

EFT Review : theory and experimental landscape / 3

VBS experimental review

Corresponding Authors: giacomo.boldrini@cern.ch, despoina.sampsonidou@cern.ch

EFT Review : theory and experimental landscape / 4

HH experimental review

Corresponding Authors: atalierc@cern.ch, laura.pereira.sanchez@cern.ch

EFT Review : theory and experimental landscape / 5

Discussion

Modelling of EFT effects / 6

Review of VBS EFT MC modelling in ATLAS and CMS

Corresponding Authors: costanza.carrivale@pg.infn.it, costanza.carrivale@cern.ch, nicolas.morange@cern.ch

Modelling of EFT effects / 7

Review of HH EFT MC modelling in ATLAS and CMS

Corresponding Authors: jona.motta@cern.ch, tom.ingebretsen.carlson@cern.ch

Modelling of EFT effects / 8

ML for reweighting

Modelling of EFT effects / 9

Pheno and theory overview : HH and VBS interplay, available tools

Corresponding Author: gudrun.heinrich@kit.edu

Modelling of EFT effects / 10

Discussion

Publishing and exploiting the results / 11

Publishing experimental results

Corresponding Authors: adinda.maite.de.wit@cern.ch, ken.mimasu@cern.ch

Track/session:

Publishing and exploiting the results / 12

Discussion

Early career scientists presentations / 13

VBS measurement in semileptonic final states with ATLAS Run-2 data

Author: Mathieu Markovitch¹

¹ *Université Paris-Saclay (FR)*

Corresponding Author: mathieu.markovitch@cern.ch

Vector boson scattering (VBS) processes probe the fundamental structure of electroweak (EW) interactions and provide a high sensitivity to new physics phenomena affecting gauge and Higgs couplings. The semileptonic final states, where one of the scattered electroweak boson decays hadronically into a quark/antiquark pair and the other boson decays leptonically into electrons, muons or neutrinos, has high statistics in the WW, WZ and ZZ processes. The hadronically decaying gauge

boson can be reconstructed as two small-radius jets or one large-radius jet in case of high boost. Semileptonic final states VBS was studied in an ATLAS Run-2 analysis using 139/fb of proton collisions at a center-of-mass energy of 13 TeV, allowing to measure the fiducial cross section for EW diboson production accompanied with jets in semileptonic final states. The results are also interpreted in an Effective Fields Theories (EFT) framework to set constraints on anomalous Quartic Gauge Couplings (aQGC) through dimension-8 operators.

Track/session:

Plenary track

Theory parallel discussion / 14

HEFT Multi-Higgs Production in VBF

Authors: Alexandre Salas-Bernárdez¹; Javier Martínez Martín²; Juan José Sanz-Cillero²; Rafael Delgado López^{None}; Raquel Gomez Ambrosio^{None}

¹ *Universidad Complutense de Madrid*

² *Universidad Complutense de Madrid & IPARCOS*

Corresponding Authors: javmar21@ucm.es, raquel.gomezambrosio@unito.it, rafael.delgado@upm.es, alexsala@ucm.es, jjsanzcillero@ucm.es

We have studied the production of 2, 3 and 4 Higgs bosons from the collision of Goldstone bosons using the equivalence theorem in HEFT. This work has been compared with results from both ATLAS and CMS. The measurements are being updated by the collaborations with the latest results.

Track/session:

Parallel Theory track

Early career scientists presentations / 15

Double Higgs Production in Vector Boson Fusion at NLO QCD in HEFT

Authors: Gudrun Heinrich¹; Jens Braun²; Marius Höfer^{None}; Pia Bredt³

¹ *KIT*

² *Karlsruhe Institute of Technology (KIT)*

³ *Siegen University*

Corresponding Authors: gudrun.heinrich@kit.edu, marius.hoefer@kit.edu, pia.bredt@uni-siegen.de, jens.braun2@kit.edu

We present a calculation of the NLO QCD corrections to Higgs boson pair production in vector boson fusion, combined with the leading operators parametrising anomalous interactions in non-linear Effective Field Theory (HEFT). Based on our Monte Carlo implementation using GoSam+Whizard, we investigate the effects of anomalous Higgs couplings on various observables.

Track/session:

Plenary track

ATLAS internal discussion / 16

The effect of single Higgs backgrounds and Higgs branching fractions in the SMEFT interpretation

Author: Ligang Xia¹

¹ *Nanjing University (CN)*

Corresponding Author: ligang.xia@cern.ch

Some SMEFT parameters contribute to both di-Higgs and single-Higgs productions. Taking the HH->bbyy channel (JHEP 01(2024) 066) as example, we present the results on the effects of single Higgs backgrounds and Higgs branching fractions and discuss about the evaluation of systematical uncertainties.

Track/session:

Parallel ATLAS track

Theory parallel discussion / 17

Double Higgs production via VBF NLO in SMEFT

Author: Michal Ryczkowski¹

¹ *Università di Padova and INFN, Sezione di Padova*

Corresponding Author: michaljakub.ryczkowski@unipd.it

The process of double Higgs boson production via vector boson fusion (VBF), although strongly suppressed compared to the gluon fusion channel, remains an important area of study. Not only can it provide valuable insights into the EW sector and the mechanism of EWSB, but it is also highly sensitive to BSM physics. In this talk, we explore the impact of dimension-6 and dimension-8 SMEFT operators on VBF double Higgs production. We estimate and present the maximal effects, taking into account available constraints on SMEFT Wilson coefficients from fits to experimental data, perturbative unitarity constraints and the validity of the EFT expansion.

Track/session:

Parallel Theory track

Early career scientists presentations / 18

A novel implementation of the Matrix Element Method at next-to-leading-order (NLO) for the measurement of the Higgs tri-linear coupling in di-Higgs production at the LHC

Authors: Jan Stark¹; Matthias Jean Tartarin¹

¹ *Laboratoire des 2 Infinis - Toulouse, CNRS / Univ. Paul Sabatier (FR)*

Corresponding Authors: stark@in2p3.fr, matthias.tartarin@l2it.in2p3.fr

One of the LHC's priorities, following the discovery of the Higgs boson, is to observe the production of Higgs pairs and to measure the Higgs tri-linear coupling λ_{3H} .

Due to the rarity of di-Higgs production, measuring λ_{3H} has proven to be highly challenging. Exclusion limits have been observed using a variety of approaches, including cut-based methods and boosted decision trees (BDTs).

To address this difficulty from a new perspective, our work explores the application of the Matrix Element Method (MEM), a technique that has demonstrated its effectiveness in multiple analyses in which measurements were performed in processes that were rare (at the time). One can mention the primordial role of the MEM in measuring the top-quark mass at Tevatron, or its role in the first evidence for single top production in the s-channel at the LHC for example.

The MEM is a statistically optimal multivariate method that maximizes the utilization of both the experimental and theoretical information available to an analysis, making it inherently well-suited to rare process searches and Standard Model measurements at particle accelerators like the LHC.

The MEM avoids the application of strict selection requirements to the data. It accounts for the entire dataset (with minimal preselection), which is very important when dealing with such rare processes.

Most MEM studies have been limited to leading-order (LO) accuracy, with extensions to next-to-leading-order (NLO) explored only in specific cases due to the additional complexities introduced by virtual and real contributions. Building a MEM at LO is already a highly challenging task, and incorporating NLO formalism substantially amplifies this difficulty due to the increased computational demands and complex theoretical requirements.

To contribute to the measurement of λ_{3H} from LHC data in the $gg \rightarrow HH \rightarrow b\bar{b}\gamma\gamma$ channel, we developed a MEM framework by working on a new NLO implementation (which can be applied in many more analyses).

This MEM framework utilizes state-of-the-art Matrix Elements at NLO from the POWHEG-BOX-V2 [1] and MG5 _@NLO software packages, and can be extended to include Higgs boson pair production at full NLO QCD in SMEFT [2].

The framework is implemented within a modified version of MoMEMta, a software designed for managing multi-variable phase-space integration, which has been extended to incorporate this new NLO implementation.

To our knowledge, this work marks the first application of the MEM at NLO accuracy to the search for HH and the measurement of λ_{3H} . This also represents the first application of the MEM using this new NLO formalism.

We have demonstrated that this NLO framework is functional and effective.

We also applied this framework to Monte Carlo (MC) simulated samples in a search for λ_{3H} , achieving promising results. This study aims to introduce this new approach to the community and position the MEM (at NLO) as a competitive alternative to other established methods to determine the Higgs self-coupling λ_{3H} .

References:

[1] G. Heinrich, S.P. Jones, M. Kerner, G. Luisoni, L. Scyboz

Probing the trilinear Higgs boson coupling in di-Higgs production at NLO QCD including parton shower effects, JHEP 06 (2019) 066 [<https://inspirehep.net/literature/1725744>]

[2] G. Heinrich, J. Lang

Renormalisation group effects in SMEFT for di-Higgs production, [<https://arxiv.org/pdf/2409.19578>]

Track/session:

Plenary track

ATLAS internal discussion / 19

VBS aQGC Reweighting study

Author: Olivier Salin¹

¹ *Université Paris-Saclay*

Corresponding Author: olivier.salin@ens-paris-saclay.fr

This talk will present an investigation into generating EFT aQGC samples using the internal Mad-graph LO matrix element reweighting method. The study focuses on the semi-leptonic VBS process applied to the new aQGC recommendation superset of the Eboli model. The aim is to assess the accuracy and reliability of the approach and, if possible, validate it as an alternative to the EFT decomposition method commonly used in ATLAS. If successful, this method could significantly reduce the number of EFT samples required.

Track/session:

Parallel ATLAS track

Theory parallel discussion / 21

The Art of Counting: where to cut-off SMEFT and HEFT in Higgs Pair Production

Corresponding Author: konstantin.schmid@pd.infn.it

Theory parallel discussion / 22

RGE effects in SMEFT predictions for HH production

Corresponding Author: stefano.dinoi@studenti.unipd.it

Theory parallel discussion / 23

HHH and HH production as probes of the Higgs self-couplings

Corresponding Author: panagiotis.stylianou@desy.de

Theory parallel discussion / 24

Triple and double VBF-Higgs production as probes of SMEFT/HEFT interactions

Corresponding Author: anisha@kit.edu

Theory parallel discussion / 25

SMEFT effects and CP violation in diboson production via gluon-fusion

Corresponding Author: marion.thomas@manchester.ac.uk

Theory parallel discussion / 26

VBS constraints on Higgs interactions in HEFT [TBC]

Corresponding Author: domenec.espriu@cern.ch

Theory parallel discussion / 27

TBC