


# Phenomenology of Dirac neutrino EFT

---

Based on 2411.17414

Anirban Biswas, **Eung Jin Chun**,  
Sanjoy Mandal, Dibyendu Nanda



# Outline

- Construct EFT with RHN assuming neutrinos are Dirac:
  - Impose B-L invariance
  - Phenomenological studies on  $D \leq 6$  BNC & BNV operators
- EWPT; LHC limits from  $pp \rightarrow l/j + \text{MET}$ ; Decays of  $t, \tau, M \rightarrow X + \text{MET}$
- Neutrino masses from effective operators
- Stellar cooling by emitting  $N$
- Contribution to  $\Delta N_{\text{eff}}$
- Proton decay from BNV operators

# Defining Lagrangian

- Introduce  $N = N_R$  with B-L charge  $-1$ .

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + i\bar{N}\gamma^\mu\partial_\mu N - \left(Y_\nu\bar{L}\tilde{H}N + \text{h.c.}\right) + \frac{1}{\Lambda^2} \sum_i^n C_i^{(6)} \mathcal{O}_i^{(6)}$$

- Assume B-L is a good quantum number and thus  $N$  has no Majorana mass.
- Tiny Dirac neutrino Yukawa  $Y_\nu = \frac{m_\nu}{\langle H \rangle} \sim 10^{-13}$  may be a consequence of Dirac Seesaw.

# BNC operators

1612.04527

$\psi^2 H X$	$\mathcal{O}_{LNB}(+\text{h.c.}) = \bar{L}\sigma^{\mu\nu} N \tilde{H} B_{\mu\nu}$ $\mathcal{O}_{LNW}(+\text{h.c.}) = \bar{L}\sigma^{\mu\nu} N \sigma_I \tilde{H} W_{\mu\nu}^I$
$\psi^2 H^2 D$	$\mathcal{O}_{HN} = \bar{N}\gamma^\mu N (H^\dagger i \overleftrightarrow{D}_\mu H)$ $\mathcal{O}_{HN_e}(+\text{h.c.}) = \bar{N}\gamma^\mu e_R (\tilde{H}^\dagger i D_\mu H)$
$\psi^2 H^3$	$\mathcal{O}_{LNH} = \bar{L} \tilde{H} N (H^\dagger H)$
RRRR	$\mathcal{O}_{NN} = (\bar{N}\gamma_\mu N)(\bar{N}\gamma^\mu N)$ $\mathcal{O}_{eeNN} = (\bar{e}_R\gamma_\mu e_R)(\bar{N}\gamma^\mu N)$ $\mathcal{O}_{uuNN} = (\bar{u}_R\gamma_\mu u_R)(\bar{N}\gamma^\mu N)$ $\mathcal{O}_{ddNN} = (\bar{d}_R\gamma_\mu d_R)(\bar{N}\gamma^\mu N)$ $\mathcal{O}_{duNe}(+\text{h.c.}) = (\bar{d}_R\gamma_\mu u_R)(\bar{N}\gamma^\mu e_R)$
LLRR	$\mathcal{O}_{LLNN} = (\bar{L}\gamma_\mu L)(\bar{N}\gamma^\mu N)$ $\mathcal{O}_{QQNN} = (\bar{Q}\gamma_\mu Q)(\bar{N}\gamma^\mu N)$
LRLR	$\mathcal{O}_{LNLe}(+\text{h.c.}) = (\bar{L}N)\epsilon(\bar{L}e_R)$ $\mathcal{O}_{LNQd}(+\text{h.c.}) = (\bar{L}N)\epsilon(\bar{Q}d_R)$ $\mathcal{O}_{LdQN}(+\text{h.c.}) = (\bar{L}d_R)\epsilon(\bar{Q}N)$
LRRL	$\mathcal{O}_{QuNL}(+\text{h.c.}) = (\bar{Q}u_R)(\bar{N}L)$

# BNV operators ( $B - L = 0$ )

$$\mathcal{O}_1 = \left( \overline{d_{\alpha R}^c} u_{\beta R} \right) \left( \overline{Q_{i\gamma L}^c} L_{jL} \right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij},$$

$$\mathcal{O}_2 = \left( \overline{Q_{i\alpha L}^c} Q_{j\beta L} \right) \left( \overline{u_{\gamma R}^c} e_R \right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij},$$

$$\mathcal{O}_3 = \left( \overline{Q_{i\alpha L}^c} Q_{j\beta L} \right) \left( \overline{Q_{k\gamma L}^c} L_{lL} \right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij} \epsilon_{kl},$$

$$\mathcal{O}_4 = \left( \overline{d_{\alpha R}^c} u_{\beta R} \right) \left( \overline{u_{\gamma R}^c} e_R \right) \epsilon_{\alpha\beta\gamma}.$$

$$\mathcal{O}_{N1} = \left( \overline{Q_{i\alpha L}^c} Q_{j\beta L} \right) \left( \overline{d_{\gamma R}^c} N_R \right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij},$$

1612.04527

$$\mathcal{O}_{N2} = \left( \overline{u_{\alpha R}^c} d_{\beta R} \right) \left( \overline{d_{\gamma R}^c} N_R \right) \epsilon_{\alpha\beta\gamma}.$$

# EWPT limits

$$\Gamma^{\text{NP}}(Z \rightarrow \text{inv}) = \frac{m_Z^3 v^2}{8\pi\Lambda^4} (C_{HN}^2 + 2C_{LNZ}^2),$$

$$\Gamma^{\text{NP}}(h \rightarrow \text{inv}) = \frac{3m_h v^4}{16\pi\Lambda^4} C_{LNH}^2,$$

$$\Delta\Gamma(W \rightarrow l\nu) = \frac{m_W^3 v^2}{48\pi\Lambda^4} C_{HNe}^2$$



$$C = 1$$

$$\Gamma(Z \rightarrow \text{inv}) < 2 \text{ MeV} \Rightarrow \Lambda_{HN, LNZ} > 1 \text{ TeV}$$


$$\text{BR}(h \rightarrow \text{inv}) < 10 \% \Rightarrow \Lambda_{LNH} > 2.8 \text{ TeV}$$

$$\Delta\Gamma(W \rightarrow l\nu) < 2 \times 10^{-3} \Gamma(W \rightarrow l\nu) \Rightarrow \Lambda_{HNe} > 0.58 \text{ TeV}$$

# Collider searches: $pp \rightarrow l/j + \text{MET}$




# Top decay




Top decays	Relevant Coefficients	$\Lambda_{\min}$ [TeV]
$t \rightarrow \ell_\alpha b + \text{Inv}$	$C_{HN_e}^{i\alpha}, C_{duNe}^{33i\alpha}, C_{LdQN}^{\alpha33i}, C_{LNQd}^{\alpha i33}, C_{QuNL}^{33i\alpha}$	0.139, 0.139, 0.137

# Meson decay: $M^\pm \rightarrow l^\pm N$



Meson decay	Decay width [GeV]	Relevant Coefficients	$\Lambda_{\min}$ [TeV]
$\pi^+ \rightarrow \mu^+ \nu_\mu$	$(2.5281 \pm 0.0005) \times 10^{-17}$	$C_{duNe}^{111\mu}, C_{HNe}^{i\mu}, C_{QuNL}^{11i\mu}, C_{LNQd}^{\mu i11}$	2.45
$\pi^+ \rightarrow e^+ \nu_e$	$(3.110 \pm 0.010) \times 10^{-21}$	$C_{duNe}^{111e}, C_{HNe}^{ie}, C_{QuNL}^{11ie}, C_{LNQd}^{ei11}$	1.23
$K^+ \rightarrow \mu^+ \nu_\mu$	$(3.379 \pm 0.008) \times 10^{-17}$	$C_{duNe}^{21i\mu}, C_{HNe}^{i\mu}, C_{QuNL}^{21i\mu}, C_{LNQd}^{\mu i12}$	1.32
$K^+ \rightarrow e^+ \nu_e$	$(8.41 \pm 0.04) \times 10^{-22}$	$C_{duNe}^{21ie}, C_{HNe}^{ie}, C_{QuNL}^{21ie}, C_{LNQd}^{ei12}$	1.12
$D_s^+ \rightarrow \mu^+ \nu_\mu$	$(7.09 \pm 0.20) \times 10^{-15}$	$C_{duNe}^{22i\mu}, C_{HNe}^{i\mu}, C_{QuNL}^{22i\mu}, C_{LNQd}^{\mu i22}$	0.73
$D_s^+ \rightarrow \tau^+ \nu_\tau$	$(6.95 \pm 0.15) \times 10^{-14}$	$C_{duNe}^{22i\tau}, C_{HNe}^{i\tau}, C_{QuNL}^{22i\tau}, C_{LNQd}^{\tau i22}$	0.77
$B^+ \rightarrow \tau^+ \nu_\tau$	$(4.4 \pm 1.0) \times 10^{-17}$	$C_{duNe}^{31i\tau}, C_{HNe}^{i\tau}, C_{QuNL}^{31i\tau}, C_{LNQd}^{\tau i13}$	0.44
$D^+ \rightarrow \mu^+ \nu_\mu$	$(2.38 \pm 0.11) \times 10^{-16}$	$C_{duNe}^{12i\mu}, C_{HNe}^{i\mu}, C_{QuNL}^{12i\mu}, C_{LNQd}^{\mu i21}$	0.63
$D^+ \rightarrow \tau^+ \nu_\tau$	$(7.6 \pm 1.7) \times 10^{-16}$	$C_{duNe}^{12i\tau}, C_{HNe}^{i\tau}, C_{QuNL}^{12i\tau}, C_{LNQd}^{\tau i21}$	0.72

# Tau decay



Tau decays	Decay width [GeV]	Relevant Coefficients	$\Lambda_{\min}$ [TeV]
$\tau \rightarrow \pi + \text{inv}$	$(2.453 \pm 0.012) \times 10^{-13}$	$C_{duNe}^{11i\tau}, C_{QuNL}^{11i\tau}, C_{LNQd}^{\tau i11}$	1.04
$\tau \rightarrow K + \text{inv}$	$(1.578 \pm 0.023) \times 10^{-14}$	$C_{duNe}^{21i\tau}, C_{QuNL}^{21i\tau}, C_{LNQd}^{\tau i12}$	1.65

Tau decays	Relevant Coefficients	$\Lambda_{\min}$ [TeV]
$\tau^- \rightarrow l^- + \cancel{E}$	$C_{HNe}^{il/\tau}, C_{LNW}^{li/\tau i}, C_{LNLe}^{\tau il(\tau/l)}, C_{LNLe}^{li\tau(\tau/l)}, C_{LNLe}^{ij(l\tau/\tau l)}, C_{LNLe}^{lj\tau}, C_{LNLe}^{\tau jil}, C_{eeNN/LLNN}^{l\tau ij}$	1.5 (1.49)

# Neutrino mass & magnetic moment

- After EWSB,  $\mathcal{O}_{LNH}$  and  $\mathcal{O}_{LNB/W}$  contribute to neutrino mass and magnetic moment, respectively:

$$\delta m_\nu = C_{LNH}(v) \frac{v^3}{2\sqrt{2}\Lambda^2} \quad C=1 \Rightarrow \Lambda_{LNH} \gtrsim 2 \times 10^8 \text{ GeV}$$

$$\frac{\mu_{\nu N}}{\mu_B} = \frac{2\sqrt{2}m_e v}{e\Lambda^2} (c_w C_{LNB}(v) + s_w C_{LNW}(v)) < 10^{-12} \quad (\text{for } \nu_e \text{ from stellar cooling})$$

$$C=1 \Rightarrow \Lambda_{LNB/W} \gtrsim 10^6 \text{ GeV}$$


# Neutrino mass & magnetic moment

- RG running with  $C_{LNH}(\Lambda) \equiv 0$  [0504134]:

$$\delta m_\nu = \frac{v^2 e}{8m_e c_w} \frac{C_{LNH}(v)}{C_{LNB}(v) + s_w C_{LNW}(v)} \frac{\mu_{\nu N}}{\mu_B}$$

$$\frac{\mu_{\nu N}}{\mu_B} \sim 10^{-15} \frac{\delta m_\nu}{0.1\text{eV}} \Leftrightarrow C_{LNH} \sim 0.02 C_{LNB/W} \quad (\text{taking the same cut-off } \Lambda)$$

$$C = 1 \Rightarrow \Lambda_{LNB/W} \gtrsim 10^7 \text{ GeV}$$



# Neutrino mass from quark condensate

$$\mathcal{O}_{LNQd}(+\text{h.c.}) = (\bar{L}N)\epsilon(\bar{Q}d_R)$$

$$\mathcal{O}_{QuNL}(+\text{h.c.}) = (\bar{Q}u_R)(\bar{N}L)$$

- $\langle \bar{u}u \rangle = \langle \bar{d}d \rangle = \langle \bar{s}s \rangle = \Lambda_{\text{QCD}}^3 \sim (300\text{MeV})^3$
- Dirac neutrino mass:  $m_\nu = \frac{\Lambda_{\text{QCD}}^3}{\Lambda^2} \Rightarrow \Lambda_{LNQd,QuNL} \gtrsim 10\text{TeV}$

# Stellar cooling

- Neutral current processes:  $f \rightarrow fNN(\nu)$  from the  $LNLe$  &  $ffNN$  ( $f = e, u, d, LL, QQ$ ) operators

limiting  $\Lambda \gtrsim 4$  TeV.

Raffelt, 9903472

- Charged current processes:  $eq \rightarrow q'N$  from the  $duNe, LNQd, LdQN, QuNL$  operators

limiting  $\Lambda \gtrsim 70$  TeV.

# Contribution to $\Delta N_{\text{eff}}$

$$\Delta N_{\text{eff}} = \frac{\sum_{\alpha} \rho_{N_R}^{\alpha}}{\rho_{\nu_L}} = 3 \times \frac{\rho_{N_R}}{\rho_{\nu_L}} = 3 \times \left( \frac{T_{N_R}}{T_{\nu_L}} \right)^4$$

- Tracking thermalization of  $N$  down to the BBN era.

$$\dot{\rho}_{N_R} + 3H(\rho_{N_R} + P_{N_R}) = \mathcal{C}_{N_R},$$

$$T \frac{d\xi_{N_R}}{dT} + (1 - \beta) \xi_{N_R} = -\frac{\beta}{4\kappa \xi_{N_R}^3 H T^4} \mathcal{C}_{N_R},$$


$$\xi_{N_R} \equiv \frac{T_{N_R}}{T_{\nu_L}} \quad \kappa = 2 \times \frac{7\pi^2}{8 \cdot 30}$$

$$\beta = \frac{g_{\star}^{1/2}(T) \sqrt{g_{\rho}(T)}}{g_s(T)} \quad g_{\star}^{1/2} = \frac{g_s}{\sqrt{g_{\rho}}} \left( 1 + \frac{1}{3} \frac{T}{g_s} \frac{dg_s}{dT} \right)$$


$$\mathcal{C}_{N_R} = \int \prod_{\alpha=1}^4 d\Pi_{\alpha} (2\pi)^4 \delta^4(\mathbf{p}_1 + \mathbf{p}_2 - \mathbf{p}_3 - \mathbf{p}_4) \times |\overline{\mathcal{M}}|_{N_R+X \rightarrow Y+Z}^2 \left( \exp\left(-\frac{E_3 + E_4}{T}\right) \frac{n_Y}{n_Y^{eq}} \frac{n_Z}{n_Z^{eq}} - \exp\left(-\frac{E_1}{T_{N_R}} - \frac{E_2}{T}\right) \frac{n_{N_R}}{n_{N_R}^{eq}} \frac{n_X}{n_X^{eq}} \right) E_1$$

# Thermalization processes


Involved Operators	Vertices
$\Lambda_{LNB}$	$\frac{v}{\sqrt{2}\Lambda_{LNB}^2} \bar{\nu}_L \sigma^{\mu\nu} N_R B_{\mu\nu}$
$\Lambda_{LNW}$	$\frac{v}{\sqrt{2}\Lambda_{LNW}^2} \bar{\nu}_L \sigma^{\mu\nu} N_R W_{\mu\nu}^3$
$\Lambda_{LNW}$	$\frac{v}{\sqrt{2}\Lambda_{LNW}^2} \bar{e}_L \sigma^{\mu\nu} N_R (W_{\mu\nu}^1 + iW_{\mu\nu}^2)$
$\Lambda_{HNe}$	$-\frac{v^2 g_L}{\sqrt{2}\Lambda_{HNe}^2} \bar{N}_R \gamma^\mu e_R W_\mu^+$
$\Lambda_{HN}$	$\frac{v^2 g_L}{2C_W \Lambda_{HN}^2} \bar{N}_R \gamma^\mu N_R Z_\mu$



# $\Delta N_{\text{eff}}$ limits



Cut-off scale	Constraint from $N_{\text{eff}}$ (TeV)
$\Lambda_{\text{LNB}}$	80
$\Lambda_{\text{LNW}}$	50
$\Lambda_{\text{HNe}}$	15
$\Lambda_{\text{HN}}$	8



$$\sigma(eN \rightarrow ev) \propto \left( 2(2m_A^2 + s) \log \left( \frac{m_A^2 + s}{m_A^2} \right) - 4s \right) \quad m_A^2 = \frac{e^2 T^2}{6}$$

# BNV operators ( $B - L = 0$ )

$$\mathcal{O}_1 = \left( \overline{d_{\alpha R}^c} u_{\beta R} \right) \left( \overline{Q_{i\gamma L}^c} L_{jL} \right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij},$$

$$\mathcal{O}_2 = \left( \overline{Q_{i\alpha L}^c} Q_{j\beta L} \right) \left( \overline{u_{\gamma R}^c} e_R \right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij},$$

$$\mathcal{O}_3 = \left( \overline{Q_{i\alpha L}^c} Q_{j\beta L} \right) \left( \overline{Q_{k\gamma L}^c} L_{lL} \right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij} \epsilon_{kl},$$

$$\mathcal{O}_4 = \left( \overline{d_{\alpha R}^c} u_{\beta R} \right) \left( \overline{u_{\gamma R}^c} e_R \right) \epsilon_{\alpha\beta\gamma}.$$

$$\mathcal{O}_{N1} = \left( \overline{Q_{i\alpha L}^c} Q_{j\beta L} \right) \left( \overline{d_{\gamma R}^c} N_R \right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij},$$

1612.04527

$$\mathcal{O}_{N2} = \left( \overline{u_{\alpha R}^c} d_{\beta R} \right) \left( \overline{d_{\gamma R}^c} N_R \right) \epsilon_{\alpha\beta\gamma}.$$

# Proton decay

- Nucleon transition to meson+lepton induced by BNV operators:

JLQCD, 9911026

$$\bar{v}_\ell^c(\vec{q}, s) \langle P(\vec{p}) | \mathcal{O}^{\chi L} | N(\vec{k}, s) \rangle = \bar{v}_\ell^c(q, s) P_L \left[ W_0^\chi(Q^2) - \frac{i\not{q}}{m_N} W_1^\chi(Q^2) \right] u_N(k, s)$$

$$Q^2 = -q^2 = -(E_N - E_P)^2 + (\vec{k} - \vec{p})^2$$

$$\Gamma(N \rightarrow P\bar{\ell}) = \frac{1}{32\pi m_N} \lambda^{\frac{1}{2}} \left( 1, \frac{m_\ell^2}{m_N^2}, \frac{m_P^2}{m_N^2} \right) \left( (m_\ell^2 + m_N^2 - m_P^2) \left( |\tilde{W}_0|^2 + \frac{m_\ell^2}{m_N^2} |\tilde{W}_1|^2 \right) - 4m_\ell^2 \text{Im} \left( \tilde{W}_0 \tilde{W}_1^* \right) \right)$$

$$\tilde{W}_{0,1} = \sum_I \frac{C^I}{\Lambda^2} W_{0,1}^I(N \rightarrow P).$$

# Form factors [2111.01608]


$$\begin{aligned}\mathcal{O}_1 &= \left(\overline{d_{\alpha R}^c} u_{\beta R}\right) \left(\overline{Q_{i\gamma L}^c} L_{jL}\right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij}, \\ \mathcal{O}_2 &= \left(\overline{Q_{i\alpha L}^c} Q_{j\beta L}\right) \left(\overline{u_{\gamma R}^c} e_R\right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij}, \\ \mathcal{O}_3 &= \left(\overline{Q_{i\alpha L}^c} Q_{j\beta L}\right) \left(\overline{Q_{k\gamma L}^c} L_{lL}\right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij} \epsilon_{kl}, \\ \mathcal{O}_4 &= \left(\overline{d_{\alpha R}^c} u_{\beta R}\right) \left(\overline{u_{\gamma R}^c} e_R\right) \epsilon_{\alpha\beta\gamma}, \\ \mathcal{O}_{N1} &= \left(\overline{Q_{i\alpha L}^c} Q_{j\beta L}\right) \left(\overline{d_{\gamma R}^c} N_R\right) \epsilon_{\alpha\beta\gamma} \epsilon_{ij}, \\ \mathcal{O}_{N2} &= \left(\overline{u_{\alpha R}^c} d_{\beta R}\right) \left(\overline{d_{\gamma R}^c} N_R\right) \epsilon_{\alpha\beta\gamma}.\end{aligned}$$

Matrix element	$W_0$	$W_1$
$\langle \pi^+   (ud)_L d_L   p \rangle$	0.1032 0.105	-0.130 -0.132
$\langle \pi^+   (ud)_R d_L   p \rangle$	-0.1125 -0.1139	0.116 0.118
$\langle K^0   (us)_L u_L   p \rangle$	0.0395 0.0397	0.0256 0.0254
$\langle K^0   (us)_R u_L   p \rangle$	0.0688 0.0693	-0.0250 -0.0254
$\langle K^+   (us)_L d_L   p \rangle$	0.0263 0.0266	-0.0448 -0.0453
$\langle K^+   (us)_R d_L   p \rangle$	-0.0301 -0.0307	0.0452 0.0458
$\langle K^+   (ud)_L s_L   p \rangle$	0.0923 0.0932	-0.0638 -0.0653
$\langle K^+   (ud)_R s_L   p \rangle$	-0.0835 -0.0846	0.0588 0.0605
$\langle K^+   (ds)_L u_L   p \rangle$	-0.0651 -0.0658	0.0192 0.0201
$\langle K^+   (ds)_R u_L   p \rangle$	-0.0394 -0.0393	-0.0203 -0.0204

$$\langle \pi^0 | (ud)_\chi u_L | p \rangle = \sqrt{2} \langle \pi^+ | (ud)_\chi d_L | p \rangle$$

# Nucleon decay limits: $p \rightarrow \pi^+ / K^+ + N$

Process	$\tau$ ( $10^{33}$ years)	operators	$\Lambda_{\min}$ [ $10^{15}$ GeV]
$p \rightarrow \pi^0 e^+$ ( $\pi^0 \mu^+$ )	24 (16) [58]	$C_{1,2,3,4}^{111\ell}$	3.88 (3.52)
$n \rightarrow \pi^- e^+$ ( $\pi^- \mu^+$ )	5.3 (3.5) [59]		2.66 (2.41)
$p \rightarrow \eta^0 e^+$ ( $\eta^0 \mu^+$ )	10 (4.7) [59]	$C_{1,2,3,4}^{111\ell}$	2.56 (2.12)
$n \rightarrow \eta^0 \nu$	0.158 [60]	$C_{1,3}^{111\ell}, C_{N1,N2}^{111i}$	0.9
$p \rightarrow K^+ \nu$	6.61 [61]	$C_{1,3,N1}^{211i}, C_{1,3,N1,N2}^{112i}, C_{3,N1,N2}^{121i}$	3.46
$n \rightarrow K^0 \nu$	0.13 [62]		1.3
$p \rightarrow K^0 e^+$ ( $K^0 \mu^+$ )	1 (1.6) [63, 64]	$C_{1,2,3,4}^{211\ell}, C_{2,3}^{121\ell}$	1.1 (1.23)
$p \rightarrow \pi^+ \nu$	0.39 [62]	$C_{1,3,N1,N2}^{111i}$	1.64
$n \rightarrow \pi^0 \nu$	1.1 [62]		2.14



Cf) Flavor dependent limits, e.g.,  $t^c b^c s^c N$  (?)

# Conclusion

- Study of Dirac neutrino EFT (B-L invariance with RHN  $N$ ).
- RHN appears as MET in collider searches:  $\Lambda_{\text{relevant}} \gtrsim O(\text{TeV})$ .
- Neutrino mass generation:  
 $\Lambda_{LNH} \gtrsim 10^8 \text{ GeV}, \Lambda_{LNB/W} \gtrsim 10^7 \text{ GeV}, \Lambda_{LNQd, QuNL} \gtrsim 10^4 \text{ GeV}$ .
- Stellar cooling:  $\Lambda_{(ffNN, LNLe; duNe, LNQd, LdQN, QuNL)} \gtrsim (4; 70) \text{ TeV}$
- $\Delta N_{\text{eff}}$  limits:  $\Lambda_{HNe, HN} \gtrsim \text{TeV}, \Lambda_{LNB(W)} \gtrsim 8(5) \text{ TeV}$ .
- Proton decays to RHN:  $\Lambda_{N1,2}^{(u,d,s)} \gtrsim 10^{15} \text{ GeV}$  (extendable to general flavor structure).