

Abstract

Introduction

Bound State

QCD sum rules approach for the masses and decay constants of the

Phenomenological part

QCD side

Numerical analysis

Properties of Mesons Containing 4G quark

V. Bashiry

Cyprus International University
Collaborators: K. Azizi, S. Sultansoy

September 23-25, 2011
Istanbul, Turkey



Outline

1

Abstract

2

Introduction

- Fourth Generation what If?

3

Bound State

4

QCD sum rules approach for the masses and decay constants of the

5

Phenomenological part

6

QCD side

- perturbative part
- Non-perturbative part

7

Numerical analysis

we calculate the masses and decay constants of mesons containing one fourth family and the other from observed SM quarks.

Outline

1

Abstract

2

Introduction

- Fourth Generation what If?

3

Bound State

4

QCD sum rules approach for the masses and decay constants of the

5

Phenomenological part

6

QCD side

- perturbative part
- Non-perturbative part

7

Numerical analysis

- Among the mysteries of nature is the number of generations. We observe three generations, however...
- Data from the LHC-CMS (under some circumstances) restricts the masses of the t' and b' quarks in a fourth generation to be greater than about 500 GeV.

- Among the mysteries of nature is the number of generations. We observe three generations, however...
- Data from the LHC-CMS (under some circumstances) restricts the masses of the t' and b' quarks in a fourth generation to be greater than about 500 GeV.

- bound state condition:

$$m_Q < 100|V_{Qq}|^{-2/3} \text{ (see Bigi 1986 PLB) and} \\ 50 < |m_{t'} - m_{b'}| < m_W$$

- $|V_{Qq}| < 0.1$ If it is!!?

- If the fourth generation quarks have a very small mixing with the ordinary quarks, they can be long enough lived that bound $\bar{q}'q'$ states decay through $\bar{q}'q'$ annihilation and not via q' decay to a lower generation quark and a W boson. In this case the production of these bound states at the LHC may have important experimental consequences.

- bound state condition:
 $m_Q < 100|V_{Qq}|^{-2/3}$ (see Bigi 1986 PLB) and
 $50 < |m_{t'} - m_{b'}| < m_W$
- $|V_{Qq}| < 0.1$ If it is!!?
- If the fourth generation quarks have a very small mixing with the ordinary quarks, they can be long enough lived that bound $\bar{q}'q'$ states decay through $\bar{q}'q'$ annihilation and not via q' decay to a lower generation quark and a W boson. In this case the production of these bound states at the LHC may have important experimental consequences.

- bound state condition:
 $m_Q < 100|V_{Qq}|^{-2/3}$ (see Bigi 1986 PLB) and
 $50 < |m_{t'} - m_{b'}| < m_W$
- $|V_{Qq}| < 0.1$ If it is!!?
- If the fourth generation quarks have a very small mixing with the ordinary quarks, they can be long enough lived that bound $\bar{q}'q'$ states decay through $\bar{q}'q'$ annihilation and not via q' decay to a lower generation quark and a W boson. In this case the production of these bound states at the LHC may have important experimental consequences.

- These bound states can be: scalar, pseudoscalar, vector and axial vector type ground states or higher states mesons. The masses and bound states spectrum can be calculated in different methods, for instance, Ishiwata et al PRD 2011.

- QCD Sum Rules Can be applied just for the ground states.
- In one side we have QCD language where everything can be calculated in terms quarks, gluon parameters. In the other side we have hadronic parameters. The result of the QCD calculation is then matched, via dispersion relation, to a sum over hadronic states.

- QCD Sum Rules Can be applied just for the ground states.
- On one side we have QCD language where everything can be calculated in terms of quarks, gluon parameters. On the other side we have hadronic parameters. The result of the QCD calculation is then matched, via dispersion relation, to a sum over hadronic states.

The two point correlation function corresponding to the scalar (S) and pseudoscalar (PS) cases can be written as:

$$\Pi^{S(PS)} = i \int d^4x e^{ip \cdot x} \langle 0 | \mathcal{T} \left(J^{S(PS)}(x) \bar{J}^{S(PS)}(0) \right) | 0 \rangle, \quad (1)$$

where \mathcal{T} is the time ordering product and $J^S(x) = \bar{u}_4(x)q(x)$ and $J^{PS}(x) = \bar{u}_4(x)\gamma_5 q(x)$ are the interpolating currents of the heavy scalar and pseudoscalar bound states, respectively.

Similarly for the vector (V) and axial vector (AV), the correlation function can be written as:

$$\Pi_{\mu\nu}^{V(AV)} = i \int d^4x e^{ip \cdot x} \langle 0 | \mathcal{T} \left(J_{\mu}^{V(AV)}(x) \bar{J}_{\nu}^{V(AV)}(0) \right) | 0 \rangle, \quad (2)$$

where, the currents $J_{\mu}^V = \bar{u}_4(x) \gamma_{\mu} q(x)$ and $J_{\mu}^{AV} = \bar{u}_4(x) \gamma_{\mu} \gamma_5 q(x)$ are responsible for creating the vector and axial vector quarkonia, respectively from the vacuum with the same quantum numbers as the interpolating currents.

- The aforesaid correlation functions must be treated in two alternative ways.
- In physical or phenomenological side, we calculate them in terms of hadronic parameters such as masses and decay constants.
- In QCD or theoretical side, they are calculated in terms of QCD degrees of freedom such as quark masses and gluon condensates by the help of operator product expansion (OPE) in deep Euclidean region.

- The aforesaid correlation functions must be treated in two alternative ways.
- In physical or phenomenological side, we calculate them in terms of hadronic parameters such as masses and decay constants.
- In QCD or theoretical side, they are calculated in terms of QCD degrees of freedom such as quark masses and gluon condensates by the help of operator product expansion (OPE) in deep Euclidean region.

- The aforesaid correlation functions must be treated in two alternative ways.
- In physical or phenomenological side, we calculate them in terms of hadronic parameters such as masses and decay constants.
- In QCD or theoretical side, they are calculated in terms of QCD degrees of freedom such as quark masses and gluon condensates by the help of operator product expansion (OPE) in deep Euclidean region.

- Equating these two representations of the correlation function through dispersion relations, we acquire the QCD sum rules for the masses and decay constants.




Figure: (a): Bare loop diagram (b): Diagrams corresponding to gluon condensates.

- These sum rules relate the hadronic parameters to the fundamental QCD parameters.
- To suppress the contribution of the higher states and continuum, the Borel transformation with respect to the momentum squared is applied to both sides of the correlation functions.

- These sum rules relate the hadronic parameters to the fundamental QCD parameters.
- To suppress the contribution of the higher states and continuum, the Borel transformation with respect to the momentum squared is applied to both sides of the correlation functions.

- Perform the integral over x and isolating the ground state, we obtain

$$\Pi^{S(PS)} = \frac{\langle 0 | J^{S(PS)}(0) | S(PS) \rangle \langle S(PS) | J^{S(PS)}(0) | 0 \rangle}{m_{S(PS)}^2 - p^2} + \dots \quad (3)$$

where \dots represents the contributions of the higher states and continuum and $m_{S(PS)}$ is mass of the heavy scalar(pseudoscalar) meson. From the similar manner, for the vector (axial vector) case, we obtain

$$\Pi_{\mu\nu}^{V(AV)} = \frac{\langle 0 | J_{\mu}^{V(AV)}(0) | V(AV) \rangle \langle V(AV) | J_{\nu}^{V(AV)}(0) | 0 \rangle}{m_{V(AV)}^2 - p^2} + \dots \quad (4)$$

To proceed, we need to know the matrix elements of the interpolating currents between the vacuum and mesonic states. These matrix elements are parametrized in terms of the leptonic decay constants as:

$$\begin{aligned}
 \langle 0 | J(0) | S \rangle &= f_S m_S, \\
 \langle 0 | J(0) | PS \rangle &= f_{PS} \frac{m_{PS}^2}{m_{u4} + m_q}, \\
 \langle 0 | J_\mu(0) | V(AV) \rangle &= f_{V(AV)} m_{V(AV)} \varepsilon_\mu,
 \end{aligned} \tag{5}$$

Using the summation over the polarization vectors in the $V(AV)$

$$\epsilon_\mu \epsilon_\nu^* = -g_{\mu\nu} + \frac{p_\mu p_\nu}{m_{V(AV)}^2}, \quad (6)$$

The physical sides of the correlation functions as:

$$\begin{aligned} \Pi^S &= \frac{f_S^2 m_S^2}{m_S^2 - p^2} + \dots \\ \Pi^{PS} &= \frac{f_{PS}^2 \left(\frac{m_{PS}^2}{m_{u4} + m_q}\right)^2}{m_{PS}^2 - p^2} + \dots \\ \Pi_{\mu\nu}^{V(AV)} &= \frac{f_{V(AV)}^2 m_{V(AV)}^2}{m_{V(AV)}^2 - p^2} \left[-g_{\mu\nu} + \frac{p_\mu p_\nu}{m_{V(AV)}^2} \right] + \dots, \end{aligned}$$

For each correlation function we write

$$\Pi^{\text{QCD}} = \Pi_{\text{pert}} + \Pi_{\text{nonpert}}. \quad (8)$$

The bare loop diagram in figure (1) part (a)). The long distance contributions (diagrams shown in figure (1) part (b)) are parameterized in terms of gluon condensates.

$$\Pi^{\text{QCD}} = \int \frac{ds \rho(s)}{s - p^2} + \Pi_{\text{nonpert}}, \quad (9)$$

where, $\rho(s)$ is called the spectral density.

Outline

1

Abstract

2

Introduction

- Fourth Generation what If?

3

Bound State

4

QCD sum rules approach for the masses and decay constants of the

5

Phenomenological part

6

QCD side

- perturbative part
- Non-perturbative part

7

Numerical analysis

The Feynman amplitude of the bare loop diagram is calculated by the help of the Cutkosky rules, where the quark propagators are replaced by Dirac delta function, i.e.,

$$\frac{1}{p^2 - m^2} \rightarrow (-2\pi i)\delta(p^2 - m^2).$$

$$\rho(s) = \frac{3s}{8\pi^2} \left(1 - \frac{(m_1 \pm m_2)^2}{s}\right) \sqrt{1 - 2\frac{m_1^2 + m_2^2}{s} + \frac{(m_1^2 - m_2^2)^2}{s^2}} \quad (10)$$

where + sign in $(m_1 \pm m_2)$ is chosen for scalar and axial vector cases and - sign is for pseudoscalar and vector channels. Here, $m_1 = m_{u_4}$ and m_2 is either m_{u_4} or $m_{c(b)}$.

Outline

1

Abstract

2

Introduction

- Fourth Generation what If?

3

Bound State

4

QCD sum rules approach for the masses and decay constants of the

5

Phenomenological part

6

QCD side

- perturbative part
- Non-perturbative part

7

Numerical analysis

we calculate the gluon condensate diagrams represented in part (b) of figure (1). The vacuum gluon field is expressed as:

$$A_{\mu}^a(k') = -\frac{i}{2}(2\pi)^4 G_{\rho\mu}^a(0) \frac{\partial}{\partial k'_{\rho}} \delta^{(4)}(k'), \quad (11)$$

where k' is the gluon momentum and the quark-gluon-quark vertex as:

$$\Gamma_{ij\mu}^a = ig\gamma_{\mu} \left(\frac{\lambda^a}{2} \right)_{ij}, \quad (12)$$

After straightforward but lengthy calculations, the non-perturbative part for each case in momentum space is obtained as:

$$\Pi_{nonpert}^i = \int_0^1 \langle \alpha_s G^2 \rangle \frac{\Theta^i + \Theta^i(m_1 \leftrightarrow m_2)}{96\pi(m_2^2 + m_1^2 x - m_2^2 x - p^2 x + p^2 x^2)^4} dx \quad (13)$$

where $\Theta^i(m_1 \leftrightarrow m_2)$ means that in Θ^i , we exchange m_1 and m_2 . The explicit expressions for Θ^i are given as:

$$\begin{aligned}
\Theta^S &= \frac{1}{2}x^2 \left\{ 3m_1^4 x(m_2^2(x(17 - 2x(2x(9x - 26) + 47)) + 8) \right. \\
&+ p^2 x(x(27x - 25) - 7)(x - 1)^2) + 2m_2 m_1^3 (m_2^2(x(x(x(21x - 58) + 39) \\
&+ 12) - 15) - p^2(x - 1)x(x(x(7x - 13) - 3) + 12)) \\
&+ m_1^2(-m_2^2 p^2(x - 1)x(x(x(2x(81x - 242) + 455) - 96) - 33) \\
&+ m_2^4(x(x(x(3x(36x - 145) + 652) - 414) + 72) + 15) + 3p^4(x - 1)^3 \\
&x^2(24x^2 - 22x - 5)) - m_2 m_1(x - 1)(-m_2^2 p^2(x^2 - 2)(x(14x - 27) + 15) \\
&+ m_2^4(3x - 5)(x(7x - 12) + 6) + p^4(x - 1)x(x(2x(7x - 13) + 3) + 12)) \\
&+ (x - 1)(-m_2^2 p^4(x - 1)x(2x(x(2x(18x - 55) + 109) - 30) - 9) \\
&+ m_2^4 p^2(x(x(x(x(81x - 328) + 490) - 299) + 42) + 15) \\
&- m_2^6(2x - 3)(x(6x(3x - 8) + 47) - 15) + 3p^6(x - 1)^3 x^2(6(x - 1)x - 1)) \\
&\left. + 9m_1^6(x - 1)^2 x^2(4x + 1) + 3m_2 m_1^5 x((8 - 7x)x + 2) - 4) \right\},
\end{aligned}$$

(14)



The next step is to match the phenomenological and QCD sides of the correlation functions to get sum rules for the masses and decay constants of the bound states. To suppress the contribution of the higher states and continuum, Borel transformation over p^2 as well as continuum subtraction are performed. As a result of this procedure, we obtain the following sum rules:

$$\begin{aligned}
 m_{S(V)(AV)}^2 f_{S(V)(AV)}^2 e^{-\frac{m_{S(V)(AV)}^2}{M^2}} &= \int_{(m_1+m_2)^2}^{s_0} ds \rho^{S(V)(AV)}(s) e^{-\frac{s}{M^2}} + \hat{B}\Pi_{nonpert}^{S(V)(AV)}, \\
 \frac{m_{PS}^4 f_{PS}^2}{(m_{u_4} + m_q)^2} e^{-\frac{m_{PS}^2}{M^2}} &= \int_{(m_1+m_2)^2}^{s_0} ds \rho^{PS}(s) e^{-\frac{s}{M^2}} + \hat{B}\Pi_{nonpert}^{PS}, \quad (15)
 \end{aligned}$$

where M^2 is the Borel mass parameter and s_0 is the continuum threshold. The sum rules for the masses are obtained applying derivative with respect to $-\frac{1}{M^2}$ to the both sides of the above sum rules and dividing by themselves. i.e.,

$$m_{S(PS)(V)(AV)}^2 = \frac{-\frac{d}{d(\frac{1}{M^2})} \left[\int_{(m_1+m_2)^2}^{s_0} ds \rho^{S(PS)(V)(AV)}(s) e^{-\frac{s}{M^2}} + \hat{B}\Pi_{nonpert}^{S(PS)(V)(AV)} \right]}{\int_{(m_1+m_2)^2}^{s_0} ds \rho^{S(PS)(V)(AV)}(s) e^{-\frac{s}{M^2}} + \hat{B}\Pi_{nonpert}^{S(PS)(V)(AV)}}, \quad (16)$$

where

$$\hat{B}\Pi_{nonpert}^i = \int_0^1 e^{\frac{m_2^2 + x(m_1^2 - m_2^2)}{M^2 x(x-1)}} \frac{\Delta^i + \Delta^i(m_1 \leftrightarrow m_2)}{\pi 96 M^6 (x-1)^4 x^3} \langle \alpha_s G^2 \rangle dx, \quad (17)$$

and

$$\begin{aligned}
\Delta^S &= -m_2 m_1^3 (x-1)x^2(m_2^2(14x^2 - 29x + 14) \\
&+ 2M^2x(7x^2 - 13x + 6)) + m_1^4(x-1)x^3(m_2^2(9x^2 - 14x + 6) \\
&+ 3M^2x(3x^2 - 4x + 1)) + m_2 m_1(x-1)(m_2^2 M^2 x \\
&(14x^4 - 53x^3 + 71x^2 - 36x + 6) + m_2^4(7x^4 - 28x^3 + 40x^2 - 25x + 6) \\
&+ 2M^4 x^2(14x^4 - 40x^3 + 29x^2 + 9x - 12)) + m_1^2 x(m_2^2 M^2 x \\
&(-18x^5 + 70x^4 - 105x^3 + 77x^2 - 27x + 3) + m_2^4(-9x^5 + 37x^4 \\
&- 61x^3 + 52x^2 - 21x + 3) - 12M^4 x^2(3x + 1)(x-1)^4) - (x-1) \\
&(-2m_2^2 M^4 x^3(18x^4 - 76x^3 + 123x^2 - 89x + 24) \\
&+ m_2^4 M^2 x(-9x^5 + 40x^4 - 71x^3 + 68x^2 - 33x + 6) + m_2^6(-3x^5 + 14x^4 \\
&- 27x^3 + 29x^2 - 15x + 3) + 6M^6(x-1)^3 x^3(6x^2 - 6x - 1)) \\
&- 3m_1^6(x-1)x^5 + m_2 m_1^5 x^3(7x^2 - 8x + 1),
\end{aligned}$$

(18)

we take the mass of the u_4 in the interval $m_{u_4} = (450 - 550) \text{ GeV}$, $m_b = 4.8 \text{ GeV}$, $m_c = 1.3 \text{ GeV}$ and $\langle 0 | \frac{1}{\pi} \alpha_s G^2 | 0 \rangle = 0.012 \text{ GeV}^4$. The sum rules for the masses and decay constants also contain two auxiliary parameters, namely Borel mass parameter M^2 and continuum threshold s_0 . The standard criteria in QCD sum rules is that the physical quantities should be independent of the auxiliary parameters. Therefore, we should look for working regions of these parameters such that our results be approximately insensitive to the variation of auxiliary parameters.

The working region for the Borel mass parameter is determined demanding that not only the higher state and continuum contributions are suppressed but also the contributions of the highest order operators should be small, i.e., the sum rules for the masses and decay constants should converge. As a result of the above procedure, the working region for the Borel parameter is found to be $500 \text{ GeV}^2 \leq M^2 \leq 900 \text{ GeV}^2$ for $\bar{u}_4 b$ and $\bar{u}_4 c$, and $1200 \text{ GeV}^2 \leq M^2 \leq 2000 \text{ GeV}^2$ for $\bar{u}_4 u_4$ heavy SM₄ mesons.

The continuum threshold s_0 is not completely arbitrary but it is related to the energy of the first excited states with the same quantum numbers as the interpolating currents. Our numerical calculations show that in the interval $(m_1 + m_2 + 0.3)^2 \text{ GeV}^2 \leq s_0 \leq (m_1 + m_2 + 0.5)^2 \text{ GeV}^2$ for the continuum threshold, our results have very weak dependency on these parameters.

mass (GeV)	$u_4 \bar{c}$	$u_4 \bar{b}$	$u_4 \bar{u}_4$
Scalar	552.82 ± 0.31	556.27 ± 0.31	1101.67 ± 0.60
Pseudoscalar	552.43 ± 0.18	555.78 ± 0.18	1101.11 ± 0.36
Axial Vector	552.81 ± 0.31	556.25 ± 0.31	1101.68 ± 0.60
Vector	552.42 ± 0.18	555.77 ± 0.18	1101.12 ± 0.36

Table: The values of masses of different bound states obtained using $m_{u_4} = 550 \text{ GeV}$.

Leptonic decay constant f (GeV)	$u_4 \bar{c}$	$u_4 \bar{b}$	$u_4 \bar{u}_4$
Scalar	0.10 ± 0.01	0.12 ± 0.01	0.26 ± 0.03
Pseudoscalar	0.14 ± 0.01	0.27 ± 0.01	4.19 ± 0.20
Axial Vector	0.10 ± 0.01	0.12 ± 0.01	0.26 ± 0.03
Vector	0.14 ± 0.01	0.27 ± 0.01	4.18 ± 0.20

Table: The values of decay constants of different bound states obtained using $m_{u_4} = 550 \text{ GeV}$.




Figure: Dependence of mass of the scalar $\bar{u}_4 u_4$ on the Borel parameter, M^2 at three fixed values of the continuum threshold. The upper, middle and lower lines belong to the values $s_0 = (m_1 + m_2 + 0.5)^2 \text{ GeV}^2$, $s_0 = (m_1 + m_2 + 0.4)^2 \text{ GeV}^2$ and $s_0 = (m_1 + m_2 + 0.3)^2 \text{ GeV}^2$, respectively.




Figure: Dependence of the decay constant of the scalar $\bar{u}_4 u_4$ on the Borel parameter, M^2 at three fixed values of the continuum threshold. The upper, middle and lower lines belong to the values $s_0 = (m_1 + m_2 + 3.7)^2 \text{ GeV}^2$, $s_0 = (m_1 + m_2 + 3.5)^2 \text{ GeV}^2$ and $s_0 = (m_1 + m_2 + 3.3)^2 \text{ GeV}^2$, respectively.

To sum up, against the top quark, the heavy fourth generation of quarks that have sufficiently small mixing with the three known family SM quarks form hadrons. Considering the arguments mentioned in the text, the production of such bound states will be possible at LHC. Hoping this possibility, we calculated the masses and decay constants of the bound state objects containing two quarks either both quarks from the SM₄ or one from heavy fourth generation and the other from observed SM bottom or charm quarks in the framework of the QCD sum rules. The obtained numerical results approach to the known masses and decay constants of the $\bar{b}b$ and $\bar{c}c$ heavy quarkonia, when the fourth family quark is replaced by the bottom or charm quark.