



Contribution ID: 44

Type: poster

## The composition of the primary particles at energies $3 \cdot 10^{17}$ - $3 \cdot 10^{19}$ eV

Calculations of signals in surface and underground scintillation detectors of the Yakutsk array from particles of extensive air showers in terms of QGSJET-II and GEISHA 2002 models with the help of the CORSIKA 6.616 and GEANT4 codes with parameter  $\epsilon=10^{-8}$  of the thinning procedures have been carried out to estimate muon fraction at 600 m from the shower axis at energies  $3 \cdot 10^{17}$  -  $3 \cdot 10^{19}$  eV. Comparison of results of these calculations with the Yakutsk data shows rather heavy composition of the primary radiation in this energy region. But it was shown that calculations in terms of the QGSJET-II model at energies above 100 GeV underestimate the number of muons by a factor of  $\sim 1.5$ . Besides, it was also shown that calculations in terms of the GEISHA 2002 model at energies below 100 GeV underestimate the number of muons by a factor of  $\sim 1.1$  comparing with the FLUCA results. To interpret data one has also to take into account results observed at the Telescope array which show 1.27 difference in calculated in terms of models and observed with help of fluorescent light energy estimates of extensive air showers. Taken together all these corrections show the proton composition of the primary radiation in the energy region  $10^{18}$  -  $3 \cdot 10^{19}$  eV. At energies in the interval of  $3 \cdot 10^{17}$  -  $10^{18}$  eV composition is probably heavier. It is not excluded that at energies above  $3 \cdot 10^{19}$  eV composition may be also more heavier as illustrated by the trend of data.

**Author:** Dr KNURENKO, Stanislav (Shafer Institute for cosmophysical research)

**Presenter:** Dr KNURENKO, Stanislav (Shafer Institute for cosmophysical research)