



Contribution ID: 191

Type: not specified

Measurement of $D^{*\pm}$ Meson Production and Determination of $F_2^{c\bar{c}}$ at low Q^2 in Deep-Inelastic Scattering at HERA

Wednesday, 28 March 2012 11:06 (20 minutes)

Inclusive production of D mesons in deep-inelastic ep scattering at HERA is studied in the range $5 < Q^2 < 100 \text{ GeV}^2$ of the photon virtuality and $0.02 < y < 0.7$ of the inelasticity of the scattering process. The observed phase space for the D meson is $p_T(D) > 1.25 \text{ GeV}$ and $|\eta(D)| < 1.8$. The data sample corresponds to an integrated luminosity of 348 pb^{-1} collected with the H1 detector. Single and double differential cross sections are measured and the charm contribution $F_2^{c\bar{c}}$ to the proton structure function F_2 is determined. The results are compared to perturbative QCD predictions at next-to-leading order implementing different schemes for the charm mass treatment and with Monte Carlo models based on leading order matrix elements with parton showers.

Primary author: DAUM, Karin (University of Wuppertal/DESY)

Presenter: HENNEKEMPER, Eva (Uni Heidelberg)

Session Classification: Combined: Heavy flavours/structure functions

Track Classification: Heavy flavours/structure functions