

Conformal symmetry and the relations between perturbative  
contributions to the Bjorken and Ellis-Jaffe sum rule of the  
polarized DIS

A. L. Kataev

INR, Moscow, Russia

29 March 2012

## Plan of Presentation

### Introduction – Reminding of the concept of Conformal Symmetry

Existing perturbative Quenched  $U(1)$  identity  $O(E^3)$  result: Coefficient functions of Ellis-Jaffe and Bjorken sum rules consider! **Q1** Accident or Not ?

#### A1

- Not! **Statement: true in all orders of PT**

Explanation due to Kataev- 2010 derivation of new Crewther-type relation from triangle diagram of **singlet** A- V-V currents for the Green function of AA -currents and EJ **SI** structure function and comparing with basic **non-singlet** Crewther relation for A-V-V triangle **Crewther 1972**

Positive conclusion of discussion between **Kataev 1996** and **Crewther 1997**

#### A2

The conformal-invariant result for **DIS** sum rules at  $O(A_s^3)$ -level the case of  $SU(N_c)$  **Kataev and Mikhailov 2010, Teor.Mat.Fiz. 2012**

## Conclusions

- 1) Possible applications- tests of future higher-order ' EJ SR analytical calculations
- 2) Possible relations between **polarized parton distributions** in the CI limit.

## Conformal Invariance

is valid in the quark-parton model limit, and perturbative quenched QED. It is the symmetry under the following transformations of coordinates :

1. Translations  $x'^{\mu} = x^{\mu} + \alpha^{\mu}$  with 4 parameters  $\alpha^{\mu}$ ,
2. Scale (or dilaton) transformation  $x'^{\mu} = \rho x^{\mu}$  with 1 parameter  $\rho > 0$ ,
3. Special conformal transformations  $x'^{\mu} = \frac{x^{\mu} + \beta^{\mu} x^2}{1 + 2\beta x + \beta^2 x^2}$  with 4 parameters  $\beta^{\mu}$  and
4. Homogeneous Lorentz transformations  $x'^{\mu} = \Lambda_{\nu}^{\mu} x^{\nu}$ , which also contain 4 parameters.
5. Consequences are widely studied at present, though in renormalized QFT models the CI is violated- appearance of  $\beta$ -function and the effects of running of the coupling constants- QCD, QED

Bjorken polarized sum rule:

$$B_{jp}(Q^2) = \int_0^1 (g_1^{lp}(x, Q^2) - g_1^{ln}(x, Q^2)) dx = \frac{1}{6} g_a C_{B_{jp}}^{ns}(A_s) \quad (1)$$

Depends from  $Q^2$  through the running of  $A_s(Q^2) = \alpha_s(Q^2)/(4\pi)$ ,  
 Consider pQED limit-  $C_F = 1$   $N_F = 0$ ,  $A = \alpha/(4\pi)$

$$C_{B_{jp}}^{ns} = 1 - 3A + \frac{21}{2}A^2 - \frac{3}{2}A^3 - \left( \frac{4823}{8} + 96\zeta(3) \right) A^4 + O(A^5) \quad (2)$$

$O(A^3)$  Larin and Vermaseren (1991);  $O(A^4)$ - Baikov, Chetyrkin, Kuhn (2010)

Ellis-Jaffe sum rule :

$$EJ^{p(n)}(Q^2) = \int_0^1 g_1^{lp(n)}(x, Q^2) dx = C_{B_{jp}}^{ns}(A_s(Q^2)) \left( \pm \frac{1}{12} a_3 + \frac{1}{36} a_8 \right) + C_{EJ_p}^s(Q^2) \frac{1}{9} \Delta\Sigma(Q^2) \quad (3)$$

where  $a_3 = \Delta u - \Delta d$ ,  $a_8 = \Delta u + \Delta d - 2\Delta s$ ,  $\Delta u$ ,  $\Delta d$  and  $\Delta s$  are the polarized distributions and  $\Delta\Sigma$  depends from the scheme choice. In the  $\overline{MS}$ -scheme it is defined as  $\Delta\Sigma = \Delta u + \Delta d + \Delta s$ .

Note, that in the  $\overline{MS}$ -scheme the definition of the singlet coefficient function reads (Larin (1994), Larin, van Ritbergen, Vermaseren (1997)).

$$C_{EJp}^s = \overline{C}_{EJp}^s / Z_5^s \quad . \quad (4)$$

In the perturbative quenched QED the result is

$$\overline{C}_{EJ}^s = 1 - 7A + \frac{89}{2}A^2 - \left( \frac{1397}{6} - 96\zeta(3) \right) A^3 + O(A^4) \quad . \quad (5)$$

$Z_5^s$  is the finite renormalization constants of  $\overline{\Psi}\gamma_\mu\gamma_5\Psi$  current,  $Z_5^{ns}$  is the renormalization constant of  $\overline{\Psi}\gamma_5(\lambda^a/2)\Psi$ -current . They were evaluated in Larin (1994) and Larin and Vermaseren (1991). In the limit of pQED we have

$$Z_5^s(pqQED) = Z_5^{ns}(pqQED) = 1 - 4A + 22A^2 + \left( -\frac{370}{3} + 96\zeta_3 \right) A^3 + O(A^4) \quad . \quad (6)$$

Nontrivial **s**cheme-independent pQED consequence of QCD results for SI and NS coefficient functions of EJ sum rule Kataev (2010):

$$C_{EJp}^s(A) = 1 - 3A + \frac{21}{2}A^2 - \frac{3}{2}A^3 + O(A^4) = C_{Bjp}^{ns}(A) \quad (7)$$

- Q1: What is the theoretical explanation ? Q2: Is this result true in all orders of PT ?  
 Q3 : Is there any  $Q^2$ -dependence ?

A1: Follow from Crewther-type relations for AVV diagrams in CI limit

A2: Valid in all orders of PT A3: No  $Q^2$  dependence- fixed coupling constant

Proof of A1: Using OPE for

$$T_{\mu\alpha\beta}^{ab}(p, q) = i \int \langle 0 | TA_{\mu}(y) V_{\alpha}^a(x) V_{\beta}^b(0) | 0 \rangle e^{ipx+iqy} dx dy = \delta^{ab} \Delta_{\mu\alpha\beta}^{(1-loop)}(p, q) \quad (8)$$

where  $A_{\mu} = \bar{\psi} \gamma_{\mu} \gamma_5 \psi$  and

$$T_{\mu\alpha\beta}^{abc}(p, q) = i \int \langle 0 | TA_{\mu}^a(y) V_{\alpha}^b(x) V_{\beta}^c(0) | 0 \rangle e^{ipx+iqy} dx dy = d^{abc} \Delta_{\mu\alpha\beta}^{(1-loop)}(p, q) \quad (9)$$

The consideration of the first and second triangle graphs+ the concept of conformal invariance give the following relations: [Kataev 1996](#) and [Crewther 1972](#)

$$C_{EJp}^s(A) \times C_D^A(A) = 1 \quad (1996) \quad C_{Bjp}^{ns}(A) \times C_D^{ns}(A) = 1 \quad (1972) \quad (10)$$

where  $C_D^{A(ns)}$  are defined from taking  $q^2 \frac{d}{dq^2}$  of

$$i \int \langle 0 | TA_{\mu}(x) A_{\nu}(0) | 0 \rangle e^{iqx} dx = \Pi_{\mu\nu}^{ax}(q^2) \quad (11)$$

$$i \int \langle 0 | TA_{\mu}^a(x) A_{\nu}^b(0) | 0 \rangle e^{iqx} dx = \delta^{ab} \Pi_{\mu\nu}^{ns}(q^2) . \quad (12)$$

Since in the massless limit chiral symmetry is exact, one has:

$$C_D^A(A) = C_D^{ns}(A) \quad (13)$$

Thus in the CI limit in all orders of PT

$$C_{EJp}^s(A) = C_{Bjp}^{ns}(A) \quad \text{Kataev 2010} \quad (14)$$

In pQED with  $A = \alpha/(4\pi)$  CI limit we have

$$C_{EJp}^s(A) = 1 - 3A + \frac{21}{2}A^2 - \frac{3}{2}A^3 = C_{Bjp}^{ns}(A) \quad (15)$$

In the case of the  $SU(N_c)$  with  $A_s = \alpha_s/(4\pi)$  the similar relation is:

$$\begin{aligned} C_{Bjp}^{ns} &= 1 - 3C_F A_s + \left(\frac{21}{2}C_F^2 - C_F C_A\right) A_s^2 \\ &+ \left[\left(-\frac{3}{2}C_F^3 - 65C_F^2 C_A - \left(\frac{523}{12} - 216\zeta_3\right)C_F C_A^2\right) A_s^3 + O(A_s^4)\right] = C_{EJ}^s(A_s) \end{aligned} \quad (16)$$

Numbers obtained from Crewther relation in Kataev and Mikhailov (2010-2012)

This is the CI predictions for  $O(A_s^4)$ -term in  $C_{EJp}^s(A_s)$

## Conclusion

In the CI limit in  $SU(N_c)$

1. It is possible to try to get  $O(A_s^4)$  for  $C_{Bjp}$ - from [Kataev, Mikhailov \(2012\)](#)- strong test for analytical evaluation of Ellis-Jaffe sum rule at  $A_s^4$  (hope that  $SU(N_c)$  results will be obtained soon).

2. In this CI limit we get the following constraints on polarized PDFs:

$$\frac{EJ^{p(n)}}{Bjp} = \pm \frac{1}{2} + \frac{a_8}{6 a_3} + \frac{2\Delta\Sigma}{3 a_3} \quad (17)$$

where  $a_3 = \Delta u - \Delta d$ ,  $a_8 = \Delta u + \Delta d - 2\Delta s$ , and  $\Delta\Sigma = \Delta u + \Delta d + \Delta s$ .

**Question:** Is it possible to use this relation in practice?