## XX International Workshop on Deep-Inelastic Scattering and Related Subjects



Contribution ID: 64

Type: not specified

## **Fragmentation functions at Belle**

Wednesday 28 March 2012 09:30 (25 minutes)

Fragmentation functions (FFs) describe the formation of final state particles from a partonic initial state. Precise knowledge of these functions is a key ingredient in accessing quantities such as the nucleon spin structure in semi-inclusive deep inelastic scattering and proton proton collisions. However, fragmentation functions can currently not be determined from first principles Quantum Chromodynamics and have to be extracted from experimental data. The Belle experiment at KEK, Japan, provides a large data sample for high precision measurements of quantities allowing for first-time or more precise extractions of fragmentation functions. Completed and ongoing analyses for extractions of spin-independent (unpolarized FFs) as well as spin-dependent fragmentation functions (Collins and Interference FFs) at Belle will be presented.

Primary author: Mr LEITGAB, Martin (University of Illinois at Urbana-Champaign)

**Co-authors:** Mr OGAWA, Akio (BNL); Mr VOSSEN, Anselm (Indiana University Bloomington); Mrs HULSE, Charlotte (The University of the Basque Country); Mrs GIORDANO, Francesca (University of Illinois at Urbana-Champaign); Mr SCHNELL, Gunar (The University of the Basque Country); Mr GROSSE PERDEKAMP, Matthias (University of Illinois at Urbana-Champaign); Mr KOBAYASHI, Noriaki (Tokyo Institute of Technology); Mr SEIDL, Ralf (RIKEN); Mr SHIBATA, Toshi-Aki (Tokyo Institute of Technology); Mr MIYACHI, Yoshiyuki (Tokyo Institute of Technology)

Presenter: Mr LEITGAB, Martin (University of Illinois at Urbana-Champaign)

Session Classification: Spin physics

Track Classification: Spin physics