

Recent Results on Three Loop Corrections to Heavy Quark Contributions to DIS Structure Functions

Johannes Blümlein, DESY


in collaboration : with J. Ablinger (JKU), A. De Freitas (DESY), A. Hasselhuhn (DESY),
C. Schneider (JKU), F. Wißbrock (DESY)



- Introduction
- 3-Loop Gluonic OMEs $O(n_f T_F^2 C_{F,A})$ and FVNS
- 3-Loop Graphs with m_c and m_b
- A General Method to Compute Non-Singular 3-Loop Massive Graphs
- Conclusions

Introduction

Unpolarized Deep-Inelastic Scattering (DIS):



$$\begin{aligned}
 W_{\mu\nu}(q, P, s) &= \frac{1}{4\pi} \int d^4\xi \exp(iq\xi) \langle P, s | [J_\mu^{em}(\xi), J_\nu^{em}(0)] | P, s \rangle \\
 &= \frac{1}{2x} \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) F_L(x, Q^2) + \frac{2x}{Q^2} \left(P_\mu P_\nu + \frac{q_\mu P_\nu + q_\nu P_\mu}{2x} - \frac{Q^2}{4x^2} g_{\mu\nu} \right) F_2(x, Q^2) .
 \end{aligned}$$

Structure Functions: $F_{2,L}$
 contain light and heavy quark contributions.

Factorization of the Structure Functions

At leading twist the structure functions factorize in terms of a Mellin convolution

$$F_{(2,L)}(x, Q^2) = \sum_j \underbrace{\mathbb{C}_{j,(2,L)}\left(x, \frac{Q^2}{\mu^2}, \frac{m^2}{\mu^2}\right)}_{perturbative} \otimes \underbrace{f_j(x, \mu^2)}_{nonpert.}$$

into (pert.) Wilson coefficients and (nonpert.) parton distribution functions (PDFs).

\otimes denotes the Mellin convolution

$$f(x) \otimes g(x) \equiv \int_0^1 dy \int_0^1 dz \delta(x - yz) f(y) g(z) .$$

The subsequent calculations are performed in Mellin space, where \otimes reduces to a multiplication, due to the Mellin transformation

$$\hat{f}(N) := \int_0^1 dx x^{N-1} f(x) .$$

Wilson coefficients:

$$\mathbb{C}_{j,(2,L)} \left(N, \frac{Q^2}{\mu^2}, \frac{m^2}{\mu^2} \right) = \textcolor{blue}{C}_{j,(2,L)} \left(N, \frac{Q^2}{\mu^2} \right) + H_{j,(2,L)} \left(N, \frac{Q^2}{\mu^2}, \frac{m^2}{\mu^2} \right) .$$

At $Q^2 \gg m^2$ the heavy flavor part

$$H_{j,(2,L)} \left(N, \frac{Q^2}{\mu^2}, \frac{m^2}{\mu^2} \right) = \sum_i \textcolor{blue}{C}_{i,(2,L)} \left(N, \frac{Q^2}{\mu^2} \right) \textcolor{red}{A}_{ij} \left(\frac{m^2}{\mu^2}, N \right)$$

[Buza, Matiounine, Smith, van Neerven 1996 Nucl.Phys.B]

factorizes into the light flavor Wilson coefficients C and the massive operator matrix elements (OMEs) of local operators O_i between partonic states j

$$\textcolor{red}{A}_{ij} \left(\frac{m^2}{\mu^2}, N \right) = \langle j | O_i | j \rangle .$$

→ additional Feynman rules with local operator insertions for partonic matrix elements.

The unpolarized light flavor Wilson coefficients are known up to NNLO

[Moch, Vermaseren, Vogt, 2005 Nucl.Phys.B].

For $F_2(x, Q^2)$: at $Q^2 \gtrsim 10m^2$ the asymptotic representation holds at the 1% level.

Status of OME calculations

Leading Order: [Witten, 1976 Nucl.Phys.B; Babcock, Sivers, 1978 Phys.Rev.D; Shifman, Vainshtein, Zakharov, 1978 Nucl.Phys.B; Leveille, Weiler, 1979 Nucl.Phys.B; Glück, Reya, 1979 Phys.Lett.B; Glück, Hoffmann, Reya, 1982 Z.Phys.C.]

Next-to-Leading Order : [Laenen, van Neerven, Riemersma, Smith, 1993 Nucl. Phys. B]

[Large Q^2/m^2 : Buza, Matiounine, Smith, Migneron, van Neerven, 1996 Nucl.Phys.B] IBP

[Bierenbaum, Blümlein, Klein, 2007 Nucl.Phys.B] via pF_q 's, more compact results

[Bierenbaum, Blümlein, Klein 2008 Nucl.Phys.B, 2009 Phys.Lett.B]: $O(\alpha_s^2 \varepsilon)$ contributions (all N)

NNLO: [Bierenbaum, Blümlein, Klein 2009 Nucl.Phys.B] Moments for F_2 : $N = 2\dots 10(14)$

[Blümlein, Klein, Tödtli 2009 Phys. Rev. D] contrib. to transversity: $N = 1\dots 13$

[Ablinger, Blümlein, Klein, Schneider, Wißbrock 2011 Nucl.Phys.B] contrib. $\propto n_f$ to F_2 (all N):

At 3-loop order known:

- $A_{qq,Q}^{\text{PS}}, A_{qg,Q}$: complete
- $A_{Qg}, A_{Qq}^{\text{PS}}, A_{qq,Q}^{\text{NS}} A_{qq,Q}^{\text{NS,TR}}$: all terms of $O(n_f T_F^2 C_{A/F})$
- $A_{Qq}^{\text{PS}}, A_{qq,Q}^{\text{NS}} A_{qq,Q}^{\text{NS,TR}}$: all terms of $O(T_F^2 C_{A/F})$
- $A_{gq,Q}, A_{gg,Q}$: see [this talk](#) \longrightarrow all terms of $O(n_f T_F^2 C_{A/F})$

VFNS Relations for PDFs

The matching conditions for the VFNS:

[Buza, Matiounine, Smith, van Neerven 1998 Eur.Phys.J.C] → NLO

[Bierenbaum, Blümlein, Klein 2009 Nucl.Phys.B] → NNLO


$$\begin{aligned}
 & f_k(N, n_f + 1, \mu^2, m^2) + f_{\bar{k}}(N, n_f + 1, \mu^2, m^2) \\
 &= A_{qq,Q}^{\text{NS}} \left(N, n_f, \frac{\mu^2}{m^2} \right) \otimes [f_k(N, n_f, \mu^2, m^2) + f_{\bar{k}}(N, n_f, \mu^2, m^2)] \\
 &+ \frac{1}{n_f} A_{qq,Q}^{\text{PS}} \left(N, n_f, \frac{\mu^2}{m^2} \right) \otimes \Sigma(N, n_f, \mu^2, m^2, x) + \frac{1}{n_f} A_{qg,Q} \left(N, n_f, \frac{\mu^2}{m^2} \right) \otimes G(N, n_f, \mu^2, m^2, x) \\
 & f_Q(N, n_f + 1, \mu^2, m^2) + f_{\bar{Q}}(N, n_f + 1, \mu^2, m^2) \\
 &= A_{Qq}^{\text{PS}} \left(N, n_f, \frac{\mu^2}{m^2} \right) \otimes \Sigma(N, n_f, \mu^2, m^2) + A_{Qg} \left(N, n_f, \frac{\mu^2}{m^2} \right) \otimes G(N, n_f, \mu^2, m^2) \\
 & G(N, n_f + 1, \mu^2, m^2) \\
 &= A_{gq,Q} \left(N, n_f, \frac{\mu^2}{m^2} \right) \otimes \Sigma(N, n_f, \mu^2, m^2) + A_{gg,Q} \left(N, n_f, \frac{\mu^2}{m^2} \right) \otimes G(N, n_f, \mu^2, m^2)
 \end{aligned}$$

where: $\left(\Sigma(N, n_f, \dots) = \sum_{k=1}^{n_f} (f_k + f_{\bar{k}}), \quad n_f = 3 \right)$

The ABKM09 heavy flavor PDFs: (grey band) compared to MSTW2008 NNLO (dashed line):

[Alekhin, Blümlein, Klein, Moch 2010 Phys.Rev.D]

[Martin, Stirling, Thorne, Watt 2010 Eur.Phys.J.C]



Influence on W^+ production at the LHC (e.g.
 $\sqrt{s} = 7 \text{ TeV}$):

$\text{ABM11}(n_f = 5): \sigma_{W^+} = 59.53^{+0.38 +0.88}_{-0.23 -0.88}$

$\text{ABM11}(n_f = 4): \sigma_{W^+} = 59.08^{+0.30 +0.87}_{-0.14 -0.87}$


[Alekhin, Blümlein, Moch 2012]

BMSN-scheme for the heavy quark part F_2^h in F_2 :

[Buza, Matiounine, Smith, van Neerven 1998 Eur.Phys.J.C], [Alekhin, Blümlein, Klein, Moch 2010 Phys.Rev.D]

$$F_2^{h,\text{BMSN}}(n_f + 1, x, Q^2) = F_2^{h,\text{exact}}(n_f, x, Q^2) + F_2^{h,\text{ZMVFN}}(n_f + 1, x, Q^2) - F_2^{h,\text{asymp}}(n_f, x, Q^2)$$

- $F_2^{h,\text{exact}}$ = exact heavy quark contributions to F_2 in the n_f -flavor scheme
- $F_2^{h,\text{ZMVFN}}$ = similar to a $(n_f + 1)$ -flavor scheme with PDFs defined from the matching
- $F_2^{h,\text{asymp}} \approx F_2^{h,\text{exact}}$ for $Q^2 \gg m_h^2$ without power corrections



The $O(n_f T_F^2 \alpha_s^3)$ contributions to $A_{gg,Q}$

calculation of the 1PI part of $A_{gg,Q}^{(3),n_f T_F^2}$

- generation of Diagrams with **QGRAF** [Nogueira 1993 J. Comput. Phys] → 76 Diagrams
- momentum integrals (regularized in $4 + \varepsilon$ dimensions) → Feynman parameterization → finite sums and hypergeometric functions
- All- ε representation: maximum nestedness 2, hypergeometric functions ${}_3F_2$

Moments were tested using earlier calculations based on **MATAD** by [M. Steinhauser, 2000 CPC].

Then the package **Sigma** [C. Schneider, 2005–] is used for:

- reducing the sums to a small number of key sums
- expanding the summands in ε
- simplifying by symbolic summation algorithms based on $\Pi\Sigma$ -fields [Karr 1981 J. ACM, Schneider 2005–]
- harmonic sums are algebraically reduced using the package **HarmonicSums** (Ablinger) [Ablinger, Blümlein, Schneider 2011]

→ single harmonic sums and ζ -values of max. weight 3 $S_i \equiv \sum_{j=1}^N \frac{1}{j^i}$

$$\begin{aligned}
A_{ggQ}^{n_f T_f^2 \text{1PI}} = & S_\varepsilon^3 a_s^3 n_f T_F^2 \frac{1 + (-1)^N}{2} \left(\frac{m^2}{\mu^2} \right)^{\frac{3}{2}\varepsilon} \left\{ \frac{1}{\varepsilon^3} \left(\textcolor{red}{C_A} \left[\frac{512}{27} S_1 - \frac{64 (3N^4 + 6N^3 + 13N^2 + 10N + 16)}{27(N-1)N(N+1)(N+2)} \right] \right. \right. \\
& - \textcolor{red}{C_F} \frac{512 (N^2 + N + 2)^2}{9(N-1)N^2(N+1)^2(N+2)} \Big) + \frac{1}{\varepsilon^2} \left(\textcolor{red}{C_A} \left[\frac{1280}{81} S_1 - \frac{16P_1}{81(N-1)N^2(N+1)^2(N+2)} \right] \right. \\
& + \textcolor{red}{C_F} \frac{1}{(N-1)(N+2)} \left[\frac{128 (N^2 + N + 2)^2}{9N^2(N+1)^2} S_1 - \frac{128P_2}{27N^3(N+1)^3} \right] \Big) \\
& + \frac{1}{\varepsilon} \left(\textcolor{red}{C_A} \frac{1}{(N-1)(N+2)} \left[-\frac{4P_8}{81N^3(N+1)^3} - \frac{8 (3N^4 + 6N^3 + 13N^2 + 10N + 16)}{9N(N+1)} \zeta_2 \right. \right. \\
& + \frac{32P_9}{27N^2(N+1)^2} S_1 + \frac{64}{9} (N-1)(N+2) \zeta_2 S_1 \Big] + \textcolor{red}{C_F} \frac{1}{(N-1)(N+2)} \left[-\frac{160 (N^2 + N + 2)^2}{9N^2(N+1)^2} S_1^2 \right. \\
& - \frac{64 (N^2 + N + 2)^2}{3N^2(N+1)^2} \zeta_2 + \frac{32 (N^2 + N + 2)^2}{3N^2(N+1)^2} S_2 - \frac{64P_{10}}{81N^4(N+1)^4} + \frac{64P_{11}}{27N^3(N+1)^3} S_1 \Big] \Big) \\
& + \textcolor{red}{C_A} \frac{1}{(N-1)(N+2)} \left[\frac{4P_3}{27N^2(N+1)^2} S_1^2 + \frac{8P_4}{729N^3(N+1)^3} S_1 + \frac{160}{27} (N-1)(N+2) \zeta_2 S_1 \right. \\
& - \frac{448}{27} (N-1)(N+2) \zeta_3 S_1 \Big] + \frac{P_5}{729N^4(N+1)^4} - \frac{2P_6}{27N^2(N+1)^2} \zeta_2 - \frac{4P_7}{27N^2(N+1)^2} S_2 \\
& + \frac{56 (3N^4 + 6N^3 + 13N^2 + 10N + 16)}{27N(N+1)} \zeta_3 \Big] + \textcolor{red}{C_F} \frac{1}{(N-1)(N+2)} \left[\frac{112 (N^2 + N + 2)^2}{27N^2(N+1)^2} S_1^3 \right. \\
& - \frac{16P_{12}}{27N^3(N+1)^3} S_1^2 + \frac{32P_{13}}{81N^4(N+1)^4} S_1 + \frac{16 (N^2 + N + 2)^2}{3N^2(N+1)^2} \zeta_2 S_1 + \frac{16 (N^2 + N + 2)^2}{3N^2(N+1)^2} S_2 S_1 \\
& - \frac{32P_{14}}{243N^5(N+1)^5} - \frac{16P_2}{9N^3(N+1)^3} \zeta_2 + \frac{448 (N^2 + N + 2)^2}{9N^2(N+1)^2} \zeta_3 + \frac{16P_{15}}{9N^3(N+1)^3} S_2 - \frac{160 (N^2 + N + 2)^2}{27N^2(N+1)^2} \zeta_3 \Big] \Big\}
\end{aligned}$$

Renormalization of the OME:

[Bierenbaum, Blümlein, Klein 2009 Nucl.Phys.B]

1. include contributions from **reducible** diagrams
2. perform on-shell **mass** renormalization
3. renormalize the **coupling in a MOM-scheme**, using the background field method
4. remove remaining UV singularities defining **operator Z-factors**
5. remove **collinear singularities** via coll. factorization
6. transform **coupling constant to $\overline{\text{MS}}$**
7. choice: m on-shell or $m_{\overline{\text{MS}}}$

Combining the 1PI part with the gluon self energy contribution:

$$\hat{\Pi}_{\mu\nu}^{ab}(p^2, \hat{m}^2, \mu^2, \hat{a}_s^2) = i\delta^{ab} [-g_{\mu\nu}p^2 + p_\mu p_\nu] \sum_{k=1}^{\infty} \hat{a}_s^k \hat{\Pi}^{(k)}(p^2, \hat{m}^2, \mu^2)$$

$$\hat{\Pi}^{(k)} \equiv \hat{\Pi}^{(k)}(0, \hat{m}^2, \mu^2)$$

giving:

$$\begin{aligned} \hat{A}_{gg,Q}^{(3)} &= \hat{A}_{gg,Q}^{(3),1\text{PI}} - \hat{\Pi}^{(3)} - \hat{A}_{gg,Q}^{(2),1\text{PI}} \hat{\Pi}^{(1)} - 2\hat{A}_{gg,Q}^{(1)} \hat{\Pi}^{(2)} + \hat{A}_{gg,Q}^{(1)} \hat{\Pi}^{(1)} \hat{\Pi}^{(1)} \\ &\equiv \frac{a_{gg,Q}^{(3,0)}}{\varepsilon^3} + \frac{a_{gg,Q}^{(3,1)}}{\varepsilon^2} + \frac{a_{gg,Q}^{(3,2)}}{\varepsilon} + a_{gg,Q}^{(3)} \end{aligned}$$

The renormalization determines the **coefficients of the ε -poles** in terms of coefficients from the **mass renormalization**: $\delta m_1^{(-1)}, \delta m_1^{(0)}, \delta m_1^{(1)}, \delta m_2^{(-2)}, \delta m_2^{(-1)}, \delta m_2^{(0)}$ contributions β_i to the **$\overline{\text{MS}}$ - β -function** and

$\beta_{i,Q}$ to the **MOM-beta-function** in QCD,

splitting functions: $\gamma_{ij} \equiv \sum_{l=1}^{\infty} a_s^{\overline{\text{MS}}l} \gamma_{ij}^{(l)}$ with $\hat{\gamma}_{ij}^{(k)} \equiv \gamma_{ij}^{(k)}(n_f + 1) - \gamma_{ij}^{(k)}(n_f)$ and

lower order contributions to the unrenormalized OMEs:

$$\hat{\hat{A}}_{gg,Q}^{(2)} \equiv \frac{a_{gg,Q}^{(2,0)}}{\varepsilon^2} + \frac{a_{gg,Q}^{(2,1)}}{\varepsilon} + a_{gg,Q}^{(2)} + \varepsilon \bar{a}_{gg,Q}^{(2)} + O(\varepsilon^2)$$

$$\hat{\hat{A}}_{gq,Q}^{(2)} \equiv \frac{a_{gq,Q}^{(2,0)}}{\varepsilon^2} + \frac{a_{gq,Q}^{(2,1)}}{\varepsilon} + a_{gq,Q}^{(2)} + \varepsilon \bar{a}_{gq,Q}^{(2)} + O(\varepsilon^2)$$

$$\hat{\hat{A}}_{Qg,Q}^{(2)} \equiv \frac{a_{Qg}^{(2,0)}}{\varepsilon^2} + \frac{a_{Qg}^{(2,1)}}{\varepsilon} + a_{Qg}^{(2)} + \varepsilon \bar{a}_{Qg}^{(2)} + O(\varepsilon^2)$$

[Bierenbaum, Blümlein, Klein 2007,2008 Nucl.Phys.B, 2009 Phys.Lett.B]

The structure of the unrenormalized OME: [Bierenbaum, Blümlein, Klein 2009 Nucl.Phys.B]

$$\begin{aligned}
\hat{A}_{gg,Q}^{(3)} = & \left(\frac{\hat{m}^2}{\mu^2} \right)^{3\varepsilon/2} \left[\frac{1}{\varepsilon^3} \left(-\frac{\gamma_{gq}^{(0)} \hat{\gamma}_{qg}^{(0)}}{6} \left[\gamma_{gg}^{(0)} - \gamma_{qq}^{(0)} + 6\beta_0 + 4n_f \beta_{0,Q} + 10\beta_{0,Q} \right] - \frac{2\gamma_{gg}^{(0)} \beta_{0,Q}}{3} \left[2\beta_0 + 7\beta_{0,Q} \right] \right. \right. \\
& - \frac{4\beta_{0,Q}}{3} \left[2\beta_0^2 + 7\beta_{0,Q}\beta_0 + 6\beta_{0,Q}^2 \right] \Big) + \frac{1}{\varepsilon^2} \left(\frac{\hat{\gamma}_{qg}^{(0)}}{6} \left[\gamma_{gq}^{(1)} - (2n_f - 1)\hat{\gamma}_{gq}^{(1)} \right] + \frac{\gamma_{gq}^{(0)} \hat{\gamma}_{qg}^{(1)}}{3} - \frac{\hat{\gamma}_{gg}^{(1)}}{3} \left[4\beta_0 + 7\beta_{0,Q} \right] \right. \\
& + \frac{2\beta_{0,Q}}{3} \left[\gamma_{gg}^{(1)} + \beta_1 + \beta_{1,Q} \right] + \frac{2\gamma_{gg}^{(0)} \beta_{1,Q}}{3} + \delta m_1^{(-1)} \left[-\hat{\gamma}_{qg}^{(0)} \gamma_{gq}^{(0)} - 2\beta_{0,Q} \gamma_{gg}^{(0)} - 10\beta_{0,Q}^2 - 6\beta_{0,Q}\beta_0 \right] \Big) \\
& + \frac{1}{\varepsilon} \left(\frac{\hat{\gamma}_{gg}^{(2)}}{3} - 2(2\beta_0 + 3\beta_{0,Q}) \mathbf{a}_{\mathbf{gg},\mathbf{Q}}^{(\mathbf{2})} - n_f \hat{\gamma}_{qg}^{(0)} \mathbf{a}_{\mathbf{gq},\mathbf{Q}}^{(\mathbf{2})} + \gamma_{gq}^{(0)} \mathbf{a}_{\mathbf{Qg}}^{(\mathbf{2})} + \beta_{1,Q}^{(1)} \gamma_{gg}^{(0)} + \frac{\gamma_{gq}^{(0)} \hat{\gamma}_{qg}^{(0)} \zeta_2}{16} \left[\gamma_{gg}^{(0)} - \gamma_{qq}^{(0)} \right. \right. \\
& \left. \left. + 2(2n_f + 1)\beta_{0,Q} + 6\beta_0 \right] + \frac{\beta_{0,Q} \zeta_2}{4} \left[\gamma_{gg}^{(0)} \{2\beta_0 - \beta_{0,Q}\} + 4\beta_0^2 - 2\beta_{0,Q}\beta_0 - 12\beta_{0,Q}^2 \right] \right. \\
& + \delta m_1^{(-1)} \left[-3\delta m_1^{(-1)} \beta_{0,Q} - 2\delta m_1^{(0)} \beta_{0,Q} - \hat{\gamma}_{gg}^{(1)} \right] + \delta m_1^{(0)} \left[-\hat{\gamma}_{qg}^{(0)} \gamma_{gq}^{(0)} - 2\gamma_{gg}^{(0)} \beta_{0,Q} - 4\beta_{0,Q}\beta_0 - 8\beta_{0,Q}^2 \right] \\
& \left. + 2\delta m_2^{(-1)} \beta_{0,Q} \right) + \boxed{a_{gg,Q}^{(3)}} \Big] .
\end{aligned}$$

→ use for **checking** the ε singular parts

We **confirm** the $n_f T_F^2$ part of the 3-Loop anomalous dimension:

[Moch, Vermaseren, Vogt 2004 Nucl.Phys.B]

$$\begin{aligned}\hat{\gamma}_{gg}^{(2)} &= n_f T_F^2 \mathbf{C_A} \left[-\frac{32 (8N^6 + 24N^5 - 19N^4 - 78N^3 - 253N^2 - 210N - 96)}{27(N-1)N^2(N+1)^2(N+2)} S_1 \right. \\ &\quad \left. - \frac{8 (87N^8 + 348N^7 + 848N^6 + 1326N^5 + 2609N^4 + 3414N^3 + 2632N^2 + 1088N + 192)}{27(N-1)N^3(N+1)^3(N+2)} \right] \\ &\quad + n_f T_F^2 \mathbf{C_F} \left[\frac{64 (N^2 + N + 2)^2}{3(N-1)N^2(N+1)^2(N+2)} (S_1^2 - 3S_2) - \frac{16P_1}{27(N-1)N^4(N+1)^4(N+2)} \right. \\ &\quad \left. + \frac{128 (4N^6 + 3N^5 - 50N^4 - 129N^3 - 100N^2 - 56N - 24)}{9(N-1)N^3(N+1)^3(N+2)} S_1 \right]\end{aligned}$$

$$\begin{aligned}P_1 &= 33N^{10} + 165N^9 + 256N^8 - 542N^7 - 3287N^6 - 8783N^5 - 11074N^4 - 9624N^3 \\ &\quad - 5960N^2 - 2112N - 288\end{aligned}$$


First diagrammatic recalculation

$$\begin{aligned}
A_{gg,Q}^{(3),\overline{\text{MS}}} = & \frac{1}{48} \left\{ \gamma_{gq}^{(0)} \hat{\gamma}_{qg}^{(0)} \left(\gamma_{gg}^{(0)} - \gamma_{qq}^{(0)} - 6\beta_0 - 4n_f\beta_{0,Q} - 10\beta_{0,Q} \right) - 4 \left(\gamma_{gg}^{(0)} [2\beta_0 + 7\beta_{0,Q}] + 4\beta_0^2 + 14\beta_{0,Q}\beta_0 \right. \right. \\
& \left. \left. + 12\beta_{0,Q}^2 \right) \beta_{0,Q} \right\} \ln^3 \left(\frac{\mathbf{m}^2}{\mu^2} \right) + \frac{1}{8} \left\{ \hat{\gamma}_{qg}^{(0)} \left(\gamma_{gq}^{(1)} + (1-n_f)\hat{\gamma}_{gq}^{(1)} \right) + \gamma_{gq}^{(0)} \hat{\gamma}_{qg}^{(1)} - 4\hat{\gamma}_{gg}^{(1)} [\beta_0 + 2\beta_{0,Q}] \right. \\
& \left. + 4\gamma_{gg}^{(1)}\beta_{0,Q} + 4[\beta_1 + \beta_{1,Q}]\beta_{0,Q} + 2\gamma_{gg}^{(0)}\beta_{1,Q} \right\} \ln^2 \left(\frac{\mathbf{m}^2}{\mu^2} \right) + \frac{1}{16} \left\{ 8\hat{\gamma}_{gg}^{(2)} - 8n_f a_{gq,Q}^{(2)} \hat{\gamma}_{qg}^{(0)} + 8\gamma_{gq}^{(0)} a_{Qg}^{(2)} \right. \\
& - 16a_{gg,Q}^{(2)} (2\beta_0 + 3\beta_{0,Q}) + 8\gamma_{gg}^{(0)}\beta_{1,Q}^{(1)} + \gamma_{gq}^{(0)} \hat{\gamma}_{qg}^{(0)} \zeta_2 \left(\gamma_{gg}^{(0)} - \gamma_{qq}^{(0)} + 6\beta_0 + 4n_f\beta_{0,Q} + 6\beta_{0,Q} \right) \\
& \left. + 4\beta_{0,Q}\zeta_2 \left(\gamma_{gg}^{(0)} + 2\beta_0 \right) \left(2\beta_0 + 3\beta_{0,Q} \right) \right\} \ln \left(\frac{\mathbf{m}^2}{\mu^2} \right) + 2(2\beta_0 + 3\beta_{0,Q}) \bar{\mathbf{a}}_{\mathbf{g}\mathbf{g},\mathbf{Q}}^{(2)} + n_f \hat{\gamma}_{qg}^{(0)} \bar{\mathbf{a}}_{\mathbf{g}\mathbf{q},\mathbf{Q}}^{(2)} - \gamma_{gq}^{(0)} \bar{\mathbf{a}}_{\mathbf{Q}\mathbf{g}}^{(2)} \\
& - \beta_{1,Q}^{(2)} \gamma_{gg}^{(0)} + \frac{\gamma_{gq}^{(0)} \hat{\gamma}_{qg}^{(0)} \zeta_3}{48} \left(\gamma_{qq}^{(0)} - \gamma_{gg}^{(0)} - 2[2n_f + 1]\beta_{0,Q} - 6\beta_0 \right) + \frac{\beta_{0,Q}\zeta_3}{12} \left([\beta_{0,Q} - 2\beta_0] \gamma_{gg}^{(0)} \right. \\
& \left. + 2[\beta_0 + 6\beta_{0,Q}]\beta_{0,Q} - 4\beta_0^2 \right) - \frac{\hat{\gamma}_{qg}^{(0)} \zeta_2}{16} \left(\gamma_{gq}^{(1)} + \hat{\gamma}_{gq}^{(1)} \right) + \frac{\beta_{0,Q}\zeta_2}{8} \left(\hat{\gamma}_{gg}^{(1)} - 2\gamma_{gg}^{(1)} - 2\beta_1 - 2\beta_{1,Q} \right) \\
& + \frac{\delta m_1^{(-1)}}{4} \left(8a_{gg,Q}^{(2)} + 24\delta m_1^{(0)}\beta_{0,Q} + 8\delta m_1^{(1)}\beta_{0,Q} + \zeta_2\beta_{0,Q}\beta_0 + 9\zeta_2\beta_{0,Q}^2 \right) + \delta m_1^{(0)} \left(\beta_{0,Q}\delta m_1^{(0)} + \hat{\gamma}_{gg}^{(1)} \right) \\
& + \delta m_1^{(1)} \left(\hat{\gamma}_{qg}^{(0)} \gamma_{gq}^{(0)} + 2\beta_{0,Q}\gamma_{gg}^{(0)} + 4\beta_{0,Q}\beta_0 + 8\beta_{0,Q}^2 \right) - 2\delta m_2^{(0)}\beta_{0,Q} + a_{gg,Q}^{(3)}
\end{aligned}$$

The final renormalized contribution with the $\overline{\text{MS}}$ -mass \bar{m} :

$$\begin{aligned}
A_{gg,Q}^{(3),n_f T_F^2, \overline{\text{MS}}} = & n_f T_F^2 \left\{ \left(\mathbf{C}_F \frac{64(N^2 + N + 2)^2}{9(N-1)N^2(N+1)^2(N+2)} + \mathbf{C}_A \left[\frac{128(N^2 + N + 1)}{27(N-1)N(N+1)(N+2)} - \frac{64}{27} S_1 \right] \right) \ln^3 \left(\frac{\bar{m}^2}{\mu^2} \right) \right. \\
& - \mathbf{C}_F \frac{16}{3} \ln^2 \left(\frac{\bar{m}^2}{\mu^2} \right) + \left(\mathbf{C}_A \frac{1}{(N-1)(N+2)} \left[-\frac{4P_1}{81N^3(N+1)^3} - \frac{16P_2}{81N^2(N+1)^2} S_1 \right] \right. \\
& + \mathbf{C}_F \frac{1}{(N-1)(N+2)} \left[\frac{16(N^2 + N + 2)^2}{N^2(N+1)^2} \left(S_1^2 - \frac{5}{3} S_2 \right) - \frac{4P_3}{9N^4(N+1)^4} - \frac{32P_4}{3N^3(N+1)^3} S_1 \right] \left. \right) \ln \left(\frac{\bar{m}^2}{\mu^2} \right) \\
& + \mathbf{C}_A \frac{1}{(N-1)(N+2)} \left[-\frac{4P_5}{27N^2(N+1)^2} S_1^2 - \frac{8P_6}{729N^3(N+1)^3} S_1 + \frac{512}{27}(N-1)(N+2) \zeta_3 S_1 \right. \\
& \quad \left. - \frac{2P_7}{729N^4(N+1)^4} - \frac{1024(N^2 + N + 1)}{27N(N+1)} \zeta_3 + \frac{4P_8}{27N^2(N+1)^2} S_2 \right] \\
& + \mathbf{C}_F \frac{1}{(N-1)(N+2)} \left[\frac{64(N^2 + N + 2)^2}{9N^2(N+1)^2} \left(-\frac{1}{3} S_1^3 - 8\zeta_3 + \frac{4}{3} S_3 \right) + \frac{32P_9}{27N^3(N+1)^3} S_1^2 \right. \\
& \quad \left. - \frac{64P_{10}}{81N^4(N+1)^4} S_1 - \frac{32P_{11}}{243N^5(N+1)^5} - \frac{32P_{12}}{3N^3(N+1)^3} S_2 \right] \left. \right\}
\end{aligned}$$

The $O(n_f T_F^2 \alpha_s^3)$ contributions to $A_{gq,Q}$



The all- ε result constituting the color factor $T_F^2 n_f C_F$

$$\hat{A}_{gq,T_F^2 n_f}^{(3)} = -96 a_s^3 T_F^2 n_f C_F \left(\frac{m^2}{\mu^2} \right)^{\frac{3\varepsilon}{2}} S_\varepsilon^3 \frac{1 + (-1)^N}{2} e^{-\frac{3\varepsilon}{2}\gamma} \frac{(\varepsilon - 1)^2 (\varepsilon + 2) (\varepsilon + N^2 + N + 2)}{\varepsilon (\varepsilon + 1) (\varepsilon + 3)} \\ \times \Gamma(1 - \varepsilon)^2 \Gamma\left(-\frac{\varepsilon}{2} - 4\right) \Gamma\left(\frac{\varepsilon}{2} + 2\right) \frac{\Gamma\left(\frac{\varepsilon}{2} + 5\right) \Gamma\left(-\frac{3\varepsilon}{2}\right) \Gamma(N - 1)}{\Gamma(4 - 2\varepsilon) \Gamma\left(\frac{\varepsilon}{2} + N + 2\right)}$$

yields the renormalized contribution

$$A_{gq,Q}^{(3),n_f T_F^2, \overline{\text{MS}}} = n_f T_F^2 \frac{1 + (-1)^N}{2} \left\{ \mathbf{C}_F \frac{32 (N^2 + N + 2)}{9(N-1)N(N+1)} \ln^3 \left(\frac{\bar{m}^2}{\mu^2} \right) + \mathbf{C}_F \left[-\frac{16 (N^2 + N + 2)}{3(N-1)N(N+1)} (S_1^2 + S_2) \right. \right. \\ \left. + \frac{32 (8N^3 + 13N^2 + 27N + 16)}{9(N-1)N(N+1)^2} S_1 + \frac{32 (19N^4 + 81N^3 + 86N^2 + 80N + 38)}{27(N-1)N(N+1)^3} \right] \ln \left(\frac{\bar{m}^2}{\mu^2} \right) \\ + \mathbf{C}_F \left[\frac{32 (N^2 + N + 2)}{27(N-1)N(N+1)} (S_1^3 + 3S_2 S_1 + 2[S_3] - 24\zeta_3) - \frac{32 (8N^3 + 13N^2 + 27N + 16)}{27(N-1)N(N+1)^2} (S_1^2 + S_2) \right. \\ \left. + \frac{64 (4N^4 + 4N^3 + 23N^2 + 25N + 8)}{27(N-1)N(N+1)^3} S_1 + \frac{64 (197N^5 + 824N^4 + 1540N^3 + 1961N^2 + 1388N + 394)}{243(N-1)N(N+1)^4} \right] \right\}$$

Here we **confirm** the n_f contribution to the anomalous dimension:

[Moch, Vermaseren, Vogt 2004 Nucl.Phys.B]

$$\begin{aligned}\hat{\gamma}_{gq}^{(2),n_f} &= n_f T_F^2 C_F \left(\frac{64(N^2 + N + 2)}{3(N-1)N(N+1)} - (S_1^2 + S_2) + \frac{128(8N^3 + 13N^2 + 27N + 16)}{9(N-1)N(N+1)^2} S_1 \right. \\ &\quad \left. - \frac{128(4N^4 + 4N^3 + 23N^2 + 25N + 8)}{9(N-1)N(N+1)^3} \right)\end{aligned}$$

in an independent calculation.

Furthermore we are able to **check** a result for the combination

$$\tilde{\gamma}_{gg}^{(2)} + \frac{\tilde{\gamma}_{gq}^{(2)} \gamma_{qg}^{(0)}}{\tilde{\gamma}_{gg}^{(0)} n_f}$$

of 3-loop anomalous dimensions, derived from the **large n_f** expansion in QCD by [Bennett, Gracey 1997]; where we denote with $\tilde{\gamma}_{ij}^{(k)}$ the leading n_f coefficient of $\gamma_{ij}^{(k)}$.

Graphs with m_c and m_b


$$\begin{aligned}
a_{Qg}^{(3)}(N=6) = & T_F^2 C_A \left\{ \frac{69882273800453}{367569090000} - \frac{395296}{19845} \zeta_3 + \frac{1316809}{39690} \zeta_2 + \frac{832369820129}{14586075000} x + \frac{1511074426112}{624023544375} x^2 - \frac{84840004938801319}{690973782403905000} x^3 \right. \\
& + \ln\left(\frac{m_2^2}{\mu^2}\right) \left[\frac{11771644229}{194481000} + \frac{78496}{2205} \zeta_2 - \frac{1406143531}{69457500} x - \frac{105157957}{180093375} x^2 + \frac{2287164970759}{7669816654500} x^3 \right] \\
& + \ln^2\left(\frac{m_2^2}{\mu^2}\right) \left[\frac{2668087}{79380} + \frac{112669}{661500} x - \frac{49373}{51975} x^2 - \frac{31340489}{34054020} x^3 \right] + \ln^3\left(\frac{m_2^2}{\mu^2}\right) \frac{324148}{19845} + \ln^2\left(\frac{m_2^2}{\mu^2}\right) \ln\left(\frac{m_1^2}{\mu^2}\right) \frac{156992}{6615} \\
& + \ln\left(\frac{m_2^2}{\mu^2}\right) \ln\left(\frac{m_1^2}{\mu^2}\right) \left[\frac{128234}{3969} - \frac{112669}{330750} x + \frac{98746}{51975} x^2 + \frac{31340489}{17027010} x^3 \right] + \ln\left(\frac{m_2^2}{\mu^2}\right) \ln^2\left(\frac{m_1^2}{\mu^2}\right) \frac{68332}{6615} \\
& + \ln\left(\frac{m_1^2}{\mu^2}\right) \left[\frac{83755534727}{583443000} + \frac{78496}{2205} \zeta_2 + \frac{1406143531}{69457500} x + \frac{105157957}{180093375} x^2 - \frac{2287164970759}{7669816654500} x^3 \right] \\
& + \ln^2\left(\frac{m_1^2}{\mu^2}\right) \left[\frac{2668087}{79380} + \frac{112669}{661500} x - \frac{49373}{51975} x^2 - \frac{31340489}{34054020} x^3 \right] + \ln^3\left(\frac{m_1^2}{\mu^2}\right) \frac{412808}{19845} \Big\} \\
& + T_F^2 C_F \left\{ - \frac{3161811182177}{71471767500} + \frac{447392}{19845} \zeta_3 + \frac{9568018}{4862025} \zeta_2 - \frac{64855635472}{2552563125} x + \frac{1048702178522}{97070329125} x^2 + \frac{1980566069882672}{2467763508585375} x^3 \right. \\
& + \ln\left(\frac{m_2^2}{\mu^2}\right) \left[\frac{1786067629}{204205050} - \frac{111848}{15435} \zeta_2 - \frac{128543024}{24310125} x - \frac{22957168}{3361743} x^2 - \frac{2511536080}{2191376187} x^3 \right] \\
& + \ln^2\left(\frac{m_2^2}{\mu^2}\right) \left[\frac{3232799}{4862025} + \frac{752432}{231525} x + \frac{177944}{40425} x^2 + \frac{127858928}{42567525} x^3 \right] - \ln^3\left(\frac{m_2^2}{\mu^2}\right) \frac{111848}{19845} - \ln^2\left(\frac{m_2^2}{\mu^2}\right) \ln\left(\frac{m_1^2}{\mu^2}\right) \frac{223696}{46305} \\
& + \ln\left(\frac{m_2^2}{\mu^2}\right) \ln\left(\frac{m_1^2}{\mu^2}\right) \left[\frac{22238456}{4862025} - \frac{1504864}{231525} x - \frac{355888}{40425} x^2 - \frac{255717856}{42567525} x^3 \right] + \ln\left(\frac{m_2^2}{\mu^2}\right) \ln^2\left(\frac{m_1^2}{\mu^2}\right) \frac{223696}{46305} \\
& + \ln\left(\frac{m_1^2}{\mu^2}\right) \left[- \frac{24797875607}{1021025250} - \frac{111848}{15435} \zeta_2 + \frac{128543024}{24310125} x + \frac{22957168}{3361743} x^2 + \frac{2511536080}{2191376187} x^3 \right] \\
& \left. + \ln^2\left(\frac{m_1^2}{\mu^2}\right) \left[\frac{3232799}{4862025} + \frac{752432}{231525} x + \frac{177944}{40425} x^2 + \frac{127858928}{42567525} x^3 \right] - \ln^3\left(\frac{m_1^2}{\mu^2}\right) \frac{1230328}{138915} \right\} + O(x^4 \ln^3(x))
\end{aligned}$$

These moments have been calculated referring [qexp](#) by Steinhauser et al. [with operator insertions](#). Despite being **universal**, these contribution do not belong to the charm or bottom PDF. **This is then the end of the VFNF.**

Calculation of Convergent Massive 3-Loop Graphs

- Aim:
 - Compute fixed Mellin moments of convergent 3-loop diagrams
 - Find general N representations for all convergent 3-loop topologies
- We use the α -representation to solve the momentum integrals and obtain integrals
- The corresponding graph polynomials of a graph G are given by
 - $U = \sum_T \prod_{l \notin T} \alpha_l$, where T denotes the spanning trees of G
 - $V = \sum_{l \in massive} \alpha_l$
 - different Dodgson polynomials, which can be derived from the corresponding tadpole diagram, for the operator insertions

General Values of N



$$\begin{aligned}
 I(x) &= \frac{1}{(1+N)(2+N)x} \left\{ \zeta_3 \left[2L(\{-1\}, x) - 2(-1+2x)L(\{1\}, x) - 4L(\{1, 1\}, x) \right] - 3L(\{-1, 0, 0, 1\}, x) \right. \\
 &\quad + 2L(\{-1, 0, 1, 1\}, x) - 2xL(\{0, 0, 1, 1\}, x) + 3xL(\{0, 1, 0, 1\}, x) - xL(\{0, 1, 1, 1\}, x) \\
 &\quad + (-3+2x)L(\{1, 0, 0, 1\}, x) + 2xL(\{1, 0, 1, 1\}, x) - (-1+5x)L(\{1, 1, 0, 1\}, x) + xL(\{1, 1, 1, 1\}, x) \\
 &\quad - 2L(\{1, 0, 0, 1, 1\}, x) + 3L(\{1, 0, 1, 0, 1\}, x) - L(\{1, 0, 1, 1, 1\}, x) + 2L(\{1, 1, 0, 0, 1\}, x) \\
 &\quad \left. + 2L(\{1, 1, 0, 1, 1\}, x) - 5L(\{1, 1, 1, 0, 1\}, x) + L(\{1, 1, 1, 1, 1\}, x) \right\} \\
 I(N) &= \frac{1}{(N+1)(N+2)(N+3)} \left\{ \frac{648 + 1512N + 1458N^2 + 744N^3 + 212N^4 + 32N^5 + 2N^6}{(1+N)^3(2+N)^3(3+N)^3} \right. \\
 &\quad - \frac{2(-1 + (-1)^N + N + (-1)^N N)}{(1+N)} \zeta_3 - (-1)^N S_{-3} - \frac{N}{6(1+N)} S_1^3 + \frac{1}{24} S_1^4 \\
 &\quad - \frac{(7 + 22N + 10N^2)}{2(1+N)^2(2+N)} S_2 - \frac{19}{8} S_2^2 - \frac{1 + 4N + 2N^2}{2(1+N)^2(2+N)} S_1^2 + \frac{9}{4} S_2 - \frac{(-9 + 4N)}{3(1+N)} S_3 \\
 &\quad - \frac{1}{4} S_4 - 2(-1)^N S_{-2,1} + \frac{(-1 + 6N)}{(1+N)} S_{2,1} + \frac{54 + 207N + 246N^2 + 130N^3 + 32N^4 + 3N^5}{(1+N)^3(2+N)^2(3+N)^2} S_1 \\
 &\quad \left. + 4\zeta_3 S_1 - \frac{(-2 + 7N)}{2(1+N)} S_2 S_1 + \frac{13}{3} S_3 S_1 - 7S_{2,1} S_1 - 7S_{3,1} + 10S_{2,1,1} \right\}
 \end{aligned}$$

Conclusions

- A series of moments for the transition matrix elements A_{ij} at 3-loop order were given in [Bierenbaum, Blümlein, Klein 2009 Nucl. Phys. B].
- The corresponding quarkonic 3-loop contributions of $O(n_f T_F^2 C_{A,F})$ to A_{qq} and A_{qg} were calculated in [Ablinger, Blümlein, Klein, Schneider, Wißbrock 2011 Nucl. Phys. B]. Now also $A_{gg,Q}$ and $A_{gq,Q}$ have been obtained for these color coefficients at general N .
- A series of OMEs were fully calculated A_{qq} and A_{qg} in $O(T_F^2 C_{A,F})$
- Ladder topologies, including poles, are currently calculated using **Sigma**.
- The moments $\textcolor{blue}{N = 2,4,6}$ have been calculated for graphs depending on both m_c and m_b . Starting with 3-loops, graphs exist which conflict with the ideology of the VFNS.
This was expected, since the heavy quarks are produced in the final state!
- With the help of Hyperlogarithms non-divergent 3-Loop graphs can be calculated, if moments are considered. For general values of $\textcolor{red}{N}$ first analytic results have been obtained, including Benz-topologies, performing the calculation automatically.