

On the pair correlations of neutral K , D , B and B_s mesons with close momenta produced in inclusive multiparticle processes

Valery V. Lyuboshitz[†], Vladimir L. Lyuboshitz

Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia

[†] E-mail: Valery.Lyuboshitz@jinr.ru

DOI: <http://dx.doi.org/10.3204/DESY-PROC-2012-02/17>

The phenomenological structure of inclusive cross-sections of the production of two neutral K mesons in collisions of hadrons and nuclei is investigated taking into account the strangeness conservation in strong and electromagnetic interactions. Relations describing the dependence of the correlations of two short-lived and two long-lived neutral kaons $K_S^0 K_S^0$, $K_L^0 K_L^0$ and the correlations of “mixed” pairs $K_S^0 K_L^0$ at small relative momenta upon the space-time parameters of the generation region of K^0 and \bar{K}^0 mesons have been obtained. It is shown that under the strangeness conservation the correlation functions of the pairs $K_S^0 K_S^0$ and $K_L^0 K_L^0$, produced in the same inclusive process, coincide, and the difference between the correlation functions of the pairs $K_S^0 K_S^0$ and $K_S^0 K_L^0$ is conditioned by the production of the pairs of non-identical neutral kaons $K^0 \bar{K}^0$. Analogous correlations for the pairs of neutral heavy mesons D^0 , B^0 and B_s^0 , generated in multiple processes with the charm (beauty) conservation, are analyzed, and differences from the case of neutral K mesons are discussed.

1 Consequences of the strangeness conservation for neutral kaons

In the work [1] the properties of the density matrix of two neutral K mesons, following from the strangeness conservation in strong and electromagnetic interactions, have been investigated. By definition, the diagonal elements of the non-normalized two-particle density matrix coincide with the two-particle structure functions, which are proportional to the double inclusive cross-sections.

Strangeness is the additive quantum number. Taking into account the strangeness conservation, the pairs of neutral kaons $K^0 K^0$ (strangeness $S = +2$), $\bar{K}^0 \bar{K}^0$ (strangeness $S = -2$) and $K^0 \bar{K}^0$ (strangeness $S = 0$) are produced incoherently. This means that in the K^0 - \bar{K}^0 -representation the non-diagonal elements of the density matrix between the states $K^0 K^0$ and $\bar{K}^0 \bar{K}^0$, $K^0 K^0$ and $K^0 \bar{K}^0$, $\bar{K}^0 \bar{K}^0$ and $K^0 \bar{K}^0$ are equal to zero. However, the non-diagonal elements of the two-kaon density matrix between the two states $|K^0\rangle(\mathbf{p}_1)|\bar{K}^0\rangle(\mathbf{p}_2)$ and $|\bar{K}^0\rangle(\mathbf{p}_1)|K^0\rangle(\mathbf{p}_2)$ with the zero strangeness are not equal to zero, in general. Here \mathbf{p}_1 and \mathbf{p}_2 are the momenta of the first and second kaons.

The internal states of K^0 meson ($S = 1$) and \bar{K}^0 meson ($S = -1$) are the superpositions of

the states $|K_S^0\rangle$ and $|K_L^0\rangle$, where K_S^0 is the short-lived neutral kaon and K_L^0 is the long-lived one. Neglecting the small effect of CP non-invariance, the CP -parity of the state K_S^0 is equal to $(+1)$, and the CP -parity of the state K_L^0 is equal to (-1) ; in doing so,

$$|K^0\rangle = \frac{1}{\sqrt{2}}(|K_S^0\rangle + |K_L^0\rangle), \quad |\bar{K}^0\rangle = \frac{1}{\sqrt{2}}(|K_S^0\rangle - |K_L^0\rangle).$$

It is clear that both the quasistationary states of the neutral kaon have no definite strangeness.

It follows from the Bose-symmetry of the wave function of two neutral kaons with respect to the total permutation of internal states and momenta that the CP -parity of the system $K^0\bar{K}^0$ is always positive [2] (the C -parity is $(-1)^L$, the space parity is $P = (-1)^L$, where L is the orbital momentum).

The system of two non-identical neutral kaons $K^0\bar{K}^0$ in the symmetric internal state, corresponding to even orbital momenta, is decomposed into the schemes $|K_S^0\rangle|K_S^0\rangle$ and $|K_L^0\rangle|K_L^0\rangle$ [2]:

$$\begin{aligned} |\psi^+\rangle &= \frac{1}{\sqrt{2}}(|K^0\rangle^{(\mathbf{p}_1)} \otimes |\bar{K}^0\rangle^{(\mathbf{p}_2)} + |\bar{K}^0\rangle^{(\mathbf{p}_1)} \otimes |K^0\rangle^{(\mathbf{p}_2)}) = \\ &= \frac{1}{\sqrt{2}}(|K_S^0\rangle^{(\mathbf{p}_1)} \otimes |K_S^0\rangle^{(\mathbf{p}_2)} - |K_L^0\rangle^{(\mathbf{p}_1)} \otimes |K_L^0\rangle^{(\mathbf{p}_2)}); \end{aligned} \quad (1)$$

meantime, the system $K^0\bar{K}^0$ in the antisymmetric internal state, corresponding to odd orbital momenta, is decomposed into the scheme $|K_S^0\rangle|K_L^0\rangle$ [2]:

$$\begin{aligned} |\psi^-\rangle &= \frac{1}{\sqrt{2}}(|K^0\rangle^{(\mathbf{p}_1)} \otimes |\bar{K}^0\rangle^{(\mathbf{p}_2)} - |\bar{K}^0\rangle^{(\mathbf{p}_1)} \otimes |K^0\rangle^{(\mathbf{p}_2)}) = \\ &= \frac{1}{\sqrt{2}}(|K_S^0\rangle^{(\mathbf{p}_1)} \otimes |K_L^0\rangle^{(\mathbf{p}_2)} - |K_L^0\rangle^{(\mathbf{p}_1)} \otimes |K_S^0\rangle^{(\mathbf{p}_2)}). \end{aligned} \quad (2)$$

The strangeness conservation leads to the fact that all the double inclusive cross-sections of production of pairs $K_S^0K_S^0$, $K_L^0K_L^0$ and $K_S^0K_L^0$ (two-particle structure functions) prove to be symmetric with respect to the permutation of momenta \mathbf{p}_1 and \mathbf{p}_2 .

Besides, due to the strangeness conservation, the structure functions of neutral K mesons produced in inclusive processes are invariant with respect to the replacement of the short-lived state K_S^0 by the long-lived state K_L^0 , and *vice versa* [1]:

$$\begin{aligned} f_{SS}(\mathbf{p}_1, \mathbf{p}_2) = f_{LL}(\mathbf{p}_1, \mathbf{p}_2) &= \frac{1}{4} [f_{K^0K^0}(\mathbf{p}_1, \mathbf{p}_2) + f_{\bar{K}^0\bar{K}^0}(\mathbf{p}_1, \mathbf{p}_2) + \\ &+ f_{K^0\bar{K}^0}(\mathbf{p}_1, \mathbf{p}_2) + f_{\bar{K}^0K^0}(\mathbf{p}_1, \mathbf{p}_2)] + \frac{1}{2} \text{Re} \rho_{K^0\bar{K}^0 \rightarrow \bar{K}^0K^0}(\mathbf{p}_1, \mathbf{p}_2), \end{aligned} \quad (3)$$

$$\begin{aligned} f_{SL}(\mathbf{p}_1, \mathbf{p}_2) = f_{LS}(\mathbf{p}_1, \mathbf{p}_2) &= \frac{1}{4} [f_{K^0K^0}(\mathbf{p}_1, \mathbf{p}_2) + f_{\bar{K}^0\bar{K}^0}(\mathbf{p}_1, \mathbf{p}_2) + \\ &+ f_{K^0\bar{K}^0}(\mathbf{p}_1, \mathbf{p}_2) + f_{\bar{K}^0K^0}(\mathbf{p}_1, \mathbf{p}_2)] - \frac{1}{2} \text{Re} \rho_{K^0\bar{K}^0 \rightarrow \bar{K}^0K^0}(\mathbf{p}_1, \mathbf{p}_2), \end{aligned} \quad (4)$$

where $\rho_{K^0\bar{K}^0 \rightarrow \bar{K}^0K^0}(\mathbf{p}_1, \mathbf{p}_2) = (\rho_{\bar{K}^0K^0 \rightarrow K^0\bar{K}^0}(\mathbf{p}_1, \mathbf{p}_2))^*$ are the non-diagonal elements of the two-kaon density matrix. The difference between the two-particle structure functions f_{SS} and f_{SL} is connected just with the contribution of these non-diagonal elements.

2 Structure of pair correlations of identical and non-identical neutral kaons with close momenta

Now let us consider, within the model of one-particle sources [2-7], the correlations of pairs of neutral K mesons with close momenta (see also [8-10]). In the case of the identical states $K_S^0 K_S^0$ and $K_L^0 K_L^0$ we obtain the following expressions for the correlation functions R_{SS} , R_{LL} (proportional to the structure functions), normalized to unity at large relative momenta:

$$R_{SS}(\mathbf{k}) = R_{LL}(\mathbf{k}) = \lambda_{K^0 K^0} [1 + F_{K^0}(2\mathbf{k}) + 2 b_{\text{int}}(\mathbf{k})] + \lambda_{\bar{K}^0 \bar{K}^0} [1 + F_{\bar{K}^0}(2\mathbf{k}) + 2 \tilde{b}_{\text{int}}(\mathbf{k})] + \lambda_{K^0 \bar{K}^0} [1 + F_{K^0 \bar{K}^0}(2\mathbf{k}) + 2 B_{\text{int}}(\mathbf{k})]. \quad (5)$$

Here \mathbf{k} is the momentum of one of the kaons in the c.m. frame of the pair, and the quantities $\lambda_{K^0 K^0}$, $\lambda_{\bar{K}^0 \bar{K}^0}$ and $\lambda_{K^0 \bar{K}^0}$ are the relative fractions of the average numbers of produced pairs $K^0 K^0$, $\bar{K}^0 \bar{K}^0$ and $K^0 \bar{K}^0$, respectively ($\lambda_{K^0 K^0} + \lambda_{\bar{K}^0 \bar{K}^0} + \lambda_{K^0 \bar{K}^0} = 1$). The ‘‘formfactors’’ $F_{K^0}(2\mathbf{k})$, $F_{\bar{K}^0}(2\mathbf{k})$ and $F_{K^0 \bar{K}^0}(2\mathbf{k})$ appear due to the contribution of Bose-statistics:

$$F_{K^0}(2\mathbf{k}) = \int W_{K^0}(\mathbf{r}) \cos(2\mathbf{k}\mathbf{r}) d^3\mathbf{r}, \quad F_{\bar{K}^0}(2\mathbf{k}) = \int W_{\bar{K}^0}(\mathbf{r}) \cos(2\mathbf{k}\mathbf{r}) d^3\mathbf{r}, \\ F_{K^0 \bar{K}^0}(2\mathbf{k}) = \int W_{K^0 \bar{K}^0}(\mathbf{r}) \cos(2\mathbf{k}\mathbf{r}) d^3\mathbf{r}. \quad (6)$$

where $W_{K^0}(\mathbf{r})$, $W_{\bar{K}^0}(\mathbf{r})$ and $W_{K^0 \bar{K}^0}(\mathbf{r})$ are the probability distributions of distances between the sources of emission of two K^0 mesons, between the sources of emission of two \bar{K}^0 mesons and between the sources of emission of the K^0 meson and \bar{K}^0 meson, respectively, in the c.m. frame of the kaon pair. Meantime, the quantity $b_{\text{int}}(\mathbf{k})$ describes the contribution of the S -wave interaction of two K^0 mesons, the quantity $\tilde{b}_{\text{int}}(\mathbf{k})$ describes the contribution of the S -wave interaction of two \bar{K}^0 mesons and the quantity $B_{\text{int}}(\mathbf{k})$ describes the contribution of the S -wave interaction of the K^0 meson with the \bar{K}^0 meson. Due to the CP invariance, the quantities $b_{\text{int}}(\mathbf{k})$ and $\tilde{b}_{\text{int}}(\mathbf{k})$ can be expressed by means of averaging the same function $b(\mathbf{k}, \mathbf{r})$ over the different distributions:

$$b_{\text{int}}(\mathbf{k}) = \int W_{K^0}(\mathbf{r}) b(\mathbf{k}, \mathbf{r}) d^3\mathbf{r}, \quad \tilde{b}_{\text{int}}(\mathbf{k}) = \int W_{\bar{K}^0}(\mathbf{r}) b(\mathbf{k}, \mathbf{r}) d^3\mathbf{r}.$$

The quantity $B_{\text{int}}(\mathbf{k})$ has the structure : $B_{\text{int}}(\mathbf{k}) = \int W_{K^0 \bar{K}^0}(\mathbf{r}) B(\mathbf{k}, \mathbf{r}) d^3\mathbf{r}$, where $B(\mathbf{k}, \mathbf{r}) \neq b(\mathbf{k}, \mathbf{r})$.

Let us emphasize that when the pair of non-identical neutral kaons $K^0 \bar{K}^0$ is produced but the pair of identical quasistationary states $K_S^0 K_S^0$ (or $K_L^0 K_L^0$) is registered over decays, the two-particle correlations at small relative momenta have the same character as in the case of usual identical bosons with zero spin [2].

For the pairs of non-identical kaon states $K_S^0 K_L^0$ the correlation functions at small relative momenta have the form:

$$R_{SL}(\mathbf{k}) = R_{LS}(\mathbf{k}) = \lambda_{K^0 K^0} [1 + F_{K^0}(2\mathbf{k}) + 2 b_{\text{int}}(\mathbf{k})] + \lambda_{\bar{K}^0 \bar{K}^0} [1 + F_{\bar{K}^0}(2\mathbf{k}) + 2 \tilde{b}_{\text{int}}(\mathbf{k})] + \lambda_{K^0 \bar{K}^0} [1 - F_{K^0 \bar{K}^0}(2\mathbf{k})]. \quad (7)$$

It follows from Eqs.(5) and (7) that the correlation functions of pairs of neutral K mesons with close momenta, which are created in inclusive processes, satisfy the relation

$$\begin{aligned} R_{SS}(\mathbf{k}) + R_{LL}(\mathbf{k}) - R_{SL}(\mathbf{k}) - R_{LS}(\mathbf{k}) &= 2 [R_{SS}(\mathbf{k}) - R_{SL}(\mathbf{k})] = \\ &= 4\lambda_{K^0\bar{K}^0} [F_{K^0\bar{K}^0}(2\mathbf{k}) + B_{\text{int}}(\mathbf{k})]. \end{aligned} \quad (8)$$

We see that the difference between the correlation functions of the pairs of identical neutral kaons $K_S^0 K_S^0$ and pairs of non-identical neutral kaons $K_S^0 K_L^0$ is conditioned exclusively by the generation of $K^0 \bar{K}^0$ -pairs.

The relations connecting the contribution of the S -wave strong interaction into the pair correlations of particles at small relative momenta with the parameters of low-energy scattering were obtained earlier in the papers [4-7]. It is essential that the ‘‘formfactors’’ (6) and the functions $b_{\text{int}}(\mathbf{k})$, $\tilde{b}_{\text{int}}(\mathbf{k})$ and $B_{\text{int}}(\mathbf{k})$ depend on the space-time parameters of the generation region of neutral kaons and tend to zero at high values of the relative momentum $q = 2|\mathbf{k}|$ of two neutral kaons. Concretely, the expression for the function $B(\mathbf{k}, \mathbf{r})$ in the case of the $K^0 \bar{K}^0$ system has been obtained in the paper [10]. In the same paper, the estimate of contribution of the transition $K^+ K^- \rightarrow K^0 \bar{K}^0$ has also been presented .

3 Correlations of neutral heavy mesons

Formally, analogous relations are valid also for the neutral heavy mesons D^0 , B^0 and B_s^0 . In doing so, the role of strangeness conservation is played, respectively, by the conservation of charm and beauty in inclusive multiple processes with production of these mesons . In these cases the quasistationary states are also states with definite CP parity, neglecting the effects of CP nonconservation. For example,

$$|B_S^0\rangle = \frac{1}{\sqrt{2}}(|B^0\rangle + |\bar{B}^0\rangle), \quad CP \text{ parity } (+1); \quad |B_L^0\rangle = \frac{1}{\sqrt{2}}(|B^0\rangle - |\bar{B}^0\rangle), \quad CP \text{ parity } (-1).$$

The difference of masses between the respective CP -odd and CP -even states is very insignificant in all the cases, ranging from 10^{-12} MeV for K^0 mesons up to 10^{-8} MeV for B_s^0 mesons . Concerning the lifetimes of these states, in the case of K^0 mesons they differ by 600 times, but for D^0 , B^0 and B_s^0 mesons the respective lifetimes are almost the same. In connection with this, it is practically impossible to distinguish the states of D^0 , B^0 and B_s^0 mesons with definite CP parity by the difference in their lifetimes. These states, in principle, can be identified through the purely CP -even and purely CP -odd decay channels ; however, in fact the branching ratio for such decays is very small . For example,

$$Br(D^0 \rightarrow \pi^+ \pi^-) = 1.62 \cdot 10^{-3} \quad (CP = +1); \quad Br(D^0 \rightarrow K^+ K^-) = 4.25 \cdot 10^{-3} \quad (CP = +1);$$

$$Br(B_s^0 \rightarrow J/\Psi \pi^0) < 1.2 \cdot 10^{-3} \quad (CP = +1); \quad Br(B^0 \rightarrow J/\Psi K_S^0) = 9 \cdot 10^{-4} \quad (CP = -1) .$$

Just as in the case of neutral K mesons, the correlation functions for the pairs of states of neutral D , B and B_s mesons with the same CP parity ($R_{SS} = R_{LL}$) and for the pairs of states with different CP parity (R_{SL}) do not coincide, and the difference between them is conditioned exclusively by the production of pairs $D^0 \bar{D}^0$, $B^0 \bar{B}^0$ and $B_s^0 \bar{B}_s^0$, respectively. In particular, for B_s^0 mesons the following relation holds:

$$R_{SS}(\mathbf{k}) - R_{SL}(\mathbf{k}) = 2\lambda_{B_s^0 \bar{B}_s^0} \left[F_{B_s^0 \bar{B}_s^0}(2\mathbf{k}) + B_{\text{int}}(\mathbf{k}) \right]; \quad (9)$$

here $\lambda_{B_s^0 \bar{B}_s^0}$ is the relative fraction of generated pairs $B_s^0 \bar{B}_s^0$,

$$F_{B_s^0 \bar{B}_s^0}(\mathbf{2k}) = \int W_{B_s^0 \bar{B}_s^0}(\mathbf{r}) \cos(2\mathbf{k}\mathbf{r}) d^3\mathbf{r}, \quad B_{\text{int}}(\mathbf{k}) = \int W_{B_s^0 \bar{B}_s^0}(\mathbf{r}) B(\mathbf{k}, \mathbf{r}) d^3\mathbf{r},$$

$$B(\mathbf{k}, \mathbf{r}) = |A_{B_s^0 \bar{B}_s^0}(k)|^2 \frac{1}{r^2} + 2 \operatorname{Re} \left(A_{B_s^0 \bar{B}_s^0}(k) \frac{\exp(ikr) \cos \mathbf{k}\mathbf{r}}{r} \right),$$

where $A_{B_s^0 \bar{B}_s^0}(k) \equiv A_{B_s^0 \bar{B}_s^0 \rightarrow B_s^0 \bar{B}_s^0}(k)$ is the amplitude of S -wave $B_s^0 \bar{B}_s^0$ - scattering, $k = |\mathbf{k}|$, $r = |\mathbf{r}|$. Let us remark that the B_s^0 and \bar{B}_s^0 mesons do not have charged partners (the isotopic spin equals zero) and, on account of that, in the given case the transition similar to $K^+ K^- \rightarrow K^0 \bar{K}^0$ is absent .

4 Summary

1. It is shown that, taking into account the strangeness conservation, the correlation functions for two short-lived neutral K mesons (R_{SS}) and two long-lived neutral K mesons (R_{LL}) are equal to each other. This result is the direct consequence of the strangeness conservation.
2. It is shown that the production of $K^0 \bar{K}^0$ -pairs with the zero strangeness leads to the difference between the correlation functions R_{SS} and R_{SL} of two neutral kaons.
3. The character of analogous correlations for neutral heavy mesons D^0 , B^0 , B_s^0 with nonzero charm and beauty is discussed . Contrary to the case of K^0 mesons, here the distinction of respective CP -even and CP -odd states encounters difficulties, which are connected with the insignificant difference of their lifetimes and the relatively small probability of purely CP -even and purely CP -odd decay channels .

References

- [1] V. L. Lyuboshitz. *Yad. Fiz.* **23** (1976) 1266 [*Sov. J. Nucl. Phys.* **23** (1976) 673].
- [2] V. L. Lyuboshitz and M. I. Podgoretsky. *Yad. Fiz.* **30** (1979) 789 [*Sov. J. Nucl. Phys.* **30** (1979) 407].
- [3] M. I. Podgoretsky. *Fiz. Elem. Chast. At. Yadra* **20** (1989) 628 [*Sov. J. Part. Nucl.* **20** (1989) 266].
- [4] R. Lednicky and V. L. Lyuboshitz. *Yad. Fiz.* **35** (1982) 1316 [*Sov. J. Nucl. Phys.* **35** (1982) 770].
- [5] V. L. Lyuboshitz. *Yad. Fiz.* **41** (1985) 820 [*Sov. J. Nucl. Phys.* **41** (1985) 523].
- [6] V. L. Lyuboshitz. *Yad. Fiz.* **48** (1988) 1501 [*Sov. J. Nucl. Phys.* **48** (1988) 956].
- [7] R. Lednicky, V. V. Lyuboshitz and V. L. Lyuboshitz. *Yad. Fiz.* **61** (1998) 2161 [*Phys. At. Nucl.* **61** (1998) 2050].
- [8] V. L. Lyuboshitz and V. V. Lyuboshitz. *Nukleonika*, v.**51**, Suppl. 3 (2006), p.S69 - in Proceedings of Quark Matter 2005 Conference (Budapest, Hungary, August 4 - 9, 2005).
- [9] V. L. Lyuboshitz and V. V. Lyuboshitz, in Proceedings of the IV International Conference on Quarks and Nuclear Physics – QNP-06 (Madrid, Spain, June 5 - 10, 2006), Berlin - Heidelberg, 2007, p.109 .
- [10] V. L. Lyuboshitz and V. V. Lyuboshitz. *Pis'ma v EchaYa*, **4** (2007) 654 [*Physics of Particles and Nuclei Letters*, **4** (2007) 388].