

THE INFRARED STRUCTURE OF GAUGE AMPLITUDES IN THE HIGH-ENERGY LIMIT

Lorenzo Magnea

University of Torino - INFN Torino

RadCor 2011, Mamallapuram, 29/09/11




Outline

- Infrared divergences to all orders
- The dipole formula (with E. Gardi)
- The high-energy limit (with V. Del Duca, C. Duhr, E. Gardi, C. White)
- Reggeization and beyond
- Outlook

ON INFRARED DIVERGENCES



Textbook theory ...



Singularities arise only when propagators go **on shell**

$$2p \cdot k = 2p_0 k_0 (1 - \cos \theta_{pk}) = 0,$$

$$\rightarrow k_0 = 0 \text{ (IR)}; \quad \cos \theta_{pk} = 1.$$

- ➔ Emission is **not suppressed** at long distances
- ➔ Isolated charged particles are **not true asymptotic states** of unbroken gauge theories

- 🔗 A serious **problem**: the **S** matrix **does not exist** in the usual Fock space
- 🔗 Possible **solutions**: construct finite transition probabilities (**KLN theorem**)
construct better asymptotic states (**coherent states**)
- 🔗 Long-distance singularities obey a pattern of **exponentiation**

$$\mathcal{M} = \mathcal{M}_0 \left[1 - \kappa \frac{\alpha}{\pi} \frac{1}{\epsilon} + \dots \right] \Rightarrow \mathcal{M} = \mathcal{M}_0 \exp \left[-\kappa \frac{\alpha}{\pi} \frac{1}{\epsilon} + \dots \right]$$

... and Practice

Just a **formal** issue in Quantum Field Theory? Are there **practical** applications?

🎤 **Higher order QCD calculations** at colliders hinge upon **cancellation of divergences** between virtual corrections and real emission contributions

- Cancellation must be performed analytically before numerical integrations
- Need local counterterms for matrix elements in all singular regions
- State of the art: NLO multileg, NNLO for (some) color-singlet processes

🎤 **Cancellations** leave behind **large logarithms**: they must be resummed

$$\underbrace{\frac{1}{\epsilon}}_{\text{virtual}} + \underbrace{(Q^2)^\epsilon \int_0^{m^2} \frac{dk^2}{(k^2)^{1+\epsilon}}}_{\text{real}} \implies \ln(m^2/Q^2)$$

- For inclusive observables: analytic resummation to high logarithmic accuracy.
- For exclusive final states: parton shower event generators, **(N)LL** accuracy.

🎤 **Resummation** probes the **all-order structure** of perturbation theory

- Power-suppressed corrections to QCD cross sections can be studied
- Links to the strong coupling regime can be established for SUSY gauge theories.

TOOLS



Dimensional regularization

Exponentiation of infrared poles requires solving **d-dimensional** evolution equations.

The running coupling in $d = 4 - 2\epsilon$ obeys

$$\mu \frac{\partial \bar{\alpha}}{\partial \mu} \equiv \beta(\epsilon, \bar{\alpha}) = -2\epsilon \bar{\alpha} + \hat{\beta}(\bar{\alpha}) \quad , \quad \hat{\beta}(\bar{\alpha}) = -\frac{\bar{\alpha}^2}{2\pi} \sum_{n=0}^{\infty} b_n \left(\frac{\bar{\alpha}}{\pi}\right)^n .$$


The **one-loop** solution is

$$\bar{\alpha}(\mu^2, \epsilon) = \alpha_s(\mu_0^2) \left[\left(\frac{\mu^2}{\mu_0^2}\right)^\epsilon - \frac{1}{\epsilon} \left(1 - \left(\frac{\mu^2}{\mu_0^2}\right)^\epsilon\right) \frac{b_0}{4\pi} \alpha_s(\mu_0^2) \right]^{-1} .$$

The β function develops an **IR-free fixed point**, so that the coupling **vanishes** at $\mu = 0$ for fixed $\epsilon < 0$. The **Landau pole** is at

$$\mu^2 = \Lambda^2 \equiv Q^2 \left(1 + \frac{4\pi\epsilon}{b_0\alpha_s(Q^2)}\right)^{-1/\epsilon} .$$

- ➔ Integrations over the scale of the coupling can be **analytically** performed.
- ➔ **All** infrared and collinear poles arise **by integration** over the scale of the running coupling.



For negative ϵ the beta function develops a second zero, $O(\epsilon)$ from the origin.

Factorization

All factorizations separating dynamics at different energy scales lead to **resummation** of logarithms of the ratio of scales.

A textbook example is **collinear factorization** for **DIS structure functions**.

- Collinear factorization separates the dependence on the **physical scale** Q^2 from the dependence on **collinear cutoffs** (parton masses m^2). For **Mellin moments** one gets


$$\tilde{F}_2 \left(N, \frac{Q^2}{m^2}, \alpha_s \right) = \tilde{C} \left(N, \frac{Q^2}{\mu_F^2}, \alpha_s \right) \tilde{f} \left(N, \frac{\mu_F^2}{m^2}, \alpha_s \right).$$

- Factorization requires the introduction of an **arbitrarily chosen** scale μ_F . Results must be **independent** of the arbitrary choice of μ_F .

$$\frac{d\tilde{F}_2}{d\mu_F} = 0 \quad \longrightarrow \quad \frac{d \log \tilde{f}}{d \log \mu_F} = \gamma_N(\alpha_s).$$

- The simple **functional dependence** of the factors is dictated by **separation of variables**.
- Proving **factorization** is the **difficult** step: it requires all-order diagrammatic analyses, or OPE. **Evolution** equations for parton distributions **follow** automatically.
- Solving Altarelli-Parisi evolution **resums** logarithms of Q^2/μ_F^2 into evolved **parton distributions** (or fragmentation functions).

Sudakov Factorization




Leading integration regions in loop momentum space for Sudakov factorization

- **Sudakov logarithms** are **remainders** of infrared and collinear **divergences**.
- **Divergences** arise in **scattering** amplitudes from **leading regions** in loop momentum space.
- **Soft gluons** factorize both from **hard** (easy) and from **collinear** (intricate) virtual exchanges.
- **Jet functions** **J** represent **color singlet** evolution of **external** hard partons.
- The **soft function** **S** is a **matrix** mixing the available **color representations**.
- In the **planar limit** soft exchanges are confined to **wedges**: **S** is proportional to the **identity**.
- In the **planar limit** **S** can be **reabsorbed** defining **jets** as square roots of **elementary form factors**.
- **Beyond** the planar limit **S** is determined by an **anomalous dimension matrix** Γ_S .

Color flow

In order to understand the **matrix structure** of the **soft function** it is sufficient to consider the simple case of **quark-antiquark** scattering.

At tree level



Tree-level diagrams and color flows for quark-antiquark scattering

For this process only **two color structures** are possible. A **basis** in the space of available color tensors is

$$c_{abcd}^{(1)} = \delta_{ab}\delta_{cd}, \quad c_{abcd}^{(2)} = \delta_{ac}\delta_{bd}$$


The **matrix element** is a **vector** in this space, and the Born cross section is

$$\mathcal{M}_{abcd} = \mathcal{M}_1 c_{abcd}^{(1)} + \mathcal{M}_2 c_{abcd}^{(2)} \longrightarrow \sum_{\text{color}} |\mathcal{M}|^2 = \sum_{J,L} \mathcal{M}_J \mathcal{M}_L^* \text{tr} \left[c_{abcd}^{(J)} \left(c_{abcd}^{(L)} \right)^\dagger \right] \equiv \text{Tr} [HS]_0$$

A virtual **soft gluon** will **reshuffle** color and mix the components of this vector

$$\text{QED} : \mathcal{M}_{\text{div}} = S_{\text{div}} \mathcal{M}_{\text{Born}} ; \quad \text{QCD} : [\mathcal{M}_{\text{div}}]_J = [S_{\text{div}}]_{JL} [\mathcal{M}_{\text{Born}}]_L$$

Sudakov factorization: pictorial



A pictorial representation of Sudakov factorization for fixed-angle scattering amplitudes

Operator Definitions

The precise **functional form** of this graphical factorization is

$$\mathcal{M}_L(p_i/\mu, \alpha_s(\mu^2), \epsilon) = \mathcal{S}_{LK}(\beta_i \cdot \beta_j, \alpha_s(\mu^2), \epsilon) H_K\left(\frac{p_i \cdot p_j}{\mu^2}, \frac{(p_i \cdot n_i)^2}{n_i^2 \mu^2}, \alpha_s(\mu^2)\right) \\ \times \prod_{i=1}^n \left[J_i\left(\frac{(p_i \cdot n_i)^2}{n_i^2 \mu^2}, \alpha_s(\mu^2), \epsilon\right) / \mathcal{J}_i\left(\frac{(\beta_i \cdot n_i)^2}{n_i^2}, \alpha_s(\mu^2), \epsilon\right) \right],$$

We introduced **factorization vectors** n_i^μ , $n_i^2 \neq 0$ to define the jets,

$$J\left(\frac{(p \cdot n)^2}{n^2 \mu^2}, \alpha_s(\mu^2), \epsilon\right) u(p) = \langle 0 | \Phi_n(\infty, 0) \psi(0) | p \rangle.$$

where Φ_n is the **Wilson line** operator along the direction n^μ ,

$$\Phi_n(\lambda_2, \lambda_1) = P \exp \left[ig \int_{\lambda_1}^{\lambda_2} d\lambda n \cdot A(\lambda n) \right].$$

- The vectors n^μ :
- Ensure **gauge invariance** of the jets.
 - **Separate** collinear gluons from wide-angle soft ones.
 - **Replace** other hard partons with a **collinear-safe** absorber.

Soft Matrices

The **soft function** \mathcal{S} is a **matrix**, mixing the available color tensors. It is defined by a correlator of **Wilson lines**.

$$(c_L)_{\{\alpha_k\}} \mathcal{S}_{LK}(\beta_i \cdot \beta_j, \alpha_s(\mu^2), \epsilon) = \sum_{\{\eta_k\}} \langle 0 | \prod_{i=1}^n \left[\Phi_{\beta_i}(\infty, 0)_{\alpha_k, \eta_k} \right] | 0 \rangle (c_K)_{\{\eta_k\}} ,$$

The soft function \mathcal{S} obeys a **matrix** RG evolution equation




$$\mu \frac{d}{d\mu} \mathcal{S}_{IK}(\beta_i \cdot \beta_j, \alpha_s(\mu^2), \epsilon) = - \mathcal{S}_{IJ}(\beta_i \cdot \beta_j, \alpha_s(\mu^2), \epsilon) \Gamma_{JK}^{\mathcal{S}}(\beta_i \cdot \beta_j, \alpha_s(\mu^2), \epsilon)$$

 $\Gamma^{\mathcal{S}}$ is **singular** due to overlapping **UV** and **collinear** poles.

\mathcal{S} is a **pure counterterm**. In dimensional regularization, using $\alpha_s(\mu^2 = 0, \epsilon < 0) = 0$,

$$\mathcal{S}(\beta_i \cdot \beta_j, \alpha_s(\mu^2), \epsilon) = P \exp \left[-\frac{1}{2} \int_0^{\mu^2} \frac{d\xi^2}{\xi^2} \Gamma^{\mathcal{S}}(\beta_i \cdot \beta_j, \alpha_s(\xi^2, \epsilon), \epsilon) \right] .$$

The determination of the **soft anomalous dimension matrix** $\Gamma^{\mathcal{S}}$ is the **keystone** of the resummation program for multiparton **amplitudes** and **cross sections**.


-  It **governs** the interplay of **color** exchange with **kinematics** in multiparton processes.
-  It is the only **source** of multiparton **correlations** for singular contributions.
-  **Collinear** effects are **'color singlet'** and can be extracted from **two-parton** scatterings.

THE DIPOLE FORMULA



Surprising Simplicity

- The matrix Γ_S can be computed from the **UV poles** of S .
- Computations** can be performed directly **for the exponent**: the relevant diagrams are called “**webs**”.
- Γ_S appears **highly complex** at high orders.
- g-loop** webs directly **correlate** color and kinematics of up to **g+1** Wilson lines.



A web contributing to the soft anomalous dimension matrix

The **two-loop** calculation (Aybat, Dixon, Sterman 06) leads to a **surprising result**: for **any number** of external **massless** partons

$$\Gamma_S^{(2)} = \frac{\kappa}{2} \Gamma_S^{(1)} \quad \kappa = \left(\frac{67}{18} - \zeta(2) \right) C_A - \frac{10}{9} T_F C_F .$$

- ➔ **No** new kinematic dependence; **no** new matrix structure.
- ➔ κ is the two-loop coefficient of $\gamma_K(\alpha_s)$, rescaled by the appropriate **quadratic Casimir**,

$$\gamma_K^{(i)}(\alpha_s) = C^{(i)} \left[2 \frac{\alpha_s}{\pi} + \kappa \left(\frac{\alpha_s}{\pi} \right)^2 + \mathcal{O}(\alpha_s^3) \right] .$$

The Dipole Formula

The two-loop result led to an **all-order understanding**. For **massless** partons, the soft matrix obeys a set of **exact equations** that **correlate color** exchange with **kinematics**.

The **simplest solution** to these equations is a **sum over color dipoles** (Becher, Neubert; Gardi, LM, 09). It leads to an **ansatz** for the all-order singularity structure of **all** multiparton fixed-angle **massless** scattering amplitudes: the **dipole formula**.

🔗 All **soft** and **collinear** singularities can be **collected** in a multiplicative operator **Z**

$$\mathcal{M} \left(\frac{p_i}{\mu}, \alpha_s(\mu^2), \epsilon \right) = Z \left(\frac{p_i}{\mu_f}, \alpha_s(\mu_f^2), \epsilon \right) \mathcal{H} \left(\frac{p_i}{\mu}, \frac{\mu_f}{\mu}, \alpha_s(\mu^2), \epsilon \right),$$

🔗 **Z** contains both soft singularities from **S**, and collinear ones from the jet functions. It must **satisfy** its own matrix **RG equation**

$$\frac{d}{d \ln \mu} Z \left(\frac{p_i}{\mu}, \alpha_s(\mu^2), \epsilon \right) = - Z \left(\frac{p_i}{\mu}, \alpha_s(\mu^2), \epsilon \right) \Gamma \left(\frac{p_i}{\mu}, \alpha_s(\mu^2) \right).$$

The matrix **Γ** **inherits** the **dipole structure** from the soft matrix. It reads

$$\Gamma_{\text{dip}} \left(\frac{p_i}{\mu}, \alpha_s(\mu^2) \right) = -\frac{1}{4} \hat{\gamma}_K(\alpha_s(\mu^2)) \sum_{j \neq i} \ln \left(\frac{-2 p_i \cdot p_j}{\mu^2} \right) \mathbf{T}_i \cdot \mathbf{T}_j + \sum_{i=1}^n \gamma_{J_i}(\alpha_s(\mu^2)).$$

Note that **all singularities** are **generated by integration** over the scale of the coupling.

Features of the dipole formula

- All known results for IR divergences of massless gauge theory amplitudes are recovered.
- The absence of multiparton correlations implies remarkable diagrammatic cancellations.
- The color matrix structure is fixed at one loop: path-ordering is not needed.
- All divergences are determined by a handful of anomalous dimensions.
- The cusp anomalous dimension plays a very special role: a universal IR coupling.

Can this be the definitive answer for IR divergences in massless non-abelian gauge theories?

► There are precisely two sources of possible corrections.

- Quadrupole correlations may enter starting at three loops: they must be tightly constrained functions of conformal cross ratios of parton momenta.


$$\Gamma\left(\frac{p_i}{\mu}, \alpha_s(\mu^2)\right) = \Gamma_{\text{dip}}\left(\frac{p_i}{\mu}, \alpha_s(\mu^2)\right) + \Delta(\rho_{ijkl}, \alpha_s(\mu^2)) \quad , \quad \rho_{ijkl} = \frac{p_i \cdot p_j \, p_k \cdot p_l}{p_i \cdot p_k \, p_j \cdot p_l}$$

- The cusp anomalous dimension may violate Casimir scaling beyond three loops.

$$\gamma_K^{(i)}(\alpha_s) = C_i \hat{\gamma}_K(\alpha_s) + \tilde{\gamma}_K^{(i)}(\alpha_s)$$


- The functional form of Δ is further constrained by: collinear limits, Bose symmetry and transcendentality bounds (Becher, Neubert; Dixon, Gardi, LM, 09).
- A four-loop analysis indicates that Casimir scaling holds (Becher, Neubert, Vernazza).

THE HIGH-ENERGY LIMIT



Reggeization

- Studies of the **high-energy** limit **predate** the modern era of quantum field theory.
- In the $t/s \rightarrow 0$ limit **particles** exchanged in the **t-channel** (may) **'Reggeize'**.



Gluon-gluon scattering: the t-channel gluon Reggeizes

- **Large logarithms** of s/t are **generated** by a simple replacement of the **t-channel propagator**,

$$\frac{1}{t} \longrightarrow \frac{1}{t} \left(\frac{s}{-t} \right)^{\alpha(t)}$$

- The **Regge trajectory** has a perturbative expansion, with **IR divergent** coefficients

$$\alpha(t) = \frac{\alpha_s(-t, \epsilon)}{4\pi} \alpha^{(1)} + \left(\frac{\alpha_s(-t, \epsilon)}{4\pi} \right)^2 \alpha^{(2)} + \mathcal{O}(\alpha_s^3)$$

- The **gluon** has been shown to **Reggeize** at **NLL**, and the **two-loop** Regge trajectory is known.
- For example, for **gluon-gluon scattering** the matrix element obeys **Regge factorization**


$$\mathcal{M}_{a_1 a_2 a_3 a_4}^{gg \rightarrow gg}(s, t) = 2 g_s^2 \frac{s}{t} \left[(T^b)_{a_1 a_3} C_{\lambda_1 \lambda_3}(k_1, k_3) \right] \left(\frac{s}{-t} \right)^{\alpha(t)} \left[(T_b)_{a_2 a_4} C_{\lambda_2 \lambda_4}(k_2, k_4) \right]$$

with the perturbative coefficients

$$\alpha^{(1)} = C_A \frac{\widehat{\gamma}_K^{(1)}}{\epsilon} = C_A \frac{2}{\epsilon} \quad \alpha^{(2)} = C_A \left[-\frac{b_0}{\epsilon^2} + \widehat{\gamma}_K^{(2)} \frac{2}{\epsilon} + C_A \left(\frac{404}{27} - 2\zeta_3 \right) + n_f \left(-\frac{56}{27} \right) \right]$$

Multi-Regge kinematics

- Reggeization follows from the dominance of **t**-channel **ladder** diagrams as $t/s \rightarrow 0$.
- By **unitarity**, **multi-gluon** emission must similarly **simplify** in the high-energy limit.
- Regge **factorization** extends to **multi-particle** emission in 'Multi-Regge' kinematics.



$$y_3 \gg y_4 \gg \dots \gg y_L, \quad |k_i^\perp| \simeq |k_j^\perp|, \quad \forall i, j$$

$$-s \equiv -s_{12} \simeq |k_3^\perp| |k_L^\perp| e^{y_3 - y_L} e^{i\pi}$$

$$-s_{ij} \simeq |k_i^\perp| |k_j^\perp| e^{y_i - y_j} e^{i\pi}, \quad 3 \leq i < j \leq L$$

Multi-gluon emission and Multi-Regge kinematics

- Large logarithms** of s/t_i are generated by the **Reggeization** of **t**-channel propagators, as

$$\begin{aligned} \mathcal{M}_{a_1 \dots a_L}^{gg \rightarrow (L-2)g} &= 2 g_s^3 s \left[(T^b)_{a_1 a_3} C_{\lambda_1 \lambda_3}(k_1, k_3) \right] \left[\frac{1}{t_1} \left(\frac{s_{34}}{-t_1} \right)^{\alpha(t_1)} \right] \\ &\times \left[(T^{a_4})_{bc} V_{\lambda_4}(q_1, q_2) \right] \left[\frac{1}{t_2} \left(\frac{s_{45}}{-t_2} \right)^{\alpha(t_2)} \right] \dots \left[(T^c)_{a_2 a_L} C_{\lambda_2 \lambda_L}(k_2, k_L) \right] \end{aligned}$$

- The **impact factors** **C** and the **Lipatov vertices** **V** are **universal** and independent of s .

The dipole formula at high energy

Introducing **Mandelstam color** operators, and using **color** and **momentum** conservation

$$\begin{aligned} \mathbf{T}_s &= \mathbf{T}_1 + \mathbf{T}_2 = -(\mathbf{T}_3 + \mathbf{T}_4), & s + t + u &= 0 \\ \mathbf{T}_t &= \mathbf{T}_1 + \mathbf{T}_3 = -(\mathbf{T}_2 + \mathbf{T}_4), & \mathbf{T}_s^2 + \mathbf{T}_t^2 + \mathbf{T}_u^2 &= \sum_{i=1}^4 C_i \\ \mathbf{T}_u &= \mathbf{T}_1 + \mathbf{T}_4 = -(\mathbf{T}_2 + \mathbf{T}_3) \end{aligned}$$

it is easy to see that the infrared **dipole operator Z factorizes** in the high-energy limit

$$Z\left(\frac{p_i}{\mu}, \alpha_s(\mu^2), \epsilon\right) = \tilde{Z}\left(\frac{s}{t}, \alpha_s(\mu^2), \epsilon\right) Z_1\left(\frac{t}{\mu^2}, \alpha_s(\mu^2), \epsilon\right)$$

- The operator Z_1 is **s-independent** and proportional to the **unit matrix** in color space.
- **Color** dependence and **s** dependence are **collected** in the factor

$$\tilde{Z}\left(\frac{s}{t}, \alpha_s(\mu^2), \epsilon\right) = \exp\left\{K\left(\alpha_s(\mu^2), \epsilon\right) \left[\ln\left(\frac{s}{-t}\right) \mathbf{T}_t^2 + i\pi \mathbf{T}_s^2 \right]\right\},$$

where the **coupling** dependence is (once again!) completely **determined** by the **cusp** anomalous dimension and by the **β function**, through

$$K\left(\alpha_s(\mu^2), \epsilon\right) \equiv -\frac{1}{4} \int_0^{\mu^2} \frac{d\lambda^2}{\lambda^2} \hat{\gamma}_K\left(\alpha_s(\lambda^2), \epsilon\right)$$

The **simple structure** of the high-energy operator **governs** Reggeization and its breaking.

Reggeization of leading logarithms

- At **leading logarithmic** accuracy, the (**imaginary**) **s**-channel contribution can be **dropped**, and the dipole operator becomes **diagonal** in a **t**-channel basis.


$$\mathcal{M} \left(\frac{p_i}{\mu}, \alpha_s(\mu^2), \epsilon \right) = \exp \left\{ K \left(\alpha_s(\mu^2), \epsilon \right) \ln \left(\frac{s}{-t} \right) \mathbf{T}_t^2 \right\} Z_1 \mathcal{H} \left(\frac{p_i}{\mu}, \alpha_s(\mu^2), \epsilon \right)$$

- If, at **LO** and at **leading power** in **t/s**, the scattering is **dominated** by **t**-channel exchange, then the **hard function** is an **eigenstate** of the color operator \mathbf{T}_t^2

$$\mathbf{T}_t^2 \mathcal{H}^{gg \rightarrow gg} \xrightarrow{|t/s| \rightarrow 0} C_t \mathcal{H}_t^{gg \rightarrow gg}$$

- Leading-logarithmic **Reggeization** for **arbitrary t**-channel color representations **follows**

$$\mathcal{M}^{gg \rightarrow gg} = \left(\frac{s}{-t} \right)^{C_A K(\alpha_s(\mu^2), \epsilon)} Z_1 \mathcal{H}_t^{gg \rightarrow gg}$$



- The **LL Regge trajectory** is **universal** and obeys Casimir scaling.
- Scattering of **arbitrary color representations** can be **analyzed**
Example: let **1** and **2** be **antiquarks**, **4** a **gluon** and **3** a **sextet**; use

$$\bar{\mathbf{3}} \otimes \mathbf{6} = \mathbf{3} \oplus \mathbf{15}$$

$$\bar{\mathbf{3}} \otimes \mathbf{8}_a = \bar{\mathbf{3}} \oplus \mathbf{6} \oplus \bar{\mathbf{15}}$$

LL Reggeization of the **3** and **15** **t**-channel exchanges **follows**.

Beyond leading logarithms

- The **high-energy** infrared **operator** can be **systematically expanded** beyond **LL**, using the **Baker-Campbell-Hausdorff** formula. At **NLL** one finds a series of commutators

$$\tilde{Z}\left(\frac{s}{t}, \alpha_s, \epsilon\right)\Big|_{\text{NLL}} = \left(\frac{s}{-t}\right)^{K(\alpha_s, \epsilon) \mathbf{T}_t^2} \left\{ 1 + i\pi K(\alpha_s, \epsilon) \left[\mathbf{T}_s^2 - \frac{K(\alpha_s, \epsilon)}{2!} \ln\left(\frac{s}{-t}\right) [\mathbf{T}_t^2, \mathbf{T}_s^2] + \frac{K^2(\alpha_s, \epsilon)}{3!} \ln^2\left(\frac{s}{-t}\right) [\mathbf{T}_t^2, [\mathbf{T}_t^2, \mathbf{T}_s^2]] + \dots \right] \right\}$$

- The **real part** of the amplitude **Reggeizes** also at **NLL** for **arbitrary t**-channel exchanges.

- At **NNLL** **Reggeization** generically **breaks down** also for the **real part** of the amplitude.

- At **two loops**, terms that are **non-logarithmic** and **non-diagonal** in a **t**-channel basis arise

$$\mathcal{E}_0(\alpha_s, \epsilon) \equiv -\frac{1}{2}\pi^2 K^2(\alpha_s, \epsilon) (\mathbf{T}_s^2)^2$$

- At **three loops**, the first Reggeization-breaking **logarithms** of **s/t** arise, generated by

$$\mathcal{E}_1\left(\frac{s}{t}, \alpha_s, \epsilon\right) \equiv -\frac{\pi^2}{3} K^3(\alpha_s, \epsilon) \ln\left(\frac{s}{-t}\right) [\mathbf{T}_s^2, [\mathbf{T}_t^2, \mathbf{T}_s^2]]$$

- NOTE**
 - In the **planar limit** ($N_c \rightarrow \infty$) **all commutators vanish** and Reggeization **holds** also **beyond NLL** (as perhaps expected from **string theory**).
 - Possible **quadrupole corrections** to the dipole formula **cannot** come to the rescue.


Multi-Regge kinematics

- The **dipole formula** applies for **any number** of particles: we expect similar **simplifications** in **Multi-Regge** kinematics, and similar **results** concerning **Reggeization**.
- Indeed, one can **prove** recursively that the dipole operator **Z factorizes** in **MR** kinematics, as

$$Z\left(\frac{p_l}{\mu}, \alpha_s(\mu^2), \epsilon\right) = \tilde{Z}^{\text{MR}}\left(\Delta y_k, \alpha_s(\mu^2), \epsilon\right) Z_1^{\text{MR}}\left(\frac{|k_i^\perp|}{\mu}, \alpha_s(\mu^2), \epsilon\right)$$

- The **Multi-Regge** high-energy **operator** has again a simple structure.

$$\tilde{Z}^{\text{MR}}\left(\Delta y_k, \alpha_s(\mu^2), \epsilon\right) = \exp\left\{K\left(\alpha_s(\mu^2), \epsilon\right) \left[\sum_{k=3}^{L-1} \mathbf{T}_{t_{k-2}}^2 \Delta y_k + i\pi \mathbf{T}_s^2\right]\right\}$$



Color structure in Multi-Regge kinematics

- We have **defined** the **t-channel color operators**

$$\mathbf{T}_{t_k} = \mathbf{T}_1 + \sum_{p=1}^k \mathbf{T}_{p+2}$$


- A **t-channel basis** of **common** eigenstates of \mathbf{T}_{t_k} can be **constructed** using **Clebsch-Gordan** coefficients.
- The operators \mathbf{T}_{t_k} thus **commute**, and **each color representation** contributing to the hard function in the high-energy limit **Reggeizes separately** at **LL**.

Constraining quadrupoles

- Known results on the high-energy limit of QCD amplitudes imply **new constraints** on **quadrupole** corrections to the **dipole formula** at three loops and beyond.
- Previous analyses using **collinear** constraint, **Bose** symmetry and **transcendentality** bounds **could not exclude** a class of correction, including for example

$$\Delta^{(212)}(\rho_{ijkl}, \alpha_s) = \left(\frac{\alpha_s}{\pi}\right)^3 \mathbf{T}_1^a \mathbf{T}_2^b \mathbf{T}_3^c \mathbf{T}_4^d \left[f^{ade} f^{cbe} L_{1234}^2 \left(L_{1423} L_{1342}^2 + L_{1423}^2 L_{1342} \right) + \text{cycl.} \right]$$

where $L_{ijkl} = \log(\rho_{ijkl})$.



A three-loop diagram for Δ

- In the **high-energy limit** one finds (with $L = \log(s/t)$)

$$\begin{aligned} \rho_{1234} &\equiv \frac{(-s_{12})(-s_{34})}{(-s_{13})(-s_{24})} = \left(\frac{s}{-t}\right)^2 e^{-2i\pi}; & L_{1234} &= 2(L - i\pi); \\ \rho_{1342} &\equiv \frac{(-s_{13})(-s_{24})}{(-s_{14})(-s_{23})} = \left(\frac{-t}{s+t}\right)^2; & L_{1342} &\simeq -2L; \\ \rho_{1423} &\equiv \frac{(-s_{14})(-s_{23})}{(-s_{12})(-s_{34})} = \left(\frac{s+t}{s}\right)^2 e^{2i\pi}; & L_{1423} &\simeq 2i\pi, \end{aligned}$$

- Previously **admissible** corrections display **superleading high-energy logarithms** at three loops.

$$\Delta^{(212)}(\rho_{ijkl}, \alpha_s) = \left(\frac{\alpha_s}{\pi}\right)^3 \mathbf{T}_1^a \mathbf{T}_2^b \mathbf{T}_3^c \mathbf{T}_4^d 32 i \pi \left[(-L^4 - i\pi L^3 - \pi^2 L^2 - i\pi^3 L) f^{ade} f^{cbe} + \dots \right]$$

- No known explicit example** of admissible quadrupole correction **survives**. A complete **proof** is **still lacking**: linear combinations might restore the proper Regge behavior.

OUTLOOK



Summary

- After $7.5 \cdot 10^2$ years, **soft** and **collinear singularities** in gauge theories amplitudes are still a **fertile** field of study. A **definitive solution** may be at hand.
 - ✓ We are probing the **all-order** structure of the nonabelian **exponent**.
 - ✓ **All-order** results constrain, test and complement **fixed-order** calculations.
 - ✓ Understanding singularities has **phenomenological applications** through **resummation**.
- Factorization** theorems \Rightarrow **Evolution** equations \Rightarrow **Exponentiation**.
 - ✓ Sudakov **factorization** \Rightarrow soft-gluon **resummation**.
 - ✓ Multiparton processes require **anomalous dimension matrices**.
- A simple **dipole formula** may encode **all infrared singularities** for **any massless gauge** theory, a **natural generalization** of the planar limit. The study of possible **corrections** to the dipole formula is **under way**.
- The **high-energy limit** of the dipole formula provides **insights** into **Reggeization** and **beyond**, at least for **divergent contributions** to the amplitude.
- Leading logarithmic **Reggeization** is **proved** for **generic** color **representations** exchanged in the **t** channel, and for **any number of partons** in Multi-Regge kinematics.
- Regge factorization** generically **breaks down** at **NNLL**, with **computable** corrections.
- The **high-energy limit** further **constrains** quadrupole **corrections** to the dipole formula: **no** known **examples** survive.

$$\Gamma_{\text{dip}} \left(\frac{p_i}{\mu}, \alpha_s(\mu^2) \right) = -\frac{1}{4} \hat{\gamma}_K(\alpha_s(\mu^2)) \sum_{j \neq i} \ln \left(\frac{-2 p_i \cdot p_j}{\mu^2} \right) \mathbf{T}_i \cdot \mathbf{T}_j + \sum_{i=1}^n \gamma_{J_i}(\alpha_s(\mu^2)) .$$

INFRARED



CATASTROPHE

THANK YOU!