

# Automated one-loop calculations with Golem/Samurai

Gudrun Heinrich



Max-Planck Institute for Physics, Munich





RADCOR conference  
Mamallapuram, India, 27.09.2011



# Motivation



# Motivation



we need precise theory predictions for "Expected" !


# N(N)LO

predictions beyond leading order:

- ▶ reduction of scale dependence
- ▶ reliable estimate of absolute rates
- ▶ better description of jets
- ▶ better PDF fits
- ▶ ...

# N(N)LO

## 2jet to 3jet transition parameter Y3



example:

3-jet observable  
in  $e^+e^-$  annihilation

[A. Gehrmann-De Ridder,  
T. Gehrmann, N. Glover, GH '09]

uncertainty bands:  
 $M_Z/2 < \mu < 2 M_Z$

## some NLO multi-leg highlights

- ▶  $pp \rightarrow W^+ W^- b\bar{b}$  Denner, Dittmaier, Kallweit, Pozzorini '10 ; Bevilacqua, Czakon, van Hameren, Papadopoulos, Worek '11
- ▶  $pp \rightarrow W/Z + 4$  jets BlackHat collaboration '10/'11
- ▶  $pp \rightarrow W/Z/\gamma + 3$  jets BlackHat collaboration '09/'10
- ▶  $pp \rightarrow t\bar{t} + 2$  jets Bevilacqua, Czakon, Papadopoul., Worek '10
- ▶  $pp \rightarrow t\bar{t} b\bar{b}$  Bredenstein, Denner, Dittmaier, Pozzorini '09; Bevilacqua, Czakon, Papadopoulos, Worek '09
- ▶  $pp \rightarrow W\gamma\gamma j$  Campanario, Englert, Rauch, Zeppenfeld '11
- ▶  $pp \rightarrow W^+ W^+ jj$  Melia, Melnikov, Rontsch, Zanderighi '10
- ▶  $pp \rightarrow W^+ W^- jj$  Melia, Melnikov, Rontsch, Zanderighi '11
- ▶  $pp \rightarrow 4 b$  Binoth et al '09; Greiner, Guffanti, Reiter, Reuter '11
- ▶ NGluon ( $N < \sim 14$ ) Badger, Biedermann, Uwer '11 (public)
- ▶  $e^+ e^- \rightarrow 5$  jets Frederix, Frixione, Melnikov, Zanderighi '10

## some NLO multi-leg highlights

- ▶  $pp \rightarrow W^+ W^- b\bar{b}$  Denner, Dittmaier, Kallweit, Pozzorini '10 ; Bevilacqua, Czakon, van Hameren, Papadopoulos, Worek '11
- ▶  $pp \rightarrow W/Z + 4$  jets BlackHat collaboration '10/'11
- ▶  $pp \rightarrow W/Z/\gamma + 3$  jets BlackHat collaboration '09/'10
- ▶  $pp \rightarrow t\bar{t} + 2$  jets Bevilacqua, Czakon, Papadopoul., Worek '10
- ▶  $pp \rightarrow t\bar{t} b\bar{b}$  Bredenstein, Denner, Dittmaier, Pozzorini '09; Bevilacqua, Czakon, Papadopoulos, Worek '09
- ▶  $pp \rightarrow W\gamma\gamma j$  Campanario, Englert, Rauch, Zeppenfeld '11
- ▶  $pp \rightarrow W^+ W^+ jj$  Melia, Melnikov, Rontsch, Zanderighi '10
- ▶  $pp \rightarrow W^+ W^- jj$  Melia, Melnikov, Rontsch, Zanderighi '11
- ▶  $pp \rightarrow 4 b$  Binoth et al '09; Greiner, Guffanti, Reiter, Reuter '11
- ▶ NGluon ( $N < \sim 14$ ) Badger, Biedermann, Uwer '11 (public)
- ▶  $e^+ e^- \rightarrow 5$  jets Frederix, Frixione, Melnikov, Zanderighi '10
- ▶ also: BIG advances in automation

# Automated NLO Tools

## One-loop

- ▶ FeynArts/FormCalc/LoopTools ([public](#)) Thomas Hahn et al
- ▶ Helac-NLO Bevilacqua, Czakon, van Hameren, Papadopoulos, Pittau, Worek
- ▶ MadLoop/ aMC@NLO Hirschi, Frederix, Frixione, Garzelli, Maltoni, Pittau  
uses **CutTools** ([public](#)) [Ossola, Papadopoulos, Pittau] and **MadFKS**
- ▶ Golem-Samurai Cullen, Greiner, GH, Luisoni, Mastrolia, Ossola, Reiter, Tramontano  
**Samurai** ([public](#)) [Mastrolia, Ossola, Reiter, Tramontano],  
**golem95** ([public](#)) [Binoth, Cullen, Guillet, GH, Kleinschmidt, Pilon, Reiter, Rodgers]
- ▶ dedicated programs also involve high level of automation

Denner, Dittmaier, Pozzorini et al, VBFNLO coll., Blackhat, Rocket, MCFM, ...

# Automated NLO Tools

## One-loop

- ▶ FeynArts/FormCalc/LoopTools ([public](#)) Thomas Hahn et al
- ▶ Helac-NLO Bevilacqua, Czakon, van Hameren, Papadopoulos, Pittau, Worek
- ▶ MadLoop/ aMC@NLO Hirschi, Frederix, Frixione, Garzelli, Maltoni, Pittau  
uses **CutTools** ([public](#)) [Ossola, Papadopoulos, Pittau] and **MadFKS**
- ▶ Golem-Samurai Cullen, Greiner, GH, Luisoni, Mastrolia, Ossola, Reiter, Tramontano  
**Samurai** ([public](#)) [Mastrolia, Ossola, Reiter, Tramontano],  
**golem95** ([public](#)) [Binoth, Cullen, Guillet, GH, Kleinschmidt, Pilon, Reiter, Rodgers]
- ▶ dedicated programs also involve high level of automation  
Denner, Dittmaier, Pozzorini et al, VBFNLO coll., Blackhat, Rocket, MCFM, ...

## automation of subtraction for IR divergent real radiation

- ▶ MadDipole Frederix, Greiner, Gehrmann 08
- ▶ Dipole subtraction in Sherpa Gleisberg, Krauss 08
- ▶ TevJet Seymour, Tevlin 08
- ▶ AutoDipole Hasegawa, Moch, Uwer 08,09
- ▶ Helac-Phegas Czakon, Papadopoulos, Worek 09; polarized
- ▶ MadFKS Frederix, Frixione, Maltoni, Stelzer 09

# Golem-Samurai (GoSam)



# Golem-Samurai (GoSam)

General One-Loop Evaluator of Matrix elements &  
Scattering Amplitudes from Unitarity based Reduction At Integrand level  
[ Cullen, Greiner, GH, Luisoni, Mastrolia, Ossola, Reiter, Tramontano ]

- ▶ algebraic generation of D-dimensional integrands based on Feynman diagrams
  - ▶ QCD, EW, BSM
  - ▶ uses QGraf [Nogueira], FeynRules [Duhr et al], Form/Spinney [Vermaseren/Cullen et al], Haggies [Reiter] to generate integrands

# Golem-Samurai (GoSam)

General One-Loop Evaluator of Matrix elements &  
Scattering Amplitudes from Unitarity based Reduction At Integrand level  
[ Cullen, Greiner, GH, Luisoni, Mastrolia, Ossola, Reiter, Tramontano ]


- ▶ algebraic generation of D-dimensional integrands based on Feynman diagrams
  - ▶ QCD, EW, BSM
  - ▶ uses QGraf [Nogueira], FeynRules [Duhr et al], Form/Spinney [Vermaseren/Cullen et al], Haggies [Reiter] to generate integrands
- ▶ reduction by D-dimensional extension of cut-based method options:
  - OPP-type reduction  
[Ossola, Papadopoulos, Pittau; Ellis, Giele, Kunszt, Melnikov]
  - traditional tensor reduction (using golem95 library)
  - tensorial reduction at integrand level GH, Ossola, Reiter, Tramontano '10

# Golem-Samurai (GoSam)

General One-Loop Evaluator of Matrix elements &  
Scattering Amplitudes from Unitarity based Reduction At Integrand level  
[ Cullen, Greiner, GH, Luisoni, Mastrolia, Ossola, Reiter, Tramontano ]

- ▶ algebraic generation of D-dimensional integrands based on Feynman diagrams
  - ▶ QCD, EW, BSM
  - ▶ uses QGraf [Nogueira], FeynRules [Duhr et al], Form/Spinney [Vermaseren/Cullen et al], Haggies [Reiter] to generate integrands
- ▶ reduction by D-dimensional extension of cut-based method options:
  - OPP-type reduction  
[Ossola, Papadopoulos, Pittau; Ellis, Giele, Kunszt, Melnikov]
  - traditional tensor reduction (using golem95 library)
  - tensorial reduction at integrand level GH, Ossola, Reiter, Tramontano '10
- ▶ interface with existing tools for real radiation  
(MadGraph/MadEvent, Sherpa, Powheg, ... )

# Golem-Samurai structure



# Golem-Samurai

usage:

- ▶ edit "input card"

```
in= u,d~  
out= nmu, mu+, e-, ne~, s~, c  
model=smdiag  
models can be added via FeynRules (Duhr) or LanHEP (Semenov)  
order=gw,4,4; order=gs,2,4  
zero=mB,mC,mS,mU,mD,me,mmu  
one=gs,e  
helicities=-+-+-+-  
extensions=samurai, dred
```

- ▶ golem-main.py process.in

# Golem-Samurai

usage:

- ▶ edit "input card"

```
in= u,d~  
out= nmu, mu+, e-, ne~, s~, c  
model=smdiag  
models can be added via FeynRules (Duhr) or LanHEP (Semenov)  
order=gw,4,4; order=gs,2,4  
zero=mB,mC,mS,mU,mD,me,mmu  
one=gs,e  
helicities=-+-+-+-  
extensions=samurai, dred
```

- ▶ golem-main.py process.in
- ▶ make doc ⇒ documentation and diagram pictures

# Golem-Samurai

usage:

- ▶ edit "input card"

```
in= u,d~  
out= nmu, mu+, e-, ne~, s~, c  
model=smdiag  
models can be added via FeynRules (Duhr) or LanHEP (Semenov)  
order=gw,4,4; order=gs,2,4  
zero=mB,mC,mS,mU,mD,me,mmu  
one=gs,e  
helicities=-+-+-+-  
extensions=samurai, dred
```

- ▶ golem-main.py process.in
- ▶ make doc ⇒ documentation and diagram pictures
- ▶ make source ⇒ source files

# Golem-Samurai



usage:

- ▶ edit "input card"

```
in= u,d~  
out= nmu, mu+, e-, ne~, s~, c  
model=smdiag  
models can be added via FeynRules (Duhr) or LanHEP (Semenov)  
order=gw,4,4; order=gs,2,4  
zero=mB,mC,mS,mU,mD,me,mmu  
one=gs,e  
helicities=-+-+-+  
extensions=samurai, dred
```

- ▶ golem-main.py process.in
- ▶ make doc ⇒ documentation and diagram pictures
- ▶ make source ⇒ source files
- ▶ make compile ⇒ fully compiled code

Example  $u \bar{d} \rightarrow W^- W^+ \bar{s} c \rightarrow e^- \bar{\nu}_e \mu^+ \nu_\mu \bar{s} c$




## 5 One-Loop Diagrams


### General Information

Example  $u \bar{d} \rightarrow W^- W^+ \bar{s} c \rightarrow e^- \bar{\nu}_e \mu^+ \nu_\mu \bar{s} c$

NLO sample diagrams




$$S' = S_{Q \rightarrow -q - (k_3 - k_2 + k_4)}^{\{4\}}, \text{ rk } = 3$$




$$S' = S_{Q \rightarrow q + (k_1)}^{\{4\}}, \text{ rk } = 4$$

Example  $u \bar{d} \rightarrow W^- W^+ \bar{s} c \rightarrow e^- \bar{\nu}_e \mu^+ \nu_\mu \bar{s} c$




$$S' = S_{Q \rightarrow -q-(k1)}^{\{1\}}, \text{ rk } = 3$$

$$S' = S_{Q \rightarrow q+(k1)}^{\{3\}}, \text{ rk } = 3$$



$$S' = S_{Q \rightarrow q+(k1)}^{\{1,3\}}, \text{ rk } = 2$$



$$S' = S_{Q \rightarrow -q}^{\{1,4\}}, \text{ rk } = 2$$

Example  $u \bar{d} \rightarrow W^- W^+ \bar{s}c \rightarrow e^- \bar{\nu}_e \mu^+ \nu_\mu \bar{s}c$

code generation:

```
Form is processing loop diagram 80 @ Helicity 0
  2.72 sec out of 2.74 sec
Haggies is processing abbreviations for loop diagram 80 @ Helicity 0
Form is processing loop diagram 81 @ Helicity 0
  0.71 sec out of 0.73 sec
Haggies is processing abbreviations for loop diagram 81 @ Helicity 0
Form is processing loop diagram 82 @ Helicity 0
  0.73 sec out of 0.75 sec
Haggies is processing abbreviations for loop diagram 82 @ Helicity 0
Form is processing loop diagram 83 @ Helicity 0
  0.70 sec out of 0.71 sec
Haggies is processing abbreviations for loop diagram 83 @ Helicity 0
Form is processing loop diagram 84 @ Helicity 0
  0.73 sec out of 0.73 sec
Haggies is processing abbreviations for loop diagram 84 @ Helicity 0
Form is processing loop diagram 85 @ Helicity 0
```

Example  $u \bar{d} \rightarrow W^- W^+ \bar{s}c \rightarrow e^- \bar{\nu}_e \mu^+ \nu_\mu \bar{s}c$

```
=====
          GoSam-1.0
=====

#   NLO/LO, finite part: -15.91575118714612
#   NLO/LO, single pole:  7.587050495888512
#   NLO/LO, double pole: -5.333333333333234

CPU time (secs): 1.2997999999999991E-002
```

result compared with

Melia, Melnikov, Rontsch, Zanderighi (MMRZ) 1104.2327 [hep-ph]



NLO/LO	GoSam	MMRZ
$1/\epsilon^2$	-5.333333333	-5.333333
$1/\epsilon$	7.5870504959	7.587051
finite	-15.915751119	-15.91575

## Tested 5- or 6-point processes


- ▶  $u \bar{d} \rightarrow W^+ s\bar{s} \rightarrow e^+ \nu_e s\bar{s}$
- ▶  $u \bar{d} \rightarrow W^+ gg \rightarrow e^+ \nu_e gg$
- ▶  $d \bar{d} \rightarrow Z gg \rightarrow e^+ e^- gg$
- ▶  $u \bar{d} \rightarrow W^+ b\bar{b} \rightarrow e^+ \nu_e b\bar{b}$  also with massive b's
- ▶  $\gamma\gamma \rightarrow \gamma\gamma\gamma\gamma$
- ▶  $q \bar{q} \rightarrow b\bar{b} b\bar{b}$
- ▶  $gg \rightarrow b\bar{b} b\bar{b}$
- ▶  $u \bar{d} \rightarrow W^+ W^+ s\bar{c} \rightarrow e^+ \nu_e \mu^+ \nu_\mu s\bar{c}$
- ▶  $u \bar{u} \rightarrow W^+ W^- \bar{c}c \rightarrow e^- \bar{\nu}_e \mu^+ \nu_\mu \bar{c}c$
- ▶  $u \bar{d} \rightarrow W^+ W^- s\bar{c} \rightarrow e^- \bar{\nu}_e \mu^+ \nu_\mu \bar{s}c$
- ▶  $u \bar{d} \rightarrow W^+ g \rightarrow e^+ \nu_e g$  EW corrections
- ▶ plus a large number of  $2 \rightarrow 2$  processes

# golem95 integral library

**Example:** production of a heavy neutral MSSM Higgs and a  $b\bar{b}$  pair with unstable particles (squarks, neutralinos) in the loop



real masses



complex masses

contained in **golem95C library**: 1101.5595 [hep-ph]

Binoth, Cullen, Guillet, GH, Kleinschmidt, Pilon, Reiter, Rodgers

<http://projects.hepforge.org/~golem/95/>

# Numerical stability

several detection and rescue systems

- ▶ local/global  $N = N_{\text{rec}}$  test: use decomposition of numerator function after coefficients have been determined:

$$\begin{aligned}N(\bar{q}) &= \sum_{i << m}^{n-1} \Delta_{ijklm}(\bar{q}) \prod_{h \neq i,j,k,\ell,m}^{n-1} D_h + \\&+ \sum_{i << \ell}^{n-1} \Delta_{ijkl}(\bar{q}) \prod_{h \neq i,j,k,\ell}^{n-1} D_h + \sum_{i << k}^{n-1} \Delta_{ijk}(\bar{q}) \prod_{h \neq i,j,k}^{n-1} D_h + \\&+ \sum_{i < j}^{n-1} \Delta_{ij}(\bar{q}) \prod_{h \neq i,j}^{n-1} D_h + \sum_i^{n-1} \Delta_i(\bar{q}) \prod_{h \neq i}^{n-1} D_h\end{aligned}$$

and compare with original numerator for

- ▶ local: comparison only for specific cuts
- ▶ global: compare full numerator function at arbitrary  $q$  values

# Numerical stability

- ▶ local/global  $N = N_{\text{rec}}$  test

# Numerical stability

- ▶ local/global  $N = N_{\text{rec}}$  test
- ▶ pole test

# Numerical stability

- ▶ local/global  $N = N_{\text{rec}}$  test
- ▶ pole test
- ▶ power test: check certain combinations of coefficients which should sum to zero if reconstruction was successful  
(e.g. if power of integration momentum is higher than in original function)

# Numerical stability

- ▶ local/global  $N = N_{\text{rec}}$  test
- ▶ pole test
- ▶ power test: check certain combinations of coefficients which should sum to zero if reconstruction was successful  
(e.g. if power of integration momentum is higher than in original function)
- ▶ tensorial reconstruction

# Numerical stability

**Example:** massless fermion loop with two light-like and two massive legs


$$p_{1,2} = (E, 0, 0, \pm E)$$

$$p_{3,4} = (E, 0, \pm Q \sin \theta, \pm Q \cos \theta)$$


$$E = \sqrt{M^2 + Q^2}, \quad p_{3,4}^2 = M^2$$

$$\det G = 32E^4 Q^2 \sin^2 \theta$$

investigate limit  $Q^2 \rightarrow 0$



# Numerical stability



# Tensorial reconstruction

rewrite numerator function as a linear combination of tensors

$$\mathcal{N}(q) = \sum_{r=0}^R C_{\mu_1 \dots \mu_r} q_{\mu_1} \dots q_{\mu_r}$$

$$C_{\mu_1 \dots \mu_r} q_{\mu_1} \dots q_{\mu_r} = \sum_{(i_1, i_2, i_3, i_4) \vdash r} \hat{C}_{i_1 i_2 i_3 i_4}^{(r)} \cdot (q_1)^{i_1} (q_2)^{i_2} (q_3)^{i_3} (q_4)^{i_4}$$

determine the coefficients by sampling  $q$  in a bottom-up approach

**Level 0:**

$$q = (0, 0, 0, 0), \mathcal{N}(0, 0, 0, 0) \equiv \mathcal{N}^{(0)} = C_0$$

**Level 1:** 4 systems, each sampling a monomial depending on one component of  $q$  only

$$\mathcal{N}^{(1)}(q) \equiv \mathcal{N}(q) - \mathcal{N}^{(0)}$$

$$q = (x, 0, 0, 0) \Rightarrow \mathcal{N}^{(1)}(q) \equiv x C_1 + x^2 C_{11} + \dots + x^R \underbrace{C_{11 \dots 1}}_{R \text{ times}}$$


$$q = (0, y, 0, 0) \Rightarrow \mathcal{N}^{(1)}(q) \equiv y C_2 + y^2 C_{22} + \dots + y^R \underbrace{C_{22 \dots 2}}_{R \text{ times}}$$

...

# Tensorial reconstruction

advantages:

- ▶ tensor basis avoids numerical instabilities due to vanishing Gram determinants (as the latter occur in the reduction to a scalar basis)
- ▶ "rescue-system": unstable points will be reprocessed automatically using tensorial decomposition + tensor integrals from `golem95`



# Tensorial reconstruction

further advantage:

- reconstructed tensor integrand can be used as input for the "standard" reduction (more efficient, as kinematic information is already stored in the tensorial coefficients, disentangles part of integrand depending on the loop momenta from dependence on kinematic invariants)
- ⇒ "hybrid method": even for stable phase space points, feeding the reconstructed tensor integrand to the reduction can improve the timings:

# Lines	Time ratio "hybrid" / standard	
N	Rank = 4	Rank = 6
1	1.3	1.6
10	1.1	1.4
100	0.51	0.85
1000	0.30	0.59
10000	0.27	0.55

# Options

reduction:

- ▶ **samurai**, sampling of groups of diagrams
- ▶ **samurai**, sampling of individual diagrams
- ▶ tensorial reconstruction + **samurai**
- ▶ tensor reduction with **golem95**
- ▶ **samurai** + tensor reduction with **golem95** if reconstruction fails
- ▶ tensorial reconstruction with **PJFry** [V.Yundin]

# Options

reduction:

- ▶ **samurai**, sampling of groups of diagrams
- ▶ **samurai**, sampling of individual diagrams
- ▶ tensorial reconstruction + **samurai**
- ▶ tensor reduction with **golem95**
- ▶ **samurai** + tensor reduction with **golem95** if reconstruction fails
- ▶ tensorial reconstruction with **PJFry** [V.Yundin]

scalar integral libraries: optionally

- ▶ **golem95C** Cullen et al., link to **LoopTools** [T.Hahn] possible
- ▶ **QCDLoop** Ellis, Zanderighi
- ▶ **OneLoop** A. van Hameren

# Options

reduction:

- ▶ **samurai**, sampling of groups of diagrams
- ▶ **samurai**, sampling of individual diagrams
- ▶ tensorial reconstruction + **samurai**
- ▶ tensor reduction with **golem95**
- ▶ **samurai** + tensor reduction with **golem95** if reconstruction fails
- ▶ tensorial reconstruction with **PJFry** [V.Yundin]

scalar integral libraries: optionally

- ▶ **golem95C** Cullen et al., link to **LoopTools** [T.Hahn] possible
- ▶ **QCDLoop** Ellis, Zanderighi
- ▶ **OneLOop** A. van Hameren

renormalisation/regularisation schemes:

- ▶ 't Hooft/Veltman
- ▶ DRED (dimensional reduction)
- ▶ CDR (conventional dimensional regularisation)
- ▶ on-shell (mass counter terms for massive quarks)

## Reduction options

```
! Options to control the interoperation between different
! reduction methods
integer :: reduction_interoperation = 2
! 0: use samurai only
! 1: golem95 only
! 2: try samurai first, use golem95 if samurai fails
! 3: tens. reconstruction with golem95, reduction with samurai
! 4: tens. reconstruction with golem95, reduction with samurai,
!     use golem95 if samurai fails
```

## Rational Parts

$$\mathcal{A} = C_4 \text{ (square loop)} + C_3 \text{ (triangle loop)} + C_2 \text{ (circle loop)} + C_1 \text{ (empty circle)} + \mathcal{R}$$

two categories:  $\mathcal{R} = R_1 + R_2$  [Ossola, Papadopoulos, Pittau]

$$N(q) = \hat{N}(\hat{q}) + \tilde{N}(q, \mu^2, \epsilon), \quad q^2 = (\hat{q}^{(4)})^2 - \tilde{q}^2 = \hat{q}^2 - \mu^2$$

$$R_2 = \int \frac{d^D k}{(2\pi)^4} \frac{\tilde{N}(q, \mu^2, \epsilon)}{D_0 \dots D_{n-1}}$$

# Rational Parts

$$\mathcal{A} = C_4 \text{ (square loop)} + C_3 \text{ (triangle loop)} + C_2 \text{ (circle loop)} + C_1 \text{ (empty circle)} + \mathcal{R}$$

two categories:  $\mathcal{R} = R_1 + R_2$  [Ossola, Papadopoulos, Pittau]

$$N(q) = \hat{N}(\hat{q}) + \tilde{N}(q, \mu^2, \epsilon), \quad q^2 = (\hat{q}^{(4)})^2 - \tilde{q}^2 = \hat{q}^2 - \mu^2$$


$$R_2 = \int \frac{d^D k}{(2\pi)^4} \frac{\tilde{N}(q, \mu^2, \epsilon)}{D_0 \dots D_{n-1}}$$

Golem-Samurai offers different options for calculation of  $R_2$

- ▶ **implicit:**  $\mu^2$  terms are kept in the numerator and reduced at runtime
- ▶ **explicit:**  $\mu^2$  terms are reduced analytically
- ▶ **only:** only the  $R_2$  term is kept in the final result
  - (does not require any additional libraries)
- ▶ **off:** all  $\mu^2$  terms are set to zero


# Interface

- ▶ standard interface to real radiation programs  
[\(Binoth Les Houches Accord\)](#) implemented
- ▶ tested with Sherpa and Powheg
- ▶ example  $pp \rightarrow W + \text{jet}$  [figures by G. Luisoni, J. Archibald]



# Interface

- ▶ standard interface to real radiation programs  
[\(Binoth Les Houches Accord\)](#) implemented
- ▶ tested with Sherpa and Powheg
- ▶ example  $pp \rightarrow W + \text{jet}$  [figures by G. Luisoni, J. Archibald]



Example MSSM:  $pp \rightarrow \chi_1^0 \chi_1^0$

NLO SUSY-QCD corrections




figure by G. Cullen, N. Greiner

## Summary and Outlook

- ▶ automation is necessary to have NLO predictions as a standard at LHC

## Summary and Outlook

- ▶ automation is necessary to have NLO predictions as a standard at LHC
- ▶ Golem/Samurai (GoSam) is a powerful and flexible automated tool for one-loop multi-leg amplitudes

## Summary and Outlook

- ▶ automation is necessary to have NLO predictions as a standard at LHC
- ▶ Golem/Samurai (GoSam) is a powerful and flexible automated tool for one-loop multi-leg amplitudes
- ▶ not limited to Standard Model

## Summary and Outlook

- ▶ automation is necessary to have NLO predictions as a standard at LHC
- ▶ Golem/Samurai (GoSam) is a powerful and flexible automated tool for one-loop multi-leg amplitudes
- ▶ not limited to Standard Model
- ▶ numerically robust due to stability test and rescue systems

## Summary and Outlook

- ▶ automation is necessary to have NLO predictions as a standard at LHC
- ▶ Golem/Samurai (GoSam) is a powerful and flexible automated tool for one-loop multi-leg amplitudes
- ▶ not limited to Standard Model
- ▶ numerically robust due to stability test and rescue systems
- ▶ various options to calculate rational term  $\mathcal{R}_2$

# Summary and Outlook

- ▶ automation is necessary to have NLO predictions as a standard at LHC
- ▶ Golem/Samurai (GoSam) is a powerful and flexible automated tool for one-loop multi-leg amplitudes
- ▶ not limited to Standard Model
- ▶ numerically robust due to stability test and rescue systems
- ▶ various options to calculate rational term  $\mathcal{R}_2$
- ▶ standard interface to real radiation programs up and running

## Summary and Outlook

- ▶ automation is necessary to have NLO predictions as a standard at LHC
- ▶ Golem/Samurai (GoSam) is a powerful and flexible automated tool for one-loop multi-leg amplitudes
- ▶ not limited to Standard Model
- ▶ numerically robust due to stability test and rescue systems
- ▶ various options to calculate rational term  $\mathcal{R}_2$
- ▶ standard interface to real radiation programs up and running
- ▶ Samurai is publicly available at  
<http://projects.hepforge.org/samurai/>

# Summary and Outlook

- ▶ automation is necessary to have NLO predictions as a standard at LHC
- ▶ Golem/Samurai (GoSam) is a powerful and flexible automated tool for one-loop multi-leg amplitudes
- ▶ not limited to Standard Model
- ▶ numerically robust due to stability test and rescue systems
- ▶ various options to calculate rational term  $\mathcal{R}_2$
- ▶ standard interface to real radiation programs up and running
- ▶ Samurai is publicly available at  
<http://projects.hepforge.org/samurai/>
- ▶ golem95C (tensor) integral library is available at  
<http://projects.hepforge.org/~golem/95/>

# Summary and Outlook

- ▶ automation is necessary to have NLO predictions as a standard at LHC
- ▶ Golem/Samurai (GoSam) is a powerful and flexible automated tool for one-loop multi-leg amplitudes
- ▶ not limited to Standard Model
- ▶ numerically robust due to stability test and rescue systems
- ▶ various options to calculate rational term  $\mathcal{R}_2$
- ▶ standard interface to real radiation programs up and running
- ▶ Samurai is publicly available at  
<http://projects.hepforge.org/samurai/>
- ▶ golem95C (tensor) integral library is available at  
<http://projects.hepforge.org/~golem/95/>
- ▶ public release of full package planned for this autumn !