

THE OTHER HIGGSES, AT RESONANCE, IN THE LEE- WICK EXTENSION OF THE STANDARD MODEL

**ARXIV:1108.3765, JHEP10 (2011) 145
(IN COLLABORATION WITH ROMAN ZWICKY)**

Dr. Terrance Figy

The University of Manchester

Phenomenology 2012 Symposium

May 8, 2012

OUTLINE

- The Lee-Wick Standard Model
- Higgs boson pair production
- Top quark pair production
- Conclusions

A TOY MODEL

B. Grinstein, D. O'Connell, M.B. Wise (2007)
Based on ideas by Lee and Wick (1969,1970)

(A) HD formalism:

$$\mathcal{L}_{\text{hd}} = \frac{1}{2} \partial_{\mu} \hat{\phi} \partial^{\mu} \hat{\phi} - \frac{1}{2M^2} (\partial^2 \hat{\phi})^2 - \frac{1}{2} m^2 \hat{\phi}^2 - \frac{1}{3!} g \hat{\phi}^3$$

Propagator: $\hat{D}(p) = i(p^2 - p^4/M^2 - m^2)^{-1}$

2 poles: $p^2 = m^2, M^2$

(B) AF formalism: $\hat{\phi} = \phi - \tilde{\phi}$


$$\mathcal{L} = \frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi - \frac{1}{2} \partial_{\mu} \tilde{\phi} \partial^{\mu} \tilde{\phi} + \frac{1}{2} M^2 \tilde{\phi}^2 - \frac{1}{2} m^2 (\phi - \tilde{\phi})^2 - \frac{1}{3!} g (\phi - \tilde{\phi})^3$$

Wrong sign kinetic and mass term M.

The two formulations are equivalent. Use EoM.

A TOY MODEL

B. Grinstein, D. O'Connell, M.B. Wise (2007)



$$D(p) = \frac{i}{p^2 - m^2} \quad ; \quad \tilde{D}(p) = \frac{-i}{p^2 - M^2}$$

$$\begin{aligned} \Sigma(0) &= ig \int \frac{d^4 p}{(2\pi)^4} \frac{i}{p^2 - m^2} - ig \int \frac{d^4 p}{(2\pi)^4} \frac{i}{p^2 - M^2} \\ &= ig \int \frac{d^4 p}{(2\pi)^4} \frac{i(m^2 - M^2)}{(p^2 - m^2)(p^2 - M^2)} \end{aligned}$$

Quadratic divergence is cancelled
leading to a logarithmic
divergence.

A TOY MODEL

B. Grinstein, D. O'Connell, M.B. Wise (2007)

$$\begin{aligned} D_{\tilde{\phi}}(p) &= \frac{-i}{p^2 - M^2} + \frac{-i}{p^2 - M^2} (-i\Sigma(p^2)) \frac{-i}{p^2 - M^2} + \dots \\ &= \frac{-i}{p^2 - M^2 + \Sigma(p^2)}. \end{aligned}$$

$$D_{\tilde{\phi}}(p) = \frac{-i}{p^2 - M^2 - iM\Gamma}, \quad \Gamma = \frac{g^2}{32\pi M} \sqrt{1 - \frac{4m^2}{M^2}}.$$

A LW resonance has a probability Γdt of decaying in the interval $-dt$.

Is this a problem? Shall we debate this issue further or proceed?

LWSM

Higgs Sector (AF formalism)

$$\mathcal{L} = (\hat{D}_\mu H)^\dagger (\hat{D}^\mu H) - (\hat{D}_\mu \tilde{H})^\dagger (\hat{D}^\mu \tilde{H}) + M_H^2 \tilde{H}^\dagger \tilde{H} - V(H - \tilde{H})$$

$$\hat{D}_\mu = \partial_\mu + i(\mathbf{A}_\mu + \tilde{\mathbf{A}}_\mu) \quad \mathbf{A}_\mu = gA_\mu^a T^a + g_2 W_\mu^a T^a + g_1 B_\mu Y$$

$$H^\top = [0, (v + h_0)/\sqrt{2}], \quad \tilde{H}^\top = [\tilde{h}_+, (\tilde{h}_0 + i\tilde{p}_0)/\sqrt{2}]$$

$$\langle h_0 \rangle = v, \quad \langle \tilde{h}_0 \rangle = 0$$

$$\mathcal{L}_{\text{mass}} = -\frac{\lambda}{4} v^2 (h_0 - \tilde{h}_0)^2 + \frac{M_H^2}{2} (\tilde{h}_0 \tilde{h}_0 + \tilde{p}_0 \tilde{p}_0 + 2\tilde{h}_+ \tilde{h}_-)$$

LWSM

Higgs Sector

Symplectic rotation:

$$\begin{pmatrix} h \\ \tilde{h} \end{pmatrix} = \begin{pmatrix} \cosh \phi_h & \sinh \phi_h \\ \sinh \phi_h & \cosh \phi_h \end{pmatrix} \begin{pmatrix} h_{\text{phys}} \\ \tilde{h}_{\text{phys}} \end{pmatrix}$$

Mass eigenvalues:

	h_0	\tilde{h}_0	\tilde{p}_0	h_{\pm}
CP	even	even	odd	none
$\frac{m_{\text{phys}}^2}{M_H^2}$	$\frac{1}{2} \left(1 - \sqrt{1 - 2v^2\lambda/M_H^2} \right)$	$\frac{1}{2} \left(1 + \sqrt{1 - 2v^2\lambda/M_H^2} \right)$	1	1

LWSM

Higgs Sector

Mixing angle:

$$\lambda v^2 = \frac{2m_{h_0,\text{phys}}^2}{(1 + r_{h_0}^2)}, \quad r_{h_0} \equiv \frac{m_{h_0,\text{phys}}}{m_{\tilde{h}_0,\text{phys}}},$$

$$s_H = \cosh \phi_h = \frac{1}{(1 - r_{h_0}^4)^{1/2}},$$

$$s_{H-\tilde{H}} = \cosh \phi_h - \sinh \phi_h = \frac{1 + r_{h_0}^2}{(1 - r_{h_0}^4)^{1/2}}$$

LWSM

Yukawa Interactions (in auxiliary field formalism)

$$\mathcal{L} = \overline{\Psi^t} i \eta_3 \hat{D} \Psi^t - \overline{\Psi_R^t} \mathcal{M}_t \eta_3 \Psi_L^t - \overline{\Psi_L^t} \eta_3 \mathcal{M}^\dagger \Psi_R^t$$

$$\Psi_L^{t\top} = (T_L, \tilde{t}'_L, \tilde{T}_L), \quad \Psi_R^{t\top} = (t_R, \tilde{t}_R, \tilde{T}'_R)$$

SU(2) doublet: $Q_L = (T_L, B_L)^\top$

$$\mathcal{M}_t \eta_3 = \begin{pmatrix} m_t & 0 & -m_t \\ -m_t & -M_u & m_t \\ 0 & 0 & -M_Q \end{pmatrix}, \quad \eta_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

LWSM

Diagonalization of mass matrices

$$\Psi_{L(R),\text{phys}} = \eta_3 S_{L(R)}^\dagger \eta_3 \Psi_{L(R)}, \quad \mathcal{M}_{t,\text{phys}} \eta_3 = S_R^\dagger \mathcal{M}_t \eta_3 S_L,$$

$$S_L \eta_3 S_L^\dagger = \eta_3 \quad \text{and} \quad S_R \eta_3 S_R^\dagger = \eta_3$$

Higgs-quark vertices

$$\mathcal{L} = -\frac{1}{v}(h_0 - \tilde{h}_0) \left(\overline{\Psi}_R^t g_t \Psi_L^t + \overline{\Psi}_L^t g_t^\dagger \Psi_R^t \right) - \frac{1}{v}(-i\tilde{p}_0) \left(\overline{\Psi}_R^t g_t \Psi_L^t - \overline{\Psi}_L^t g_t^\dagger \Psi_R^t \right)$$

$$g_t = \begin{pmatrix} m_t & 0 & -m_t \\ -m_t & 0 & m_t \\ 0 & 0 & 0 \end{pmatrix}, \quad g_{t,\text{phys}} = S_R^\dagger g_t S_L$$

LWSM

LW gauge bosons are massive and mix:

$$\begin{aligned}
 \mathcal{L}_{2g} = & - \frac{1}{2} \text{Tr} \left(B_{\mu\nu} B^{\mu\nu} - \tilde{B}_{\mu\nu} \tilde{B}^{\mu\nu} + W_{\mu\nu} W^{\mu\nu} - \tilde{W}_{\mu\nu} \tilde{W}^{\mu\nu} \right) \\
 & - \frac{1}{2} (M_1^2 \tilde{B}_\mu \tilde{B}^\mu + M_2^2 \tilde{W}_\mu^a \tilde{W}_a^\mu) + \frac{g_2^2 v^2}{8} (W_\mu^{1,2} + \tilde{W}_\mu^{1,2})^2 \\
 & + \frac{v^2}{8} (g_1 B_\mu + g_1 \tilde{B}_\mu + g_2 W_\mu^3 + g_2 \tilde{W}_\mu^3)^2
 \end{aligned}$$



Gauge interactions:

$$\begin{aligned}
 \mathcal{L}_{int} = & - \sum_{\psi=q_L, u_R, d_R} [g_1 \bar{\psi} (\not{B} + \tilde{\not{B}}) \psi + g_2 \bar{\psi} (\not{W} + \tilde{\not{W}}) \psi] \\
 & + \sum_{\psi=q, u, d} \left[g_1 \tilde{\bar{\psi}} (\not{B} + \tilde{\not{B}}) \tilde{\psi} + g_2 \tilde{\bar{\psi}} (\not{W} + \tilde{\not{W}}) \tilde{\psi} \right].
 \end{aligned}$$

LWSM

Couplings to gauges bosons and fermions

E. Alvarez, E. Coluccio, J.Zurita: [arXiv 1004.3496](https://arxiv.org/abs/1004.3496)




$$g_{h_0 f \bar{f}} = -g_{\tilde{h}_0 f \bar{f}} = \cosh \theta - \sinh \theta = \frac{1+r^2}{\sqrt{1-r^4}}, \quad g_{\tilde{P} f \bar{f}} = -1 \quad g_{\tilde{P} gg}^2 = \frac{\sigma(gg \rightarrow \tilde{P})}{\sigma^{SM}(gg \rightarrow H)} = \left| \frac{g_{\tilde{P} t \bar{t}} F_{1/2}^{\tilde{P}}(\beta_{\tilde{P}}^t)}{F_{1/2}(\beta_{\tilde{P}}^t)} \right|^2$$

LWSM: SUMMARY

- For each SM field add a higher derivative (HD) term.
- Auxiliary fields (AF) can be introduced to cast the theory in terms of interactions with mass dimension no greater than 4.
- The AFs are interpreted as LW partner states and have the wrong-sign propagator (aka Pauli-Villars regulators).
- The LWSM solves the hierarchy problem: the extra minus sign in the loop diagrams come from the LW field propagators. No need for opposite spin statistics!
- Unitarity is preserved, provided that the LW fields do not appear as asymptotic states in the S-matrix.
- Causality is preserved at the macroscopic level (where we live). However, there can be violations of causality at the microscopic level.

HIGGS BOSON PAIR PRODUCTION



$$pp \rightarrow h_0 h_0$$



$$\mathcal{M}(gg \rightarrow h_0 h_0) = \frac{1}{32\pi^2} \delta^{ab} \frac{g^2}{v^2} \left(\mathcal{A}_0 P_0 + \mathcal{A}_2 P_2 \right)_{\mu\nu} e(p_1)_a^\mu e(p_2)_b^\nu$$



For details see our Appendix!

HIGGS BOSON PAIR PRODUCTION





Total cross section

HIGGS BOSON PAIR PRODUCTION




Total cross section

HIGGS BOSON PAIR PRODUCTION



Total cross section

HIGGS BOSON PAIR PRODUCTION



$$pp \rightarrow h_0 h_0 \rightarrow b\bar{b}\gamma\gamma$$

HIGGS BOSON PAIR PRODUCTION




$$pp \rightarrow h_0 h_0 \rightarrow b\bar{b}\gamma\gamma$$

- Cut 1: Two isolated photons.
- Cut 2: Two kt jets.
- Cut 3: At least one b-tagged jet.
- Cut 4: $|M_{\gamma\gamma} - m_{h_0}| \leq 2 \text{ GeV}$
- Cut 5: $|M_{bj} - m_{h_0}| \leq 20 \text{ GeV}$
- Cut 6: $|M_{bj\gamma\gamma} - m_{\tilde{h}_0}| \leq \delta m_{\tilde{h}_0}$

Cuts inspired by radion studies performed by ATLAS and CMS.
A more detailed description of cuts is in our paper.

HIGGS BOSON PAIR PRODUCTION

$$pp \rightarrow h_0 h_0 \rightarrow b\bar{b}\gamma\gamma$$



HIGGS BOSON PAIR PRODUCTION

Benchmark	m_{h_0} (GeV)	$m_{\tilde{h}_0}$ (GeV)	$\delta m_{\tilde{h}_0}$ (GeV)
(a)	120	300	40
(b)	130	445	45
(c)	130	550	50

$$pp \rightarrow h_0 h_0 \rightarrow b\bar{b}\gamma\gamma$$

	QCD+EW:	$\gamma\gamma jj$	$\gamma\gamma bb$	$\gamma\gamma cc$	$\gamma\gamma bc$	$\gamma\gamma bj$	$\gamma\gamma cj$
	σ_{gen} (pb)	23.2	0.176	1.56	0.0840	0.519	6.26
	cut 1	0.390	0.370	0.306	0.295	0.344	0.354
	cut 2	0.363	0.358	0.386	0.435	0.406	0.366
	cut 3	0.0526	0.795	0.116	0.516	0.460	0.0920
	cut 4a	0.0212	0.0233	0.0247	0.0217	0.0240	0.0200
	cut 5a	0.249	0.229	0.232	0.242	0.264	0.203
	cut 6a	0.604	0.547	0.713	0.534	0.471	0.627
	ϵ_{tot}	2.37×10^{-5}	3.07×10^{-4}	5.60×10^{-5}	1.85×10^{-4}	1.93×10^{-4}	3.03×10^{-5}
(a)	σ_{eff} (fb)	0.550	0.0527	0.0873	0.0156	0.100	0.190
	cut 4b	0.0150	0.0202	0.0139	0.0167	0.0221	0.0191
	cut 5b	0.221	0.213	0.174	0.242	0.234	0.276
	cut 6b	0.136	0.0567	0.129	0.138	0.165	0.130
	ϵ_{tot}	3.37×10^{-6}	2.56×10^{-5}	6.14×10^{-6}	3.67×10^{-5}	5.46×10^{-5}	8.06×10^{-6}
(b)	σ_{eff} (fb)	0.0782	0.00431	0.00959	0.00309	0.0283	0.0505
	cut 4c	0.0150	0.0213	0.0199	0.0167	0.0221	0.0191
	cut 5c	0.221	0.213	0.174	0.242	0.234	0.274
	cut 6c	0.00723	0.0337	0.00289	0.0164	0.0303	0.00122
	ϵ_{tot}	1.79×10^{-7}	1.52×10^{-5}	1.38×10^{-8}	4.36×10^{-6}	1.00×10^{-5}	7.58×10^{-7}
(c)	σ_{eff} (fb)	0.00414	0.00261	2.15×10^{-5}	0.000366	0.00521	0.00475

HIGGS BOSON PAIR PRODUCTION

Benchmark	m_{h_0} (GeV)	$m_{\tilde{h}_0}$ (GeV)	$\delta m_{\tilde{h}_0}$ (GeV)
(a)	120	300	40
(b)	130	445	45
(c)	130	550	50

$$pp \rightarrow h_0 h_0 \rightarrow b\bar{b}\gamma\gamma$$



$pp \rightarrow h_0 Z \rightarrow \gamma\gamma b\bar{b}$	(a) $m_{h_0} = 120$ GeV, $m_{\tilde{h}_0} = 300$ GeV
σ_{gen} (fb)	32.3
cut 1	0.745
cut 2	0.489
cut 3	0.772
cut 4	0.999
cut 5	0.184
cut 6	0.422
ϵ_{tot}	0.0218
σ_{eff} (fb)	0.703

$pp \rightarrow h_0 h_0 \rightarrow \gamma\gamma b\bar{b}$	(a)	(b)	(c)
σ_{gen} (fb)	11.2	0.964	0.195
cut 1	0.594	0.675	0.693
cut 2	0.414	0.405	0.391
cut 3	0.734	0.760	0.748
cut 4	0.999	0.999	0.999
cut 5	0.601	0.567	0.586
cut 6	0.966	0.823	0.725
ϵ_{tot}	0.105	0.097	0.0861
σ_{eff} (fb)	1.18	0.0935	0.0168

HIGGS BOSON PAIR PRODUCTION

Benchmark	m_{h_0} (GeV)	$m_{\tilde{h}_0}$ (GeV)	$\delta m_{\tilde{h}_0}$ (GeV)
(a)	120	300	40
(b)	130	445	45
(c)	130	550	50

$$pp \rightarrow h_0 h_0 \rightarrow b\bar{b}\gamma\gamma$$



INTERFERENCE EFFECTS IN TOP PAIR PRODUCTION

$$gg \rightarrow R \rightarrow \bar{t}t$$

D.Dicus, A. Strange, and S. Willenbrock

$$\begin{aligned} \frac{d\hat{\sigma}}{ds}(gg \rightarrow \bar{t}t)|_{\text{interference}} &= -|c(s)| \operatorname{Re} \left[\frac{l_{\Delta}}{s - m_R^2 + im_R\Gamma_R} \right] \\ &= -|\tilde{c}(s)| \left((s - m_R^2) \operatorname{Re}[l_{\Delta}] + m_R\Gamma_R \operatorname{Im}[l_{\Delta}] \right) \end{aligned}$$

$$l_{\Delta} = l_{\Delta}(s/4m_t^2) \quad \text{loop triangle function}$$

1. If there is no loop function there will be a peak-dip.
2. For a scalar or pseudo-scalar resonance this pattern does not change.

INTERFERENCE EFFECTS IN TOP PAIR PRODUCTION

$$gg \rightarrow R \rightarrow \bar{t}t$$

$$\begin{aligned} \frac{d\hat{\sigma}}{ds}(gg \rightarrow \bar{t}t)|_{\text{LW-interference}} &= -|c(s)| \operatorname{Re} \left[\frac{-l_{\Delta}(s/4m_t^2)}{(s - m_R^2) - im_R\Gamma_R} \right] \\ &= -|\tilde{c}(s)| \left(-(s - m_R^2) \operatorname{Re}[l_{\Delta}] + m_R\Gamma_R \operatorname{Im}[l_{\Delta}] \right) \end{aligned}$$

Sign-flip in the LW
case





$$\mathcal{M}_R^2 = m_R^2 + \frac{\operatorname{Im}[l_{\Delta}]}{\operatorname{Re}[l_{\Delta}]} m_R\Gamma_R$$

Dip-peak structure

INTERFERENCE EFFECTS IN TOP PAIR PRODUCTION

Peak-dip





Dip-peak



Usual resonance

Lee-Wick type
resonance




Top pair invariant mass spectrum



CONCLUSIONS

- LW Gauge bosons and LW fermions are constrained to be in the few TeV range by EWPO and dilepton searches while the LW Higgs could be below a TeV.
- We have computed the total cross section for double Higgs boson pair production.
- Additionally, we have investigated a search at the a 14 TeV LHC using the di-photon plus di-jet channel. For LW Higgs boson masses of 300 GeV a 5 sigma discovery can be made with 20 1/fb of integrated luminosity.
- We have investigated top pair production in the LW SM. For LW Higgs boson masses above the top pair production threshold, the branching fraction of the LW Higgs boson decaying top pairs dominates. Hence, the top pair channel dominates over the double Higgs boson channel.

Higgs boson decays



Higgs to two photons




Figure 4: *The relative change in the cross-section times decay rate for the full process $gg \rightarrow h_0 \rightarrow \gamma\gamma$ in the LWSM, expressed as $|\kappa_{gg}|^2 |\kappa_{\gamma\gamma}|^2 - 1$, plotted as a function of $m_{h_0,phys}$. Lee-Wick mass scales are such that $M_Q = M_u = m_{\tilde{h},phys} = m_{\tilde{h}^+,phys} = m_{\tilde{W},phys} \equiv \tilde{M}$*

F.Krauss, T.E.J Underwood, R. Zwicky: **arXiv 0709.4054**

References

- 1 - **The Lee-Wick standard model** - Grinstein, Benjamin *et al.* Phys.Rev. D77 (2008) 025012 . arXiv:0704.1845 [hep-ph] . CALT-68-2643, UCSD-PTH-07-04
- 2 - **Negative Metric and the Unitarity of the S Matrix** - Lee, T.D. *et al.* Nucl.Phys. B9 (1969) 209-243
- 3 - **Finite Theory of Quantum Electrodynamics** - Lee, T.D. *et al.* Phys.Rev. D2 (1970) 1033-1048
- 4 - **Causality as an emergent macroscopic phenomenon: The Lee-Wick O(N) model** - Grinstein, Benjamin *et al.* Phys.Rev. D79 (2009) 105019 . arXiv:0805.2156 [hep-th] . CALT-68-2684, UCSD-PTH-08-03
- 5 - **A non-analytic S matrix** - Cutkosky, R.E. *et al.* Nucl.Phys. B12 (1969) 281-300
- 6 - **Vertex Displacements for Acausal Particles: Testing the Lee-Wick Standard Model at the LHC** - Alvarez, Ezequiel *et al.* JHEP 0910 (2009) 023 . arXiv:0908.2446 [hep-ph] . UDEM-GPP-TH-09-183, IFIBA-TH-09-001
- 7 - **Lee-wick Indefinite Metric Quantization: A Functional Integral Approach** - Boulware, David G. *et al.* Nucl.Phys. B233 (1984) 1 . DOE/ER/40048-12 P3
- 8 - **Non-perturbative quantization of phantom and ghost theories: Relating definite and indefinite representations** - van Tonder, Andre Int.J.Mod.Phys. A22 (2007) 2563-2608 . hep-th/0610185
- 9 - **Unitarity, Lorentz invariance and causality in Lee-Wick theories: An Asymptotically safe completion of QED** - van Tonder, Andre . arXiv:0810.1928 [hep-th]
- 10 - **Lee-Wick Theories at High Temperature** - Fornal, Bartosz *et al.* Phys.Lett. B674 (2009) 330-335 . arXiv:0902.1585 [hep-th] . CALT-68-2720, UCSD-PTH-09-02
- 11 - **Massive vector scattering in Lee-Wick gauge theory** - Grinstein, Benjamin *et al.* Phys.Rev. D77 (2008) 065010 . arXiv:0710.5528 [hep-ph] . CALT-68-2662, UCSD-PTH-07-10
- 12 - **Neutrino masses in the Lee-Wick standard model** - Espinosa, Jose Ramon *et al.* Phys.Rev. D77 (2008) 085002 . arXiv:0705.1188 [hep-ph] . CALT-68-2647, IFT-UAM-CSIC-07-21, UCSD-PTH-07-05
- 13 - **One-Loop Renormalization of Lee-Wick Gauge Theory** - Grinstein, Benjamin *et al.* Phys.Rev. D78 (2008) 105005 . arXiv:0801.4034 [hep-ph] . UCSD-PTH-07-11
- 14 - **Ultraviolet Properties of the Higgs Sector in the Lee-Wick Standard Model** - Espinosa, Jose R. *et al.* Phys.Rev. D83 (2011) 075019 . arXiv:1101.5538 [hep-ph]
- 15 - **A Higher-Derivative Lee-Wick Standard Model** - Carone, Christopher D. *et al.* JHEP 0901 (2009) 043 . arXiv:0811.4150 [hep-ph]
- 16 - **Higher-Derivative Lee-Wick Unification** - Carone, Christopher D. Phys.Lett. B677 (2009) 306-310 . arXiv:0904.2359 [hep-ph]
- 17 - **No Lee-Wick Fields out of Gravity** - Rodigast, Andreas *et al.* Phys.Rev. D79 (2009) 125017 . arXiv:0903.3851 [hep-ph] . HU-EP-09-13
- 18 - **A Nonsingular Cosmology with a Scale-Invariant Spectrum of Cosmological Perturbations from Lee-Wick Theory** - Cai, Yi-Fu *et al.* Phys.Rev. D80 (2009) 023511 . arXiv:0810.4677 [hep-th]
- 19 - **Searching for Lee-Wick gauge bosons at the LHC** - Rizzo, Thomas G. JHEP 0706 (2007) 070 . arXiv:0704.3458 [hep-ph] . SLAC-PUB-12481
- 20 - **Unique Identification of Lee-Wick Gauge Bosons at Linear Colliders** - Rizzo, Thomas G. JHEP 0801 (2008) 042 . arXiv:0712.1791 [hep-ph] . SLAC-PUB-13039
- 21 - **Flavor Changing Neutral Currents in the Lee-Wick Standard Model** - Dulaney, Timothy R. *et al.* Phys.Lett. B658 (2008) 230-235 . arXiv:0708.0567 [hep-ph] . CALT-68-2656
- 22 - **Electroweak Precision Data and the Lee-Wick Standard Model** - Underwood, Thomas E.J. *et al.* Phys.Rev. D79 (2009) 035016 . arXiv:0805.3296 [hep-ph] . IPPP-08-21, DCPT-08-42
- 23 - **Custodial Isospin Violation in the Lee-Wick Standard Model** - Chivukula, R.Sekhar *et al.* Phys.Rev. D81 (2010) 095015 . arXiv:1002.0343 [hep-ph] . MSUHEP-100201
- 24 - **The Process $gg \rightarrow h(0) \rightarrow \gamma\gamma$ in the Lee-Wick standard model** - Krauss, F. *et al.* Phys.Rev. D77 (2008) 015012 . arXiv:0709.4054 [hep-ph] . IPPP-07-49, DCPT-07-98
- 25 - **Constraints on the Lee-Wick Higgs Sector** - Carone, Christopher D. *et al.* Phys.Rev. D80 (2009) 055020 . arXiv:0908.0342 [hep-ph]
- 26 - **Higgs \rightarrow Gamma Gamma beyond the Standard Model** - Cacciapaglia, Giacomo *et al.* JHEP 0906 (2009) 054 . arXiv:0901.0927 [hep-ph] . LYCEN-2008-13
- 27 - **Collider Bounds on Lee-Wick Higgs Bosons** - Alvarez, Ezequiel *et al.* Phys.Rev. D83 (2011) 115024 . arXiv:1104.3496 [hep-ph] . ZU-TH-06-11