## Phenomenology 2012 Symposium



Contribution ID: 142

Type: parallel talk

## LHC Discovery Potential for Non-Standard Higgs Bosons in the 3b Channel

Monday 7 May 2012 15:15 (15 minutes)

## Abstract:

In a variety of well motivated models, such as two Higgs Doublet Models (2HDMs) and the Minimal Supersymmetric Standard Model (MSSM), there are neutral Higgs bosons that have significantly enhanced couplings to b-quarks and tau leptons in comparison to those of the SM Higgs. These so called non-standard Higgs bosons could be copiously produced at the LHC in association with b quarks, and subsequently decay into b-quark pairs. However, this production channel suffers from large irreducible QCD backgrounds. We propose a new search strategy for non-standard neutral Higgs bosons at the 7 TeV LHC in the 3b's final state topology. We perform a simulation of the signal and backgrounds, using state of the art tools and methods for different sets of selection cuts, and conclude that neutral Higgs bosons with couplings to b-quarks of about 0.3 or larger, and masses up to 400 GeV, could be seen with a luminosity of 30 fm<sup>2</sup>-1. In the case of the MSSM we also discuss the complementarity between the 3b channel and the inclusive tau pair channel in exploring the supersymmetric parameter space.

Author: Dr MENON, Arjun (University of Oregon)Presenter: Dr MENON, Arjun (University of Oregon)Session Classification: Higgs I