



# PHENO 2012

LHC LIGHTS THE WAY TO NEW PHYSICS




## NEUTRINO COSMOLOGY REDUX



Luis Anchordoqui

# EARLIEST OBSERVATIONALLY VERIFIED LANDMARKS





# OUTLINE

- > Effective number of neutrinos
- > CMB and BBN data/theory predictions
- > Right-handed neutrinos are necessary in...
- > The best of all models:  $U(1)$  for everyone
- > Joint constraints on milliweak interactions (CMB-BBN-LHC)
- > Summary and Conclusions

Work done in collaboration with Haim Goldberg

PRL 108 (2012) 081805



# EFFECTIVE NUMBER OF NEUTRINOS

- > Most straightforward variation of Standard Big-Bang Cosmology
  - ↪ extra energy contributed by new relativistic particles "X"
- > When X's don't share in energy released by  $e^\pm$  annihilation
  - ↪ convenient to account for extra contribution to SM energy density by normalizing it to that of an equivalent neutrino species

$$\rho_X \equiv \Delta N_\nu \rho_\nu = \frac{7}{8} \Delta N_\nu \rho_\gamma \quad (\text{with } \Delta N_\nu = N_\nu - 3)$$

Steigman, Schramm, and Gunn, PLB66 (1977) 202

- > For each additional relativistic degree of freedom:

$$\text{if } T_X = T_\nu \Rightarrow \begin{cases} \Delta N_\nu = 1 & \text{for } X = \text{any two-component fermion} \\ \Delta N_\nu = 4/7 & \text{for } X = \text{scalar} \end{cases}$$

- > If X's have decoupled even earlier and have failed to profit from heating when various other particle-antiparticle pairs annihilated (or unstable particles decayed)

contribution to  $\Delta N_\nu$  from each such particle will be  $\begin{cases} < 1 \\ < 4/7 \end{cases}$



# CMB

> Basic equation:

$$\frac{\Delta N_\nu^{\text{eff}}}{N_\nu^{\text{eff}}} \simeq 2.45 \frac{\Delta(\Omega_m h^2)}{\Omega_m h^2} - 2.45 \frac{\Delta z_{\text{eq}}}{z_{\text{eq}}}$$

- >  $\Delta(\Omega_m h^2)$  from galaxy distributions and precise  $H_0$  measurements  
SDSS Collaboration, MNRAS 401 (2010) 2148      Riess et al., ApJ 699 (2009) 539
- > Wilkinson Microwave Anisotropy Probe  $\rightarrow N_\nu^{\text{eff}} = 4.34_{-0.88}^{+0.86} (2\sigma)$   
WMAP Collaboration, ApJS 192 (2011) 18
- > Atacama Cosmology Telescope  $\rightarrow N_\nu^{\text{eff}} = 4.56 \pm 0.75 (68\% \text{CL})$   
ACT Collaboration, ApJ 739 (2011) 52
- > South Pole Telescope  $\rightarrow N_\nu^{\text{eff}} = 3.86 \pm 0.42 (1\sigma)$   
SPT Collaboration, ApJ 743 (2011) 28
- > Though none of these measurements individually deviates from standard value by more than about  $2\sigma$ 
  - $\rightarrow$  they collectively rule out  $N_\nu = 3$  at approximately 99% CL



# BBN

- Primordial  ${}^4\text{He}$  abundance is driven by decoupling of weak interaction (when neutrinos go out of equilibrium)

$$Y_p \propto e^{-(m_n - m_p)/T_{\text{dec}}}$$

- $T_{\text{dec}}$  determined via  $\Gamma(T_{\text{dec}}) = H(T_{\text{dec}})$

$$T_{\text{dec}}^5 (g/M_W)^4 M_{Pl} \sim \sqrt{N} T_{\text{dec}}^2 \quad (\text{with } M_W \sim 100 \text{ GeV})$$

- For BBN  $\leftarrow T \sim 5 \text{ MeV} \ \& \ N \sim 10$

- $Y_p$  increases with  $N$

- Observationally inferred primordial fractions of baryonic mass in  ${}^4\text{He}$

have been constantly favoring  $N_{\nu}^{\text{eff}} \lesssim 3$

Simha and Steigman, JCAP 06 (2008) 016



# DYNAMICAL DARK MATTER

- > To produce an increase in  $N_{\nu}^{\text{eff}}$  at CMB epoch
  - ↳ but retaining SM value  $N_{\nu}^{\text{eff}}$  for BBN



obvious possibility is production of relativistic particles  
(non-electromagnetically interacting)  
from decay of massive relic particles during/after BBN

- > This has been considered in:
  - > Ichikawa, Kawasaki, Nakayama, Senami, and Takahashi, JCAP 0705 (2007) 008
  - > Fischler and Meyers, PRD 83 (2011) 063520
  - > Hasenkamp, PLB 707 (2012) 121
  - > Menestrina and Scherrer, PRD 85 (2012) 047301
  - > Hooper, Queiroz, and Gnedin, PRD 85 (2012) 063513
- > Required delicate balance can be framed within recently proposed multi-component framework in which dark matter comprises vast ensemble of interacting fields with variety of different masses, mixings, and abundances  
Dienes and Thomas, PRD 85 (2012) 083523



## HOWEVER...

> Unexpectedly  $\rightarrow$  recent determination of primordial  $^4\text{He}$  mass fraction

leads to

$$Y_p = 0.2565 \pm 0.0010(\text{stat}) \pm 0.0050(\text{syst})$$

( $2\sigma$  higher than value given by standard BBN)



For  $\tau_n = 878 \pm 0.8 \text{ s} \rightarrow N_\nu^{\text{eff}} = 3.80_{-0.70}^{+0.80} (2\sigma)$

Izotov and Thuan, ApJ 710 (2010) L67

>  $^4\text{He}$  observed primordial abundance has relative large systematic errors

Aver, Olive, and Skillman, JCAP 1103 (2011) 043

>  $Y_p$  is predicted with precision of  $\sim 0.2\%$

D,  $^3\text{He}$ , and  $^7\text{Li}$  with precisions of roughly 5%, 4% and 8%

**BUT**

because of very precise measurement  $\rightarrow$  constraint on  $N_\nu^{\text{eff}}$

from D/H is competitive with that from  $Y_p$

> Setting aside  $^4\text{He}$  constraints

and combining CMB with BBN theory and observed D/H

$$N_\nu^{\text{eff}} = 3.9 \pm 0.44 (1\sigma)$$

Nollett and Holder, arXiv:1112.2683






## HOW DO WE GET $\Delta N_\nu^{\text{eff}} \sim 1$ WITH RIGHT-HANDED NEUTRINOS?

- Find model in which  $\nu_R$  decouples during quark-hadron transition
- From previous equation  $\rightarrow$  need non-zero coupling to gauge fields  
with  $M \sim \text{TeV}$
- D-brane candidate:  $SU(3)_C \times SU(2)_L \times U(1)_B \times U(1)_L \times U(1)_{I_R}$
- $Y = \frac{1}{2}(B - L) + I_R$
- $B - L$  is non-anomalous if 3  $\nu_L$ 's are accompanied by 3  $\nu_R$ 's
- Matter fields consist of six sets of Weyl fermion-antifermion pairs  
(labeled by index  $i = 1 \dots 6$ )
- Gauging of  $B$  prevents fast proton decay
  - $\rightarrow$  mass via Green-Schwarz/Stueckelberg mechanism
- Gauging of  $L$  disallows heavy Majorana
  - $\rightarrow$  just 3 Dirac neutrinos with tiny Yukawas



# PICTORIAL REPRESENTATION OF D-BRANE CONSTRUCT



Cremades, Ibañez, and Marchesano, JHEP 0307 (2003) 0388



# MODEL PARAMETERS

- > 3 couplings  $g_B, g_L, g_{I_R}$
- > 3 Euler angles  $\rightarrow$  field rotation to coupling diagonal in  $Y$  fixes 2 angles
- > Orthogonal nature of rotation  $\rightarrow$  one constraint on couplings

$$\frac{1}{g_Y^2} = \left(\frac{1}{2g_L}\right)^2 + \left(\frac{1}{6g_B}\right)^2 + \left(\frac{1}{2g_{I_R}}\right)^2$$

- > Baryon number coupling  $g_B$ 
  - fixed to be  $1/\sqrt{6}$  of  $SU(3)$  coupling at  $U(3)$  unification
  - $\rightarrow$  determined elsewhere via RG running
- > 2 remaining d.o.f. allow further rotation leaving
  - $Z'$  to couple to  $B$  at 95%
  - $Z''$  to couple to  $B - L$  at 91%
  - only boson masses are free --



# THE DRAMATIS PERSONAE

Index	Fields	Sector	$SU(3)_C \times SU(2)_L$	$U(1)_B$	$U(1)_L$	$U(1)_{I_R}$	$U(1)_Y$	$g'$	$g''$
1	$U_R$	$3 \rightarrow 1^*$	$(3, 1)$	$\frac{1}{3}$	0	$\frac{1}{2}$	$\frac{2}{3}$	0.368	-0.028
2	$D_R$	$3 \rightarrow 1$	$(3, 1)$	$\frac{1}{3}$	0	$-\frac{1}{2}$	$-\frac{1}{3}$	0.368	-0.209
3	$L_L$	$4 \rightarrow 2$	$(1, 2)$	0	1	0	$-\frac{1}{2}$	0.143	0.143
4	$E_R$	$4 \rightarrow 1$	$(1, 1)$	0	1	$-\frac{1}{2}$	-1	0.142	0.262
5	$Q_L$	$3 \rightarrow 2$	$(3, 2)$	$\frac{1}{3}$	0	0	$\frac{1}{6}$	0.368	-0.119
6	$N_R$	$4 \rightarrow 1^*$	$(1, 1)$	0	1	$\frac{1}{2}$	0	0.143	0.443
-	$H$	$2 \rightarrow 1$	$(1, 2)$	0	0	$\frac{1}{2}$	$\frac{1}{2}$	$2.5 \times 10^{-4}$	0.090

$$\mathcal{L}_{\text{Yukawa}} = -Y_d^{ij} \bar{Q}_i H D_j - Y_u^{ij} \epsilon^{ab} \bar{Q}_{ia} H_b^\dagger U_j - Y_\ell^{ij} \bar{L}_i H E_j + Y_\nu^{ij} \epsilon^{ab} \bar{L}_{ia} H_b^\dagger N_j + \text{h.c.}$$

LAA, Antoniadis, Goldberg, Huang, Lüst, and Taylor, PRD 85 (2012) 086003



# OBTAINING DECOUPLING TEMPERATURE

- Adiabatic reheating of all particles except  $\nu_R$ 's after decoupling gives relation





$$\Delta N_{\nu}^{\text{eff}} = 3 \left( \frac{N(T_{\text{end}})}{N(T_{\text{dec}})} \right)^{4/3}$$

- $T_{\text{end}}$   $\leftarrow$  temperature at end of reheating phase
- $N(T) = r(T)(N_B + \frac{7}{8}N_F)$   $\leftarrow$  effective number of r.d.o.f. at  $T$
- $r(T) = 1$  for lepton/photon and  $r(T) = s(T)/s_{\text{SB}}$  for  $qg$  plasma
- $N(T_{\text{dec}}) = 37 r(T_{\text{dec}}) + 14.25$
- $N(T_{\text{end}}) = 10.75$



# LATTICE QCD

- > Lower  $T$  coincides with most rapid rise of entropy
- >  $N(T)$  based on energy curve rather than entropy-similar



Bazavov et al., PRD 80 (2009) 014504



# QUARK-HADRON CROSSOVER TRANSITION

- > Excess r.d.o.f. within  $1\sigma$  of central value of each if

$$0.46 < \Delta N_{\nu}^{\text{eff}} < 1.08$$

$$\rightarrow 23 < N(T_{\text{dec}}) < 44$$

$$\rightarrow 0.24 < r(T_{\text{dec}}) < 0.80$$

- > From lattice QCD study  $\rightarrow$  this translates to a temperature range

$$175 \text{ MeV} < T_{\text{dec}} < 250 \text{ MeV}$$

Bazavov et al., PRD 80 (2009) 014504

- > Decoupling of  $\nu_R$  occurs when

$$\nu_R \text{ m.f.p.} \geq \text{horizon size} \Rightarrow \Gamma^{\text{int}}(T_{\text{dec}}) = H(T_{\text{dec}})$$



Thermal equilibrium  $\rightarrow$  int = scatt + ann

Chemical equilibrium  $\rightarrow$  int = ann




$$H(T) = 1.66 \langle N(T) \rangle^{1/2} T^2 / M_{\text{Pl}}$$



AS A CHECK ...

behavior of trace anomaly

(which is very sensitive to behavior in crossover region)



shows a sharp peak at 200 MeV

and our range for  $T_{\text{dec}}$  straddles this region

➤ Including  $s \rightarrow 0.18 < r(T) < 0.63$



# CROSS SECTIONS

- > All is fixed except for  $Z'$  and  $Z''$  masses
- > For interaction rate
  - ↳ take average over angles and thermal average over energies

$$\Gamma^{\text{scat}}(T) \simeq 2.0 G_{\text{eff}}^2 T^5$$

$$\Gamma^{\text{ann}}(T) \simeq 0.50 G_{\text{eff}}^2 T^5$$

$$G_{\text{eff}}^2 \sim \sum G_i^2 \quad \text{with} \quad 4 \frac{G_i}{\sqrt{2}} = \frac{g'_6 g'_i}{M_{Z'}^2} + \frac{g''_6 g''_i}{M_{Z''}^2}$$

> By setting in turn  $\Gamma^{\text{ann}}(T) = H(T) \simeq 10.4 T^2 / M_{\text{Pl}}$

and

$$\Gamma^{\text{ann}}(T) + \Gamma^{\text{scatt}}(T) = H(T) \simeq 10.4 T^2 / M_{\text{Pl}}$$

one arrives at two values of  $T_{\text{dec}}$



Chemical equilibrium  $\rightarrow T_{\text{dec}} = 2.75 (G_{\text{eff}}^2 M_{\text{Pl}})^{-1/3}$

Thermal equilibrium  $\rightarrow T_{\text{dec}} = 1.60 (G_{\text{eff}}^2 M_{\text{Pl}})^{-1/3}$

- > When each of these is required to lie between 175 MeV and 250 MeV
  - ↳ allowed regions of  $Z'$  and  $Z''$  masses are defined in each case



# CONSTRAINTS



- Dark shaded areas show region allowed from decoupling requirements to accommodate CMB and BBN data
- Light shaded regions indicate masses excluded by LHC7 dijet searches
- These two estimates should serve to bracket size of actual effect
- Designation of B corresponds to  $Z'$  and  $B - L$  to  $Z''$

LAA and Goldberg, PRL 108 (2012) 081805



# SUMMARY AND CONCLUSIONS

- We developed dynamic explanation of recent hints that relativistic component of energy during BBN and CMB epochs is equivalent to about 1 extra Weyl neutrino
- We work within (string base)  $U(3)_C \times SU(2)_L \times U(1)_R \times U(1)_L$  gauge theory
  - Model endowed with  $3U(1)$  gauge symmetries coupled to  $B, L, I_R$
  - Rotation of gauge fields to basis exactly diagonal in  $Y$  and very nearly diagonal in  $B - L$  and  $B$  fixes all mixing angles and gauge couplings
  - Requiring  $B - L$  current be anomaly free implies existence of 3 right-handed Weyl neutrinos
- Task then reverts to explain why there are not 3 additional r.d.o.f.
- We find that for certain ranges of  $M_B$  and  $M_{B-L}$  decoupling of  $\nu_R$ 's occurs during course of quark-hadron crossover transition
  - ↳ just so that they are only partially reheated compared to  $\nu_L$ 's
- Corresponding upper and lower bounds on gauge field masses yield ranges to be probed at LHC